# STATISTICAL METHODS IN EDUCATION AND PSYCHOLOGY

# PRENTICE-HALL SERIES IN EDUCATIONAL MEASUREMENT, RESEARCH, AND STATISTICS Gene V Glass, Editor



# STATISTICAL METHODS IN EDUCATION AND PSYCHOLOGY

# GENE V GLASS

Laboratory of Educational Research University of Colorado

# JULIAN C. STANLEY

- Departments of Education and Psychology The Johns Hopkins University

PRENTICE-HALL, INC., Englewood Cliffs, New Jersey

### © 1970 by PRENTICE-HALL, INC. ENGLEWOOD CLIFFS, NEW JERSEY

No part of this book may be reproduced in any form or by any means without permission in writing from the publisher.

Current printing (last digit): 10 9 8 7 6 5 4 3 2

13-844928-7

Library of Congress catalog card number: 79-101579

Printed in the United States of America

Prentice-Hall International, Inc., London
Prentice-Hall of Australia, Pty. Ltd., Sydney
Prentice-Hall of Canada, Ltd., Toronto
Prentice-Hall of India Private Ltd., New Delhi
Prentice-Hall of Japan, Inc., Tokyo

# PREFACE

n

This textbook is for use in either a one-semester statistics course or a twosemester sequence of statistics courses in education and the social sciences. Chapters 1 through 8 can be covered thoroughly in a one-semester course meeting three hours weekly, with time left for sampling topics from Chapters 9 and 14. Intensive study of Chapters 10 through 9 in a second course would constitute thorough preparation in the fundamentals of the inferential statistical techniques most useful for research in education, psychology, sociology, and other social sciences.

In writing this text, we sought to produce a more thorough coverage of analysis of variance techniques than now appears in basic statistics texts for social scientists. In addition, an attempt was made to cover correlational techniques comprehensively while emphasizing their interrelationships. We recognize only too clearly the possible shortcomings of this work; but at the same time, we respectfully offer it to the profession with some small measure of pride.

The generosity of publishing companies in allowing the reproduction of scientific material marks them as one of a rapidly dwindling brotherhood of altruistic agents in our society. The publication of this book—and probably

any other contemporary statistics textbook—would not have been possible without the cooperation of several publishers. We are indebted to the following; the Literary Executor of the late Sir Ronald A. Fisher, F.R.S., Dr. Frank Yates, F.R.S., and Oliver & Boyd Ltd., Edinburgh, for their permission to reprint tables from their books Statistical Methods for Research Workers and Statistical Tables for Biological, Agricultural and Medical Research; Biometrika Trustees; Charles Griffin & Company, Ltd.; Cambridge University Press; Chandler Publishing Company; Annals of Mathematical Statistics; McGraw-Hill Book Company; Prentice-Hall, Inc.; The RAND Corporation; The Free Press; The Psychometric Society; The Psychological Corporation; Psychological Bulletin; The National Industrial Conference Board.

During the five years while this text was being written, our colleagues and students contributed in innumerable ways to our efforts. We cannot name them all here, but the following colleagues deserve special thanks for contributing recently to our education in statistical methods: Frank B. Baker. R. Darrell Bock, Raymond O. Collier, Leon J. Gleser, Chester W. Harris, J. Thomas Hastings, John L. Horn, Henry F. Kaiser, William Kruskal, Leonard A. Marascuilo, Leslie D. McLean, Donald L. Meyer, Ellis B. Page. Robert M. Pruzek, Ronald G. Ragsdale, Robert E. Stake, George C. Tiao. and David E. Wiley. Several persons assisted in various ways during their graduate training: Alan Abrams, Donald Bosshart, Glenn Bracht, Russell Chadbourn, James Collins, Ralph Hakstian, Thomas Maguire, Masahito Okada, Perc Peckham, Andrew Porter, Robert Smith, and Peter Taylor. The assistance rendered by four persons merits special mention. Jason Millman and Kenneth D. Hopkins graciously consented to the inclusion of materials which they earlier prepared with the first author for classroom use. Robert Mendro assisted in the preparation of solutions for the problem sets which follow each chapter. Marilyn D. Wang made excellent detailed suggestions for improving much of the book. The task of typing the various drafts of the manuscript was shared by Ann Beadleston, Harriet Clutterbuck. Linda Schmale, and Linda Venter.

GENE V GLASS
JULIAN C. STANLEY

# CONTENTS

# INTRODUCTION (

### 2 MEASUREMENT, SCALES, AND STATISTICS 7

2.1 Measurement, 7 2.2 Measurement Scales, 8 2.3 Variables and Their Measurement, 14 2.4 Symbolizing Data and Operations, 17 2.5 Sigma (£) Notation, 18 Problems and Exercises, 24

# 3 TABULATING AND DEPICTING DATA 26

3.1 Tabulation of Data, 26 3.2 Quantiles, 33 3.3 Determining Percentiles, 34 3.4 Graphing Data, 38 3.5 Representing a Feoquency Distribution Graphically, 42 3.6 Misleading Graphs, 50 3.7 General Suggestions for Constructing Graphs, 22 Problems and Exercises, 54

# MEASURES OF CENTRAL TENDENCY 57

4.1 Introduction, 57 4.2 The Mode, 58 4.3 Conventional Use of the Mode, 58 4.4 The Median, 59 4.5 Calculation of the Median, 59 4.6 The Mean, 60 4.7 Calculation of the Mean, 61 4.9 Mean, Median, and Mode of Combined Groups, 65 4.10 Interpretation of Mode, Median, and Mean, 67 4.11 Choice of a Measure of Central Tendency, 68 4.12 Other Measures of Central Tendency, 71 Problems and Exercises, 22

### 5 MEASURES OF VARIABILITY 75

5.1 Introduction, 75 5.2 The Range, 76 5.3 D, the 90th-to-10th Percentile Range, 77 5.4 The Semi-interquartile Range, 77 5.5 The Variance, 79 5.6 Calculation of the Variance 3, 81 5.7 The Standard Deviation 3, 82 5.8 Some Properties of the Variance, 83 5.9 The Mean Deviation, 85 5.10 Standard Scores, 86 5.11 Skewness, 88 5.12 Kurtosis, 90 Problems and Exercises, 92

# THE NORMAL DISTRIBUTION 95

6.1 Introduction, 95 6.2 History of the Normal Distribution, 95 6.3 The Normal Curve, 98 6.4 The Family of Normal Curves, 100 6.5 The Unit Normal Distribution as a Standard, 102 6.6 Uses of the Normal Curve, 102 6.7 The Bivariate Normal Distribution, 105 Problems and Exercises, 107

# MEASURES OF RELATIONSHIP 109

7.1 Introduction, 109 7.2 The Pearson Product-moment Correlation Coefficient, 109 7.3 A Computational Formula for r<sub>gr</sub>, 113 7.4 Illustration of the Calculation of r<sub>gr</sub>, 114 7.5 Range of Values of r<sub>gr</sub>, 117 7.6 The Effect on r<sub>gr</sub> of Transforming Scores, 119 7.7 Interpreting Correlation Coefficients, 121 7.8 Parther Remarks on the Interpretation of r<sub>gr</sub>, 127 7.9 The Variance of Sums and Differences of Variables, 127 7.10 Other Material on Correlation, 130 Problems and Exercises, 130

# PREDICTION AND ESTIMATION 133

8.1 Preliminaries, 133 8.2 The Problem of Estimating Y from X (or X from Y), 135 8.3 Homoscedasticity and the Standard Error of Estimate, 141 88. Relationships Between  $b_0$  and  $b_1$  and Other Descriptive Statistics, 144 8.5 Verification that the Least-squares Criterion is Satisfied by  $b_1$  and  $b_0$ , 149 8.6 Measuring Nonlinear Relationships Between Variables; the Correlation Ratio  $\eta^2$ , 150 8.7 Additional Reading on Prediction, 152 Problems and Exercises, 152

### 9 ADDITIONAL MEASURES OF RELATIONSHIP 155

9.1 Introduction, 155 9.2 An Overview of the Chapter, 156 9.3 Measures of Relationship, 158 9.4 Part Correlation and Partial Correlation, 182 9.5 Multinle Correlation and Prediction, 186 Problems and Exercises, 191

# PROBABILITY 195

10.1 Introduction, 195 10.2 Probability as a Mathematical System, 196 10.3 Combining Probabilities, 198 10.4 Permutations and Combinations, 203 10.6 Indomness and Random Sampling, 212 10.7 Random Variable, 215 10.8 Types of Random Variables, 217 10.9 Probability as an Area, 218 10.10 Expectations and Moments, 220 Problems and Exercise, 226

### II THEORETICAL DISTRIBUTIONS FOR USE IN STATISTICAL INFERENCE 228

11.1 Introduction, 228 11.2 Normal Distribution, 228 11.3 Chi-square Distributions, 229 11.4 F-distributions, 233 11.5 t-distributions, 234 11.6 Re-lationships among the Normal, t-, Chi-square, and F-distributions, 236 Problems and Exercise, 238

### 12 STATISTICAL INFERENCE: ESTIMATION 240

12.1 Populations and Samples; Parameters and Statistics, 240 12.2 Random Sampling, 242 12.3 The Concept of a Sampling Distribution, 243 12.4 Properties of Estimators, 250 12.5 Interval Estimation, 256 12.6 Conclusion, 268 Problems and Exercises, 269

# STATISTICAL INFERENCE: HYPOTHESIS TESTING 271

13.1 Introduction, 271 13.2 Scientific and Statistical Hypotheses, 272 13.3 Testing a Statistical Hypothesis, 274 134 Typs 1 Error, Level of Significance, and Critical Region, 279 13.5 Type II Error, B, and Power, 283 13.6 Non-directional and Directional Alternatives: "Two-tailed vs. One-tailed Tests," 288 137 Postscript, 250 Problems and Exercise, 250

# 14 SELECTED INFERENTIAL TECHNIQUES 292

14.1 Introduction, 292 14.2 Inferences about the Mean,  $\mu$ , of a Population, 293 14.3 Inferences about  $\mu_1 = \mu_2$  Using Independent Samples, 295 14.4 Inferences

about  $p_1 - p_2$  Using Dependent Samples, 297 14.5 Inferences about the Variance,  $\sigma_1^2$ , of a Population, 301 14.6 Inferences about  $\sigma_1^2/\sigma_2^2$  Using Independent Samples, 303 14.7 Inferences about  $\sigma_1^2/\sigma_2^2$  Using Dependent Samples, 306 14.8 Inferences about  $\rho_1$ , the Population Product-moment Correlation Coefficient, 308 14.9 Inferences about  $\rho_2$ ,  $\rho_2$  Using Independent Samples, 311 14.10 Inferences about  $\rho_3$ ,  $\rho_4$ ,  $\rho_2$  Using Dependent Samples, 311 14.10 Inferences about  $\rho_3$ ,  $\rho_4$ ,  $\rho_2$ , Using Dependent Samples, 313 14.11 Inferences about Other Correlation Coefficients, 315 14.12 Inferences about  $\rho_1$ ,  $\rho_2$  Using Independent Samples, 316 14.15 Inferences about  $\rho_3$  Dependent Samples, 326 14.15 Inferences about  $\rho_1$ ,  $\rho_2$  Using Dependent Samples, 326 14.15 Inferences About the Independence of Classifications in a Contingency Table, 329 14.16 Relationship between Interval Estimation and Hypothesis Testing, 333 Problems and Exercises, 334

# 15 THE ONE-FACTOR ANALYSIS OF VARIANCE—FIXED EFFECTS 238

15.1 Layout of Data, 338 15.2 A Model for the Data, 339 15.3 Estimates of the Terms in the Model, 342 15.4 Sums of Squares, 343 15.5 Restatement of the Null Hypothesis in Terms of Population Means, 345 15.6 Degrees of Freedom, 345 15.7 Mean Squares, 347 15.8 Expectations of MS, and MS<sub>20</sub>, 347 15.9 Some Distribution Theory, 350 15.10 The F-test of the Null Hypothesis: Rationale and Procedure, 353 15.11 The One-way ANOVA with n Observations per Cell (Summary), 357 15.12 The One-way ANOVA with Unequal n's, 362 15.13 Consequences of Failure to Meet the ANOVA Assumptions: the "Robustness" of ANOVA, 368 15.14 Testing Homogeneity of Variances, 374 15.15 Power of the F-test, 376 Problems and Exercises, 377

# 16 MULTIPLE COMPARISON PROCEDURES 381

16.1 Introduction, 381 16.2 The T-method, 383 16.3 Confidence Intervals around Contrasts by the T-method, 385 16.4 The S-method, 388 16.5 Confidence Intervals around Contrasts by the S-method, 393 16.6 The T- and S-methods Compared, 395 Problems and Exercises, 397

### IT THE TWO-FACTOR ANALYSIS OF VARIANCE—FIXED EFFECTS 400

17.1 The Layout and Symbolization of Data, 400 17.2 A Model for the Data, 402 17.3 Least-squares Estimation of the Model, 403 17.4 The Nature of Interaction, 406 17.5 Statement of Null Hypotheses, 411 17.6 Sums of Squares in the Two-factor ANOVA, 4/4 17.7 Degrees of Freedom, 4/6 17.8 Mean Squares, 48 17.9 Computational Procedures, 4/8 17.10 Expected Values of Mean Squares, 421 17.11 The Distributions of the Mean Squares, 424 17.12 The Distributions of the Mean Squares, 424 17.12 Hypothesis Tests of the Null Hypothesis, 428 17.13 Review of the Two-factor ANOVA with Unequal n's, 432 17.15 Multiple Comparisons in the Two-factor ANOVA, 443 Problems and Exercises, 446

### IB ONE-FACTOR AND MULTIFACTOR ANALYSIS OF VARIANCE: RANDOM, MIXED, AND FIXED EFFECTS 452

18.1 Introduction, 452 18.2 The Random-effects ANOVA Model, 452 18.3 The Mixed effects ANOVA Model, 463 18.4 Rules of Thumb for Writing the ANOVA Table, 471 18.5 Definitions of Terms, 473 18.6 Determining the Possible Lines (Sources of Variation) of the ANOVA Table, 474 18.7 Determining the Degree of Freedom for Sources of Variation, 473 18.8 Computing Sums of Squares, 476 18.9 Determining the Expectations of Mean Squares, 479 18.10 Significance Testing, 481 Problems and Exercises, 482

# 19 FUNDAMENTALS OF EXPERIMENTAL DESIGN 486

19.1 Introduction, 486 19.2 An Example, 488 19.3 Advantages and Disadvantages of the Factorial Design, 490 19.4 Blocking, 49 19.5 Ordered levels of Factors, 495 19.6 Random Selection of Factor Levels, 495 19.7 Natural and Controlled Experiments, 495 19.8 Other Experimental Designs, 498 19.9 The Analysis of Covariance, 499 19.10 Quasi-experimentalion, 501 19.11 The Experimental Unit and the Unit of Statistical Analysis: Comparative Experiments with Intact Groups, 501

APPENDIX A: STATISTICAL TABLES 510

APPENDIX B: ANSWERS TO PROBLEMS AND EXERCISES 555

APPENDIX C: VERIFICATION OF STANDARD SOLUTION TO LEAST-SQUARES CRITERION 570

BIBLIOGRAPHY 572

**AUTHOR INDEX 586** 

SUBJECT INDEX 589

# INTRODUCTION

Popular attitudes toward statistics contain a mixture of awe, cynicism, suspicion, and contempt. Statisticians have been uncomplimentarily placed in the company of liars and accused of "statisticulation"—the art of lying with statistics while maintaining an appearance of objectivity and rationality. Someone once remarked that "If all the statisticians in the world were laid end to end—it would be a good thing." A statistican has been scornfully depicted as a man who drowns while wading in a river of average depth 3 feet or who sits with his head in a refrigerator and his feet in an oven and reports that "On the average, I feel fine." In a weekly news magazine, the author of an essay on "The Science and Snares of Statistics" concluded that "... perhaps the time has come for society to be less numerically conscious and therefore less willing to be ruled by statistics."

Persons beginning the study of statistics will profit from discarding the popular image of statistics and statisticians. They should realize that non-popular image of statistics are statistically as it can be expressed quantitatively. See the property of the property of

Time, September 8, 1967, p. 29.

CHAP. 1

acceptance of verbal nonsense, and a knowledge of statistics is the best defense against quantitative nonsense. The first step toward replacing popular images of statistics with more realistic ones is the study of the structure of the discipline of "statistical methods" and its historical antecedents.

There were two widely divergent influences on the early development of statistical methods. Statistics had a mother who was dedicated to keeping orderly records of governmental units (state and statistics come from the same Latin root, status) and a gentlemanly gambling father who relied on mathematics to increase his skill at playing the odds in games of chance. From the mother sprang counting, measuring, describing, tabulating, ordering and the taking of censuses—all of which led to modern descriptive statistics. From the adventurous, intellectual father eventually came modern inferential statistics, which is based squarely on theories of probability. A recent addition called the design of experiments relies heavily on a combination of probability theory and rather elementary but uncommon logic. This text offers an introduction to and a useful explanation of descriptive statistics, inferential statistics, and the design of experiments. Chapters 2 through 9 cover a large portion of descriptive statistics. Beginning with "Probability" in Chapter 10 and extending through Chapter 14, several tonics from inferential statistics are covered. Chapters 15 through 19 present the considerations and inferential techniques fundamental to the design and analysis of experiments.

Descriptive statistics involves the tabulating, depicting, and describing of collections of data. These data may be either quantitative, such as measures of height and weight, or qualitative, such as sex and personality type. Large masses of data must generally undergo a process of summarization or reduction before they are interpretable by the human mind. A monkey is unsuccessful in his clumsy attempt to untie a simple knot because the complexity of the problem of untying the knot surpasses the resolving power of the poor creature's intellect. The unsuccessful fumbling attempt of a fisherman to unsnarl a backlash in his fishing reel is analogous to the monkey's plight. For the fisherman, that backlash is a Gordian knot; it presents too great a problem for his finite intellect. Similarly, but at a different level, the human mind cannot extract the full import of a mass of data (How do they vary? About how large are they? Is one set useless in reducing uncertainty about the other?) without the aid of special techniques (swords to cut the Gordian knot). Thus descriptive statistics serves as a tool to describe or summarize or reduce to manageable form the properties of a mass of data.

Inferential statistics is a formalized body of techniques for solving another class of problems that present great difficulties for the unaided human mind. This general class of problems characteristically involves attempts to infer

the properties of a large collection of data from inspection of a sample of the collection. For example, a school nurse wishes to determine the proportion of children in the fifth grades of a large school system who have never had chicken pox. It would be unnecessary to question each child if the proportion could be reliably estimated from a sample of as few as 100 children. But how does the proportion of children in the sample of 100 who haven't had chicken pox relate to the analogous proportion in the entire fifth-grade population? The answer can be obtained through inferential statistics. Thus the purpose of inferential statistics is to surmise the properties of a population from a knowledge of the properties of only a sample of the population. Inferential statistics builds upon descriptive statistics. The inferences are drawn from particular properties of samples to particular properties of populations; the descriptions of the properties of both thesamples and the populations are obtained by methods of descriptive statistics.

The design and analysis of experiments is a third important branch of statistical methods. These methods were developed for the discovery and confirmation of causal relationships among variables. Researchers in the social sciences are concerned with causation, a very complex concept in philosophy. Experimental design is so important for the study of causal relationships that in some philosophical systems an experiment constitutes an operational definition of a causal relationship. Adults make causal inferences during all of their waking moments. The frequent use of the word "because" reveals this: "The school bond failed to pass because it was not well publicized," or "He scored poorly on the intelligence test because he was overly annious about the consequences of the score."

The sentence, "Drug A kills pain faster than Drug B," does not contain the word "because," but it implies that "More of this group of patients than of that group gained fast relief from pain because Drug A was administered to the former, whereas Drug B was administered to the latter." The weakness of the "because" explanation is its potential vagueness. This weakness is betrayed by the favorite remark of many young children when, at the prelogical stages of their thinking, they are confronted with evidence of their misbehavior. If asked, "Why did you do that?" they respond, "Just 'cause." Obviously, the word has many denotations and connotations.

Statistical methods assist researchers in describing data, in drawing inferences to larger bodies of data, and in studying causal relationships. They can be a useful tool in answering such questions as the following: How old is the average man on the day he receives a Bachelor of Arts degree from a certain college? What percentage of these new graduates have blue eyes? What percentage of them are presently married? How many of them already have 0, 1, 2, . . . children? Do those who earned good grades as undergraduates attend graduate schools in greater proportions than those who earned mediocre grades? Has the international situation affected

INTRODUCTION CHAP. 1

attendance at graduate schools? In an experiment, will college students who are received with friendliness by a group conform more to that group's judgments than will college students shunned by a group? Is this differential reaction, if found, contingent on the sex of the student? For instance, are women more or less amenable than men to group influence?

Mastering statistical methods requires some mathematical skill. The subject, statistics, is a branch of applied mathematics. Statistics is inadequately described in the dictionary as "the science of compiling facts." If statistics were only that simple, this text would be short indeed. In its more rigorous form, statistics is usually called mathematical statistics. For social scientists and other nonmathematicians it is termed "applied statistics" and includes much use of intuition, simple arithmetic, and elementary algebra. To study mathematical statistics seriously, one needs a background including at least advanced calculus and matrix theory; however, much of the rationale of applied statistics and many of its techniques can be learned without this mathematical maturity, although of course not as deenly understood. Perhaps this is partly the reason why various social sciences tend to be technique oriented. In large universities, separate courses in "educational and psychological statistics," "sociological statistics," "economic statistics," and the like can usually be found outside the Department of Statistics. Fortunately, however, the most fundamental principles are useful in neatly all disciplines, from agriculture to zoology. A knowledge of statistics is becoming necessary for the pursuit of a career of scholarship in any empirical discipline. Many graduate schools have recently acknowledged its importance by accepting course work in statistics as a replacement for one of the two foreign language courses that are traditionally required for the PhD degree. The substitution is apt; statistics is an increasingly important means of communicating knowledge. The increasing recognition of statistics as a tool of scholarship brings to mind the description of the education of children in B. F. Skinner's utopian community, Walden Two: "We help them in every way short of teaching them. We give them new techniques of acquiring knowledge and thinking . . . We give them an excellent survey of the methods and techniques of thinking, taken from logic, statistics, scientific method, psychology, and mathematics. That's all the 'college education' they need. They get the rest by themselves in our libraries and laboratories."

The word statistic is defined by Kendall and Buckland (1957)† as "A summary value calculated from a sample of observations, usually but not necessarily as an estimator of some population parameter; a function of

<sup>\*</sup> Skinner, B. F. Walden Two. New York: The Macmillan Company (paperback edition), 1962, p. 121.

<sup>†</sup> For complete information on this and later references, consult the bibliography at the end of this text.

CHAP. 1 INTRODUCTION

sample values." The contrasting term is "parameter," which we shall define later. Thus, the arithmetic mean of the numbers 1, 4, and 4, which is 3, is a statistic. The fact that a certain man has 2 children is a datum, whereas the average number of children in a town is a statistic. (You can actually see those 2 children, but not the average child.) This distinction between "statistic" and "datum" is not always preserved, however. Some applied statisticians and researchers use "statistic" to cover both, even saying that a person's name or hair color is a statistic.

Individual statistical techniques are unified by an underlying method. We shall attempt to show this unity and the interrelations as clearly as possible by using only the elementary mathematics that those studying this book will normally have learned in secondary school. Some special symbols will be introduced as needed; these will be explained carefully. They must be mastered at the point of introduction because thinking in statistics is facilitated by such symbols.

The approach of this text is from descriptive to inferential, with the statistics and logic of controlled experimentation gradually introduced. Two goals toward which this textbook is directed are (1) ability to read reports of surveys, studies, investigations, and experiments in your substantive field with moderate competence (given that you understand the substantive problems being researched) and (2) appreciable ability to plan your own studies and analyze data resulting from them.

The amount you learn will depend on your quantitative aptitude and distance, plus the efforts of your instructor. The range of acquistion of statistics during an academic year is usually great for the students in a given class. Some develop considerable expertise, and others build a sound basis for further study in class and out. A few students, especially those traumatized by "symbol shock," find the pace too fast and the explanations too scanty. If they are to avoid a life of stuttering every time they even try to pronounce "statistics," their best hope is a skilled tutor to ease their anxiety, help them develop arithmetic and symbolic competence, and promote overlearning of the basic aspects needed for subsequent sections of the book. The sooner such a tutor is acquired the better.

If you have not studied mathematics, logic, or any other rigorous and deductive body of knowledge for some time, you may find studying statistics uncomfortable for a while. In many disciplines characterized by vague verbal discourse and personalistic use of language, a student can sustain sloppy and erroneous thinking for long periods without being confronted with its inadequacy. It is simply impossible to refute the statement that "We must educate the whole child" or that "Both heredity and environment are important in determining human intelligence." However, the student of statistics is likely to be confronted abruptly and uncomfortably with the results of loose thinking, as when a calculated quantity that cannot possibly

INTRODUCTION CHAP, 1

be negative resists one's best attempts to make it come out positive. If you are inclined toward critical and precise thought, this restrictive and confining mantle will soon begin to feel comfortable. The satisfying reassurance of knowing that you are mastering a logical and unambiguous language will outweigh the occasional pang of anxiety produced when you are discovered speaking illogically. Being openly and clearly wrong is the price we must pay for knowing when we are correct. Never to know whether you are speaking sense or nonsense is too expensive a luxury to enjoy in an age when sense is precious and nonsense is rampant.

It is easy to succumb to the delusion that you are learning a great deal about statistics from simply reading the text. A statistics text is not a novel; none will ever become a "Book-of-the-Month Club" selection. This statistics text must be studied carefully and thoughtfully. Above all, the exercises and problems that follow each chapter must not be slighted. Working these exercises will put a fine edge on your knowledge of the subject. Skip the exercises and you may never know what you don't know about statistics. In statistics, as in most human endeavors, "A little learning is a dangerous thing; drink deep, or taste not the Pierian Spring...."

Almost anyone can learn a great deal about statistics. Even many high-school students find it interesting, so do not think it accessible only to the specially appointed. A good textbook helps; in the hands of a gifted teacher it is doubly effective. While writing this book we have held in mind three functions of a textbook: it must be an effective pedagogic instrument; it must serve as a reference work once the material is learned; it must serve to direct the student toward the larger body of knowledge that it only samples. Clearly the first function is primary, for there is no point in referring to what is not known nor should one attempt to read more advanced material until the fundamentals have been mastered. Master the fundamentals on and use them the rest of your life.

Alexander Pope, Essay on Criticism (1711).

2

# MEASUREMENT. SCALES. AND STATISTICS

# 2.1 MEASUREMENT

Everyone chooses to define "measurement" in slightly different terms from slightly different points of view. What is common to all definitions seems to be this: Measurement is the assigning of numbers to things according to rules. To measure a person's height is to assign a number to the distance between the top of a person's head and the bottom of his feet with the use of a ruler. Measurement of a child's IQ is the assignment of a number to the pattern of response that he makes to a group of standard problems. Measurement transforms certain attributes of our perceptions into familiar, tractable things called "numbers." What an impossible world it would be if we did not measure! How useful to a physicist would be the knowledge that iron melts at a high temperature, or to a traveler that Chicago is "down the road a piece"? A little reflection will reveal the important role played by measurement in education and almost every other social enterprise. 7

### 2.2 MEASUREMENT SCALES

The ideas of "scales of measurement" constitute a useful set of concepts. Behavioral scientists, and very few other scientists, have been concerned with these problems. We shall now discuss briefly the different scales and their implications for statistics.

## Nominal Measurement\*

Nominal measurement (giving a name or names) scarcely deserves to be called "measurement." It is the process of grouping objects into classes so that all of those in a single class are equivalent (or nearly so) with respect to some attribute or property. The classes are then given names: that the classes could as well receive, and often do receive, numerals for identification instead of names may account for the title "nominal measurement." Classificatory schemes in biology are examples of nominal measurement. Psychologists often code "sex" by assigning 0 to "female" and 1 to "male": this is nominal measurement, too. We would perform nominal measurement if we assigned I to Englishmen, 2 to Germans, and 3 to Frenchmen. Does one Englishman plus one German equal one Frenchman (1 + 2 = 3)? Obviously not. The numerals we assign in nominal measurement have all the properties of any other numerals. We can add them, subtract them, divide them, or simply see which is larger than another. But if the process by which we assigned the numerals to objects was nominal measurement. then our playing with the size, order, and other properties of the numerals will imply nothing at all about the objects themselves because we took no cognizance of the size, order, and other properties of the numerals when we assigned them. When measurement is nominal only, one uses only the property of numbers that I is distinct from 2 or 4 and that if object A has a I and object B a 4, then A and B are different with respect to the attribute measured. It does not necessarily follow that B has any more of the attribute than A. The three remaining scales of measurement we shall encounter make use of three additional properties of numbers: numbers can be ordered with respect to size; they can be added; and they can be divided.

### Ordinal Measurement

Ordinal measurement is possible when the measurer can detect differing degrees of an attribute or property in objects. This being possible, he makes

<sup>\*</sup> The names of the scales of measurement used here and many of the concepts are due to S. S. Stevens (1951)

use of the "orderedness" property of numbers and assigns numbers to objects so that if the number assigned to object A is larger than the number assigned to B, then A possesses more of the property in question than B.

Suppose we ask a person to rank-order Mary, Jane, Alice, and Betty from least beautiful to most beautiful. We may rank them as follows: Betty, Jane, Mary, Alice. Ordinal measurement occurs when we assign the numbers 1, 2, 3, and 4 to Betty, Jane, Mary, and Alice respectively. Notice that the numbers 0, 23, 49, and 50 would have served just as well since the distance between any two adjacent numbers is of no significance. We may not feel that the measurer was capable of discerning, for example, whether the difference between the amount of beauty possessed by Betty and Jane is greater or less than the difference in beauty between Jane and Mary. Hence, no significance should be attached to the fact that the difference between Betty's and Jane's scores is the same as the distance between Mary's and

Alice's scores.

Notice how the numbers take the place of the objects of concern. The numbers are a partial representation of the objects; we agree to treat the numbers as if both the facts that they are distinct and they can be ordered numbers as if both the facts that they are distinct and they can be ordered are important. At the ordinal measurement stage, the numbers constitute something of a reduction in effort to convey information. Instead of having to report that "Betty was judged least beautiful, Jane next least, Mary second most beautiful, and Alice most beautiful," we can report simply:

Person	Score
Mary	2
Jane	4
Alice	i
Betty	,

A scale of hardness of minerals is an ordinal scale. If mineral A can scratch mineral B, it is harder; hence, it receives a higher number. Suppose minerals A, B, C, and D have been assigned respectively numbers 12, 10, 8, and  $\delta$  in this manner. We know the hardest and the softest mineral. Is the difference between the hardness of A and B the same or different from the difference between the hardness of C and D? We have no way of knowing because the numbers were assigned so that only their uniqueness and ordinal properties were regarded—measurement was ordinal.

properties were regarded—measurement was ordinat.

Another common ordinal scale is "rank in high school class." Numbers are assigned from 1 for "highest grade-point average" to n for "lowest grade-point average" in a class of n students. (If, for example, the three top students all had perfect averages, each would receive a rank of 2, which is students all had perfect averages, each would receive a rank of resolving the average of the first three ranks, 1, 2, and 3. This means of resolving

10

ties is conventional, because it keeps the sum of the tied and untied ranks the same: 1+2+3=2+2+2.)

There is no law preventing one from adding, subtracting, multiplying, etc. numbers that have been assigned to objects by ordinal measurement. However, the results of these operations may reflect nothing about the amounts of the property in question that the objects corresponding to the numbers possess. For example, the difference between the "beauty socres" of Alice and Betty is 3; the difference between the socres of Mary and Jane is 1. Does this mean that the difference in beauty between Alice and Betty is three times as great as the difference in beauty between Mary and Jane? Of course it doesn't. The results of the arithmetic cannot be interpreted as saying anything about the amounts of the property actually possessed by the objects. You can do what you wish with the numbers you obtain, but you are always faced with the question, "Do the results of these operations have any meaning for me?"

### Interval Measurement

Interval measurement is possible when the measurer can distinguish not only between different amounts of the property in objects (the characteristic of ordinal measurement) but can also discern equal differences between objects. For interval measurement a unit of measurement (degree, inch, foot, ounce, etc.) has been defined. A number is assigned to an object that equals the number of units of measurement equivalent to the amount of the property possessed. For example, the temperature of a certain metal bar is 86° centigrade. An important feature which distinguishes interval measurement from ratio measurement (to be studied next) is that an object with a measurement of zero does not necessarily lack the attribute being measured. Hence, water at 0°C is not absolutely without temperature. The zero point on an interval scale is an arbitrary one.

The numbers assigned in the process of interval measurement have the properties of distinctness and order, but in addition the difference between the numbers is meaningful. The number assigned to the object is the number of units of measurement it has. Today's temperature is 60°F; yesterday's temperature was 55°F. Today is 5° warmer than yesterday. If tomorrow's temperature is 70°F then we know that yesterday and today are more alike in temperature than are today and tomorrow. The difference between 55 and 60 is half as large as the difference between 60 and 70; furthermore, the sizes of these differences tell us something about the temperature of the air.

The numbering of the years is an interval scale. The year I was arbitrarily set originally as the year of the birth of Christ. The unit of measure-

11 MEASUREMENT SCALES SEC. 2.2

ment is a span of 3651 days. The year 1931 is more recent than any other year with a smaller number. Finally, the time between 1776 and 1780 equals the length of time between 1920 and 1924. James K. Polk was president for half as long (1845-1849) as Dwight D. Eisenhower (1953-1961).

nor nan as iong (1843–1849) as Dwight D. Eisenhower (1953–1961).

Interval measurement involves assigning numbers to objects in such a way that equal differences in the numbers correspond to equal differences in the amounts in the objects of the property or attribute measured. The zero point of the interval scale can be placed arbitrarily and does not indicate absence of the property measured.

# Ratio Measurement

Ratio measurement differs from interval measurement only in that the zero point is not arbitrary but indicates total absence of the property measured. point is not arourary out indicates total absence of the property measured. The measurer can perceive the absence of the property, and he has a unit of measurement with which he records differing amounts of the property. of incasurement with which he records different amounts of the property.

Equal differences between the numbers assigned in measurement reflect equal differences in the amount of the property possessed by the things measured. Furthermore, since the zero point is not arbitrary but absolute, it is meaningful to say that A has two, three, or four times as much of the property as B. o say that A has two, three, or roat times as much of the property as B.
Height and weight are examples of ratio measurement scales. Zero

height is no height at all, and a man six-feet tall is twice as tall as a three-foot boy. The ratio scale is so named because the ratios of numbers on a ratio tooy. The ratio scale is so manifed occasion the factors of numbers on a ratio scale are meaningful. These ratios can be interpreted as ratios of amounts of the objects measured. A ratio statement about a strictly interval scale has no meaning in terms of the amounts of the attribute in the objects. nas no meaning in terms of the amounts of the attroute in the objects. For example, if June 3 had a high temperature of 90°F and March 17 had a high of 45°F, it is not correct to say that June 3 had twice as much temperature as March 17.

e as march 11. Most measurement in educational research and in the behavioral sciences occurs at the nominal, ordinal, and interval levels. Few important variables occurs at the nominat, occurs, and their revers. Few important variables in these fields as yet lend themselves to ratio measurement; in fact, one in these nears as yet rend themselves to take measurement; in fact, one must search diligently to find scales of measurement that will satisfy the must search unigently to the scales of measurement that will satisfy the conditions of an interval scale. Occasionally, ratio-scale variables such as time (to solve a problem or learn a list of words), height, weight, or distance time (to solve a problem of learn a use of words), neight, weight, or distance will be of interest, but such occasions arise infrequently. You must undertake will be of interest, but such occasions arise intrequently. I ou must undertake to recognize measurement at the nominal and ordinal levels and prepare to recognize measurement at the nominal and ordinal revers and prepare for the problems that the analysis and interpretation of such data present. Table 2.1 summarizes and supplements what has been said thus far

about scales of measurement.

Ratio

TABLE 2.1 SUMMARY OF CHARACTERISTICS AND EXAMPLES OF MEASUREMENT SCALES
Scale Characteristics Examples

Nominal Objects are classified and classes are denoted by numbers. That the number for one class is greater or less than another number reflects nothing about the properties of the objects other than that they are different.

Racial origin, eye color, numbers on football jerseys, sex, clinical diagnoses, automobile license numbers, social security numbers

Ordinal The relative sizes of the numbers assigned to the objects reflect the amounts of the attribute the objects possess. Equal differences between the numbers do not imply equal differences in the amounts of the attributes.

Hardness of minerals, grades for achievement, ranking on a personality trait, military ranks

Interval A unit of measurement exists by which the objects not only can be ordered but may also be assigned numbers so that equal differences, between the numbers assigned to objects reflect equal differences in the amounts of the attribute measured. The zero point of the interval scale is arbitrary and does not reflect absence of the attribute.

Calendar time, Fahrenheit and centigrade temperature scales

The numbers assigned to objects have all the properties of those of the interval scale, and in addition an absolute zero point exists on the scale. A measurement of 0 indicates absence of the property measured. Ratios of the numbers assigned in measurement reflect ratios in amounts of the property measured.

Height, weight, numerosity, time, temperature on the Kelvin (absolute zero) scale

The discussion of measurement scales presented above has been dogmatic. We have attempted to plead the case of a small group of psychologists who have strong opinions about the level at which measurement is performed. We cannot present their arguments as well as they, so we recommend that you go to their own writings before you pass judgment on their position. Read such works as the following in which both pro and con opinions about the didney of the above concepts are presented. Several of these articles will be beyond the grasp of the typical beginning student. However, they ought not to be overlooked at some later stage in one's studies.

Anderson, N. H., "Scales and statistics: parametric and nonparametric," Psychological Bulletin, 1961, 58, No. 4, 305-16.

Kaiser, H. F., "Review of Measurement and Statistics by Virginia Senders," Psychometrika, 1960, 25, 411-13. This review of Senders' text is highly critical of the position held by Stevens and Senders.

Lord, F. M., "On the statistical treatment of football numbers," American Psychologist, 1953, 8, 750-51. This satirical article by a noted psychometrician and statistician is a cogentargument against the idea that one's scale of measurement ductates which statistics can be used.

MEASUREMENT SCALES

Senders, Virginia L., Measurement and Statistics (Oxford University Press, New York, 1958). This textbook is organized around Stevens' concepts; her position is one of the more extreme ones taken by psychologists.

13

Siegel, S., Nonparametric Statistics (McGraw-Hill, New York, 1956). Siegel's position is identical to that of Stevens. Siegel's text emphasizes which position is identical techniques are "appropriate" to which scales of measurement. Although a useful text in many respects, its emphasis on "permissible" and "appropriate" statisties is perhaps misplaced.

Stevens, S. S. (Ed.), "Mathematics, measurement and psychophysics," in *Handbook of Experimental Psychology* (Wiley, New York, 1951), pp. 1-49. This early article revived interest in the problem of measurement scales and precipitated debate on the issue.

You are likely to get the impression from the works of Stevens, Senders, Siegel, and others that in some metaphysical way a "scale" underlies certain attributes. A certain set of numbers assigned to a group of objects definitely fits into this one category or that one: the scale is either nominal, ordinal, ints into this one category of that care in between. This position can lead interval, or ratio; and there is nothing in between. to chaos if held to in the face of the less well organized feelings of those who actually perform psychological and educational measurements. Those in the Stevens camp maintain, for example, that IQ scores form an ordinal scale, not an interval scale. Uncritical acceptance of this decree forces one scare, not an interval scare to disregard completely the magnitude of the differences between IQ scores. Suppose Joe has an IQ of 50, Sam an IQ of 110, and Bob an IQ of 112. Suppose Joe and Ril 1Q of Jo, Sall that can be said is that Bob is more in-If IQ is truly an ordinal scale, all that can be said is that Bob is more in-If 1Q is truly an ordinal searc, an arms of the statement that Bob telligent than Sam who is more intelligent than Joe. The statement that Bob telligent than Sam who is more intelligent than Joe. telligent than Sam who is more about the IQ than are Sam and Joe would not and Sam are more alike with respect to IQ than are Sam and Joe would not and Sam are more ance with respect to the defensible. To say that this last statement cannot be made because IQ's be detensible. 10 say that this area only ordinal would be anarchy. Ask the person who administered the are only ordinal would be analous. As he person who administered the IQ test, and he would have told you before he tested the children that Joe IQ test, and ne would have took book and are enlighten that Joe is far less intelligent than Sam and Bob, who are much closer together. Try is far less intelligent than same and the street together. Try to tell the tester that he must pay no attention to the sizes of the differences to tell the tester that he mild tell you to mind your own business, as he should between scores, and he will tell you to mind your own business, as he should. between scores, and ne will can joe to minus joet own unsiness, as he should. Even though an IQ unit is not a completely equivalent unit of measurement at Even though an IQ unit is not a company of equation, unit of measurement at different IQ levels, IQ scores are not on a par with a lowly ordinal scale. different 1Q ievers, 1Q secretarion as strictly ordinal or interval; perhaps it The IQ scale defies categorization as strictly ordinal or interval; perhaps it is better to speak of it as "quasi-interval."

Is better to speak or important for a researcher to attempt to categorize his scale of measurement. If the numbers a measurer assigns to n different objects are no more than the n ranks  $1, 2, \dots, n$  (an ordinal scale), some statements with the numbers are meaningless in terms of the amounts of the attribute possessed by the objects. The measurer should be forewarned that this is so. He must also realize that if he has arbitrarily given "males" a 2 and "females" a 2 (nominal measurement), the fact that 3 is greater than 2 means nothing about the attribute measured, namely "sex." In this

manner, the distinctions between the various scales can be useful. However, except for a few infrequently used measures (such as time, length, and mass), educational and psychological measurements, especially clinical measurement, defy any easy categorization as "ordinal" or "interval." Surely, the author of a textbook is in no position to pass judgment on the level of measurement at which one is working unless he, too, is intimately involved in the particular problem.

We shall not develop the notions of scales of measurement further. Only a few of the statistical techniques to be discussed in this text were developed with an eye toward the relationship of measures to the things being measured. The nature of this relationship is the concern of the measurement specialist. Statistical methods are means of analyzing numbers as numbers, not as true amounts of some attribute. Any statistical technique can be applied to any conglomeration of numbers (with some limitations, of course), but we know of no technique which refuses to be performed because the numbers put into it are not "proper." Statistical methods (with the possible exception of some psychometric scaling methods) do not add to or subtract from the meaningfulness of the numbers on which they are performed. This point was made with humor and insight by Kaplan (1964, pp. 205-6):

Mathematics can spare us the painful necessity of doing our own thinking, but we must pay for the privilege by taking pains with our thinking both before and after mathematics comes into play.

I recall a childhood puzzle which takes advantage of just this necessity. Three men registered at a hotel, paying ten dollars each for their rooms. The clerk, later realizing that the three rooms constituted a suite, for which the charge was only twenty-five dollars, gave five dollars to the bellhop to refund to the guests. Since five dollars is not evenly divisible by three, as well as for other less subtle reasons, the bellhop kept two dollars for himself and returned only three dollars as a refund. On his way back he calculated as follows. "They each paid ten dollars, making thirty dollars in all. I returned three dollars, or one dollar to each of them, so they each really paid nine. Now three times nine is twenty-seven, and two dollars I kept, making twenty-nine. Where is the thirtieth dollar?" Of course, if his two dollars is subtracted from the twenty-seven, not added, the remainder is twenty-five, the amount paid the hotel. We are quite free to add the numbers if we wish, but not to expect the sum to represent anything in the situation. What is missing in the bellhop's manipulations is not the dollar but good sense; his logic was no better than his morals.

### 2.3 VARIABLES AND THEIR MEASUREMENT

Variables are characteristic of persons or things, e.g., weight, age, reaction time, ideational fluency, reading speed, number of children, number of

students. Intuition and experience tell us that some of these variables are continuous (i.e., measurements of them can take on any value within a certain range), such as weight, age, and reaction time. We know surely that some variables are discrete (i.e., measurements of them can take on only separated values), such as number of children. The most familiar discrete variables are those that are measured by counting. "Number of children" can give rise to the numbers 0, 1, 2, 3, .... It is not normally possible for this variable to take on intermediate values such as 1.75.

On the other hand, we feel confident that if we only had the instrumentation, resources, and time, we could measure continuous variables to as fine a degree as we wished. We choose to stop measuring elapsed time in a footrace after we have determined tenths of seconds. Even though it is reported that the hundred-yard dash was won in 10.4 seconds, more precise timing equipment might have revealed the winning time to be 10.416 seconds. But even this time is not exact; it is merely correct to thousandths of a second. The actual or exact measurement of a variable is something that can never be attained because measurement must always stop short of the exact value.\* Standing opposite the exact value of a variable is a reported value. The reported value is the value which the measuring process produced. We do not expect the reported and actual values of a variable to coincide, but the former expect the reported and actual values of a variable is person's height is 62 in. measured to the nearest inch, then his actual height at that time and under those conditions is between 61.5 and 62.5 in.

The measurement of any continuous variable should always be accompanied by a statement of the accuracy of the measuring process. Races are timed to the nearest tenth of a second; heights may be measured to the nearest inch; ages might be measured to the nearest day. The sensitivity of a measuring process is the smallest unit of the number scale which is reported. Thus the sensitivities in the above three examples are tenths of a second, inches, and days, respectively.

Second, incnes, and uays, respectively.

We often wish to establish limits around any reported value within which the exact value lies. For example, what are the lowest and highest actual heights that will result in a reported height of 58 in. if measurement of height is to the nearest inch? The limits for the exact value around any reported value are found by adding and subtracting one-half the sensitivity of reported value are found by adding and subtracting one-half the sensitivity of the measuring process from the reported value. Thus a person with a reported height of 58 in. has an actual height between 58 in. - (1 in./2) = 57.5 in. and 58 in. + (1 in./2) = 58.5 in. The examples below should clarify this procedure.

An exact value or score must not be interpreted as a "true" or perfectly stable score, which it is not. The actual score may be unstable from one time to the next.

MEASUREMENT, SCALES, AND STATISTICS 18

> A complicated experiment might produce data which are tabulated in a square with rows and columns; and a symbol X could need three subscripts, i, j, and k, to specify a particular number. For example, X1.3.8 (usually written X138) stands for the sixth number in the cell formed by the intersection of row I and column 3.

# 2.5 SIGMA (Σ) NOTATION

You may wish to concentrate only on the first three-quarters of this section the first time through. You may safely omit the material in this section beyond "Rule 3" below; then you can return to studying the remaining fourth of the section before embarking upon Chapter 15.

The analysis of most data involves adding, subtracting, multiplying, and dividing numbers, among other things. Since we want to talk about performing these operations on a group of numbers in general, we will perform operations on the symbols for the numbers.

 $X_1, X_2, \ldots, X_n$  stands for a group of n numbers, any one of which can be referred to as  $X_1$ , the ith number.  $X_1 + X_2$  stands for the sum of the first and second numbers. The ordering of the subscripts is usually completely arbitrary;  $X_2 + X_1$  could be used to designate the sum of the first and second numbers, instead.  $X_1 + X_2 + X_{10}$  stands for the sum of the first, second, and tenth numbers.

Often we want to add up all of the numbers in a group. If there are five numbers in the group, n = 5 and the sum of all the numbers is  $X_1 + X_2 +$  $\dots + X_5$ .  $X_1 + X_2 + \dots + X_n$  stands for the sum of all n numbers in a group when the exact value of n is not specified.

An abbreviation for  $X_1 + X_2 + \ldots + X_n$  which is frequently used is  $\sum_{i=1}^{n} X_{i}$ .

$$\sum_{i=1}^{5} X_{i} \quad \text{means} \quad X_{1} + X_{2} + \ldots + X_{n}.$$

$$\sum_{i=1}^{5} X_{i} = X_{1} + X_{2} + X_{3}. \quad \sum_{i=3}^{5} X_{i} = X_{3} + X_{4} + X_{5}.$$

 $\Sigma$  is the Greek capital letter "sigma."  $\sum_{i=1}^{5} X_i$  is read "the sum of  $X_i$  as i goes from 1 to 5."  $\sum_{i=1}^{n} X_i$  is read "the sum of  $X_i$  as i goes from 1 to n."

Admittedly, this compact Σ-notation is economical. Statisticians make great use of it. You will have to learn how to interpret and use Σ-notation before you learn much about statistics. The notation takes the place of a set of directions; at first you will have to translate the symbols into a set of verbal directions. As you gain greater facility with  $\Sigma$ -notation, however, you will respond automatically to  $\sum_{i=1}^{n} X_i$  without first having to say to yourself, "the sum of  $X_i$  as i runs from 1 to n."

Adding up numbers after something has been done to them, such as multiplying each number by 6, or squaring each number (that is, multiplying it by itself), is as common as simply adding the numbers as they are. Suppose one wants to multiply each of n numbers by 2 and add together the resulting n products. The desired sum will be

$$2X_1 + 2X_2 + \ldots + 2X_n$$

But surely you see that this sum is the same as

$$2(X_1 + X_2 + \ldots + X_n)$$

Using the shorthand  $\Sigma$ -notation discussed above, we can replace  $(X_1+X_2+\ldots+X_n)$  by  $\sum_{i=1}^n X_i$ . The result can be summarized as follows:

$$2X_1 + 2X_2 + \ldots + 2X_n = \sum_{i=1}^n 2X_i = 2\sum_{i=1}^n X_i$$

This result did not come about because of any magic in the number 2; with 4, 60, or 131.4 the result is the same. In fact, if e stands for any constant number (i.e., a number that does not change regardless of what i is), then

$$cX_1 + cX_2 + \ldots + cX_n = \sum_{i=1}^n cX_i = c\sum_{i=1}^n X_i$$
. (Rule 1)

If a constant number c is to be added to each of n numbers, one writes  $X_1+c$ ,  $X_2+c$ , ...,  $X_n+c$ .

The sum of the above values is

$$(X_1+c)+(X_2+c)+\ldots+(X_n+c)=\sum_{i=1}^n(X_i+c).$$

Always with addition of numbers we can regroup them in any way before adding.

$$\sum_{i=1}^{n} (X_i + c) = (X_1 + X_2 + \ldots + X_n) + (c + c + \ldots + c).$$

The first sum in parentheses on the right-hand side above is  $\sum_{i=1}^{n} X_i$ . What about the second sum in parentheses? How many c's are added together? The answer is n. So the second sum equals nc. Consequently,

$$\sum_{i=1}^{n} (X_i + c) = \sum_{i=1}^{n} X_i + \sum_{i=1}^{n} c = \sum_{i=1}^{n} X_i + nc.$$
 (Rule 2)

If c is one constant and d is another, how else can you write  $\sum_{i=1}^{n} (cX_i + d)$ ? (Use Rules 1 and 2.)

Another important expression is the sum of n numbers after each individual number has been squared,

$$(X_1 \cdot X_2) + (X_2 \cdot X_2) + \ldots + (X_n \cdot X_n) = X_1^2 + X_2^2 + \ldots + X_n^2$$
  
which is symbolized as

 $\sum_{i} X_{i}^{*}$ .

Similarly.

$$X_1^3 + X_2^3 + \ldots + X_n^3 = \sum_{i=1}^n X_i^3$$

although in elementary statistics there will seldom be an occasion to use this expression.

Notice that  $\sum X_i$  stands for a single number; the one that results from the addition of n numbers.  $\sum_{i=1}^{n} X_i$  might be 10, 13, or 1300.  $c \sum_{i=1}^{n} X_i$  is the product of two numbers, a and  $\sum_{i=1}^{n} X_i$ .  $\left(\sum_{i=1}^{n} X_i\right) \left(\sum_{i=1}^{n} X_i\right)$  is the product of a number (a certain sum) and itself. We also write it as follows:

$$\left(\sum_{i=1}^{n} X_{i}\right) \left(\sum_{i=1}^{n} X_{i}\right) = \left(\sum_{i=1}^{n} X_{i}\right)^{2}.$$

If  $X_1 = 3$ ,  $X_2 = 6$ , and  $X_3 = 1$ , then  $\sum_{i=1}^{n} X_i = 10$  and  $\left(\sum_{i=1}^{n} X_i\right)^{\frac{1}{n}} = 100$ . Is  $\sum_{i=1}^{n} X_i^{\frac{n}{n}}$  always the same number as  $\left(\sum_{i=1}^{n} X_i\right)^{\frac{n}{n}}$  [Hint: When does  $a^2 + b^2 = (a + b)^2$ ?] Calculate each when  $X_1 = 2$ ,  $X_2 = 1$ ,  $X_3 = 4$ , and

A common expression in statistical analysis is

$$\sum_{i=1}^{n} (X_i + c)^2 = (X_1 + c)^2 + (X_2 + c)^2 + \dots + (X_n + c)^2.$$

 $(X_i + c)^2$ , which is  $(X_i + c)(X_i + c)$ , can be written in a different way:

$$X_t + c$$

$$\frac{X_t + c}{cX_1 + c^2}$$

$$\frac{X_t^2 + cX_t}{X_1^2 + 2cX_1 + c^2}$$

21

It is true, then, that  $\sum_{i=1}^{n} (X_i + c)^2 = \sum_{i=1}^{n} (X_i^2 + 2cX_i + c^2)$ . The expression within the parentheses may be written n times, as follows:

$$X_{1}^{2} + 2cX_{1} + c^{2}$$

$$X_{2}^{2} + 2cX_{2} + c^{2}$$

$$\vdots$$

$$X_{n}^{2} + 2cX_{n} + c^{2}$$

What is the sum of the first column above? It is  $X_1^2 + X_2^2 + \ldots +$  $X_n^2 = \sum_{i=1}^n X_i^2$ . What is the sum of the second column above? It is  $2cX_1 +$  $2cX_2 + \ldots + 2cX_n = 2c(X_1 + X_2 + \ldots + X_n)$ , which may be written compactly as  $2c\sum_{i=1}^{n}X_{i}$ . What is the sum of the third column above? It is

Putting together these three column sums we see that

$$\sum_{i=1}^{n} (X_i + c)^2 = \sum_{i=1}^{n} X_i^2 + 2c \sum_{i=1}^{n} X_i + nc^2.$$
 (Rule 3)

Though it is correct to proceed in this way, by writing each individual expression ar a summing columns, it is not necessary. Instead, one can "distribute t' e summation sign" before each term, as follows, and secure the same result more directly:

$$\sum_{i=1}^{n} (X_i + c)^2 = \sum_{i=1}^{n} (X_i^2 + 2cX_i + c^2)$$

$$= \sum_{i=1}^{n} X_i^2 + \sum_{i=1}^{n} 2cX_i + \sum_{i=1}^{n} c^2$$

$$= \sum_{i=1}^{n} X_i^2 + 2c\sum_{i=1}^{n} X_i + nc^2.$$

$$= \sum_{i=1}^{n} X_i + \sum_{i=1}^{n} \sum_{i=1}^{n} x_i + nc^2.$$

Note carefully how the summation sign was placed in front of each of the three terms after the squaring had been done. Also, verbalize to yourself as follows: The sum from 1 to n of the product of a constant (here, 2c) and a variable (here, X) equals the constant (2e) times the sum from 1 to n of the values  $(X_1, X_2, \dots, X_n)$  taken by the variable. Similarly, the sum from 1 to n of a constant (here,  $e^2$ ) is simply n times the constant, this being a fundamental relationship between addition  $(c^2 + c^2 + ... + c^2)$  and multiplication (nc2).

The data below are symbolized by X written with two subscripts, as  $X_{ii}$ ,

where  $i=1,2,\ldots,n$  indicates the position of the *i*th observation or measurement (X) within the *j*th column  $(j=1,2,\ldots,J)$ 

These data could be from an experiment in which n persons were given level 1 of a treatment, a different n persons were given level 2, and so on up to the n persons given level 3. There would be nl different persons in such an experiment. Alternatively, each person might be given all l treatments. The two situations are quite different, as we shall see in later chapters. (For the former, it is not necessary that the number of l in each column be n, the same for all columns. Instead, the number could be  $n_l$ , where l = 1, 2, . . . , l; r<sub>l</sub> sould then mean the number of l in the third column, n<sub>l</sub> the number in the second column, etc. Here we keep the n<sub>l</sub>'s equal, for simplicity.

The sum of all the numbers at level 1 (i.e., column 1) is  $X_{11} + X_{12} + \dots + X_{n1}$ . Notice that the first number in the subscript tells what row the observation is in, and the second number tells the column of the observation. To find the sum for column 1, which is  $X_{11} + X_{21} + \dots + X_{n1}$ , we sum i from 1 to n while j keeps the value 1. We write this as  $\sum_{i=1}^{n} X_{in}$ . The expression  $\sum_{i=1}^{n} X_{in}$  denotes the sum over i from 1 to n while j remains 2, i.e., the sum of the observation in column 2.

 $\sum_{j=1}^{n} X_{1j} \text{ is } X_{11} + X_{12} + \ldots + X_{1J} \text{ (read "the sum over } j \text{ from 1 to } J \text{ for } i = 1).$  This expression denotes the sum of the observations in row 1 of the layout.  $\sum_{i=1}^{n} X_{ii} \text{ is the sum of the } n \text{ numbers in column } j; \sum_{j=1}^{n} X_{ij} \text{ is the sum of the } J \text{ numbers in row } i.$ 

How could we denote the grand sum of all nJ numbers from the experiment? One way would be to add up the numbers in each column individually and then add the J column sums together:

Grand Sum = 
$$\sum_{i=1}^{n} X_{i1} + \sum_{i=1}^{n} X_{i2} + ... + \sum_{i=1}^{n} X_{i,i}$$
.

This sum of J numbers can be denoted more simply:

Grand Sum = 
$$\sum_{i=1}^{J} \left( \sum_{i=1}^{n} X_{ij} \right) = \sum_{j=1}^{J} \sum_{i=1}^{n} X_{ij}$$
.

The grand sum squared has the form  $\left(\sum_{j=1}^{J}\sum_{i=1}^{N}X_{ij}\right)^{2}$ . The symbol for the sum of all nJ numbers which result from squaring each original observation and then adding together the squared numbers is  $\sum_{j=1}^{J}\sum_{i=1}^{N}X_{ij}^{2}$ . This expression is read as follows: "The double summation of X sub ij squared as j runs from 1 to J and I runs from 1 to J." First j is given the value 1 as i runs from 1 to J, etc. 1 to J, etc.

The sum, across columns, of the squared values of each column sum is denoted by

$$\sum_{i=1}^{J} \left( \sum_{i=1}^{n} X_{ij} \right)^{2}.$$

There will be a time when we want to talk about adding one constant value to every number in column 1 of an array and a different constant to every number in column 2. Since the value of the constant depends only on the column and not on its position in the column, the subscript i is not needed to identify the constant. It is sufficient, then, to speak of  $c_i$  and  $c_2$ . We can denote a constant value for the jth column by  $c_i$ . Thus, if we wanted to talk about  $X_{ij}$  plus a constant which is different for each column but the same for all n observations in one column, we denote this quantity by  $X_{ij}$ ,  $+c_i$ . The set of such quantities for the jth column is

$$X_{1j} + c_j$$

$$X_{2j} + c_j$$

$$\vdots$$

$$\vdots$$

$$X_{nj} + c_{n}$$

Thus

$$\sum_{i=1}^{n} (X_{ij} + c_{j}) = \sum_{i=1}^{n} X_{ij} + nc_{j},$$
 (Rule 4)

because " $c_j$  is a constant with respect to summation over i." For double summation,

$$\sum_{i=1}^{J} \sum_{j=1}^{n} (X_{ij} + c_j) = \sum_{j=1}^{J} \left( \sum_{i=1}^{n} X_{ij} + nc_j \right) = \sum_{j=1}^{J} \sum_{i=1}^{n} X_{ij} + n \sum_{j=1}^{J} c_j.$$

If a constant value d were to be added to all nJ observations, then no subscripts for d would be needed. The value of d is the same regardless of

24

what row and column it is in. You should be convinced that

$$\sum_{i=1}^{J} \sum_{i=1}^{n} (X_{ii} + d) = \sum_{j=1}^{J} \sum_{i=1}^{n} X_{ij} + nJ \cdot d,$$
 (Rule 5)

Because d is added to observations n times in each of J rows, it figures in the grand sum a total of nJ times.

If you are not yet convinced, list the symbols for every one of the nJ observations or measurements, as below, and sum them all:

Obviously, there are n d's in each column, and there are J columns, so there are n.i d's in all.

### PROBLEMS AND EXERCISES

- 1. Categorize each of the following as either nominal, ordinal, interval, or ratio measurement:
  - a. Zip-code numbers
  - b. Academic rank (assistant professor, associate professor, professor) as a measure of length of service
  - c. Metric system of measuring distance
  - d. Telephone numbers
  - 2. A teacher builds a spelling test by selecting a representative sample of 200 words from a particular dictionary.
    - a. What scale of measurement is being employed if the teacher scores the test as follows:
      - 0-the student spelled at least one plural word incorrectly 1-the student spelled all plurals correctly
    - b. The teacher counts the number of correct spellings and calls this number a measure of "general spelling ability." What scale of measurement is being used (nominal, ordinal, interval, or ratio)?
  - 3. Determine the limits of the exact value corresponding to the reported value in each of the following instances:

Variable	Sensitivity of measurement	Reported value	Limits of exact value	
a. Age b. Weight	Nearest month Nearest half ounce	6 yr 5 months		

Nearest half ounce c. Monetary value Nearest dollar

2 lb 13.0 oz

- 4. Let the symbol X., denote the ith datum in the ith group of data.
  - a. Write the symbol for the 1st datum in the 2nd group
  - b. Write the symbol for the 4th datum in the 1st group.
  - c. Write the symbol for the 2nd datum in an arbitrary group.
- 5. Let  $X_1 = 2$ ,  $X_2 = 7$ ,  $X_3 = 1$ ,  $X_4 = 3$ ,  $X_5 = 2$ , and  $X_5 = 4$ . Evaluate each of the following:

a. 
$$\sum_{i=1}^{6} X_i =$$
 b.  $\sum_{i=1}^{2} X_i =$  c.  $\sum_{i=1}^{4} X_i =$  d.  $\sum_{i=1}^{6} X_i - \sum_{i=1}^{5} X_i =$ 

$$c. \sum_{i=2}^4 X_i =$$

d. 
$$\sum_{i=1}^{6} X_i - \sum_{i=3}^{5} X_i =$$

6. Let 
$$X_1 = 0$$
,  $X_2 = 4$ ,  $X_3 = 8$ ,  $X_4 = 2$ , and  $X_5 = 1$ . We see that  $\sum_{i=1}^{5} X_i = 15$ . Evaluate each of the following without operating on the original five numbers.

a. 
$$\sum_{i=1}^{5} 4X_i =$$
 (Rule 1) b.  $\sum_{i=1}^{5} (X_i + 3.1) =$  (Rule 2)  
c.  $\sum_{i=1}^{5} (X_i - 2) =$  d.  $\left(\sum_{i=1}^{5} X_i\right)^3 =$ 

7. Consider the following data:

$$X_{11} = 4$$
  $X_{12} = 3$   
 $X_{21} = 2$   $X_{22} = 2$   
 $X_{31} = 6$   $X_{32} = 1$   
 $X_{43} = 3$   $X_{42} = 5$ 

a. 
$$\sum_{j=1}^{2} \sum_{i=1}^{4} X_{ij} \approx$$
 b.  $\sum_{i=1}^{4} X_{i1} \approx$ 

c. 
$$\sum_{j=1}^{2} X_{3j} = \text{d. } \sum_{i=1}^{4} X_{i2}^{2} =$$

8. Write out the following expressions:

a. 
$$\sum_{i=1}^{4} X_i^2 = b. \sum_{i=4}^{6} X_i = c. \left(\sum_{i=1}^{3} X_i\right)^2 =$$

9. Change the following expressions into sigma-notation:

a. 
$$3X_1 + 3X_2 + 3X_3 =$$

b. 
$$(X_1 + ... + X_{10})^2 =$$

c. 
$$(X_1 + \ldots + X_n) + 7n =$$
  
d.  $(X_1^2 + X_1) + (X_2^2 + X_2) + \ldots + (X_5^2 + X_5) =$ 

3

# TABULATING AND DEPICTING DATA

### 3.1 TABULATION OF DATA

Before quantitative data can be understood and interpreted, it is usually necessary to summarize them. Table 3.1 shows a class record for a reading readiness test administered at the beginning of the school year. The scores appear in alphabetical order as they are recorded in the teacher's class roll book. However, the scores do not mean very much in this form, and we can tell only with some difficulty whether, for example, the first-listed pupil (David A), with a score of 90 points out of a possible 128, is superior or just average in reading readiness, compared with his classmath.

### Rank Order

Ordinarily the first step is to arrange the scores in order of size, usually from highest to lowest. This is called an ungrouped series. In a small class, this is often all that is necessary. Table 3.2 shows the same 38 scores as Table

3.1, arranged in order of size from 112 to 44. This table also shows the rank order of the pupils (1st, 2nd, ..., 38th) and the scores tabulated without further grouping. It is now easy to see that David A's score of 90 gives him a rank of 13 in a class of 38, or about one-third of the way from the top. Similarly, it is easy to interpret each of the other scores in terms of rank. But ties are likely to occur, especially in classes of 20 or more pupils. Notice, for example, that two pupils made a score of 97. Since it is not correct to say that one ranks higher than the other, we must assign them the same rank. Since there are six pupils who rank higher (1, 2, 3, 4, 5, 6), the next two ranks, 7 and 8, are averaged, giving 7.5. In like manner the average of ranks 9 and 10 is 9.5, and so on for the other pupils with tied scores. There are three pupils with scores of 75, and there are 21 pupils who rank above this score; the average of the next three ranks (22, 23, and 24) is 23, which is the rank assigned to each of the scores of 75. In addition to the time and trouble required to determine these ranks, the list is long, unwieldy, and inadequate for making comparisons with other classes that are much larger or much smaller; ranking 19th in a class of 38 pupils is poorer than ranking 19th in an equally capable class of 70 pupils.

TABLE 3.1 A CLASS RECORD FOR A READING READINESS TEST (38 PUPILS)

Pupil	Score	Pupil	Score	Pupil	Score	Pupil	Score
David A.	90	Robert D.	59	Jerome L.	75	Paul S.	81
Barbara B.	66	Dan F.	95	Rosa M.	75	Richard S.	71
Charles B.	106	Larry F.	78	Billy N.	51	Robert S.	68
Robert B.	84	Richard G.	70	Nancy O.	109	William S.	112
Mildred C.	105	Grover H.	47	Carrie P.	89	Jean T.	62
Robbin C.	83	Robert H.	95	Ralph R.	58	Adolfo W.	91
Robert C.	104	Sylvia H.	100	George S.	59	Dolores W.	93
Diney D.	82	Warren H.	69	Gretta S.	72	Richard W.	84
Jım D.	97	Clarence K.	44	Jack S.	74		
John D.	97	David K.	80	Mary S.	75		

# The Frequency Distribution

The list of scores can be made shorter by arranging the scores in a frequency distribution, sometimes simply called a distribution. The third and fourth columns of Table 3.2 show the simplest form of a distribution. The various scores are arranged in order of size, here from 112 to 44, and to the right of each score is recorded the number of times it occurs. Each entry to the right of a score is called a frequency, abbreviated f, and the total of the frequencies is represented by n.

READING READINESS SCORES FROM TABLE 3.1 ARRANGED IN ORDER OF SIZE. TABLE 3.2 RANKED, AND TABULATED

Order of size Rank		Tabulated without further grouping Score Frequency (1)		
Order of size	Kank	Score	Frequency (f)	
112	1	112	11	
109	2	109	il	
106	3	106	il	
105	4	105	il	
104	5	104	il	
100	6	100	i l	
97)	<b>[7.5</b> ]	97	2	
971	17.5	95	2	
951	(9.5 19.5	93	Sum = 19	
95)	19.5	91	i l	
93	11	90	i l	
91	12	89	i	
90	13	84	2	
89	14	83	îl	
84)	115.5	82	il	
841	115,5	81	1.1	
83	17		Midpoint of	
82	18	80	1) frequencies	
81	19	78	il	
80	20	75	اذ	
78	21	74	íl	
75)	(23	72	il	
75}	23	71	îl	
75) 74	(23	70	il	
72	25	69	11	
71	26	68	1 Sum = 19	
70	27	66	i	
69	28	62	i	
68	29 30	59	2	
66	31	58	1	
62	32	51	1	
59)	j33,5	47	1	
59	33.5	44	1)	
58	35		n = 38 = 19 + 19	
51	36	1	** 1 17	
47	37	1		
44	38	1		

# Grouped Frequency Distribution

For a large number of scores—say 100 or more—it may be desirable to carry the summarization of data one step further. As a rule, there is such a wide range of scores that it is economical to group them according to size. such as a group including all scores from 105 to 109, inclusive, from 110 to 114, inclusive, and so on. Each group is called a score class. The complete grouping arrangement is usually referred to as a grouped frequency distribution. Although there is no fixed rule for the number of score classes, it is usually best to make not fewer than 12 classes nor more than about 15. To have fewer than 12 classes is to run the risk of distorting the results, whereas more than 15 classes produce a table that is inconvenient to handle.

#### Constructing the Grouped Frequency Distribution

There are four steps in making the ordinary grouped frequency distribution. These are shown in Table 3.3, using the scores given in Table 3.1.

- 1. Determine the inclusive range, which is 1 plus the difference between the highest score and the lowest. Of these scores, the highest is 112 and the lowest is 44, which gives a range of (112 - 44) + 1 = 69. Actually, 112 is considered to cover the one-point score interval 112.5-111.5, and 44 the interval 44.5-43.5. Notice, therefore, that the range is 69 1(112-44)+1, or 112.5-43.51. The real score limits are not always fractional, however. If age is reckoned at the last (most recent) birthday, then persons who report themselves as being 44 years old (that is, not yet 45) lie within the interval 44.00-44.99 ... (almost, but not quite 45.00), whose midpoint is 44.5. If they report age to nearest birthday, the interval is 43.5-44.5, with a midpoint of 44. Similarly, if they report themselves "going on 44," the interval is 43.00-43.99 ..., with midpoint 43.5. There will be a difference of almost two years between the youngest possible "going on 44" person, who has just reached the age of 43, and the oldest possible "44 last birthday" respondent, who is almost 45. When we ask merely for "Age\_\_\_\_," without specifying the reckoning system, we will not be able to interpret our results precisely.
- 2. Select the score-class grouping interval, which is the width of the groups into which the scores are to be classified, so that there will be not fewer than 12 score classes nor more than 15. To do this, divide the range by 12 to find the largest group, or score-class interval to be used. Divide the range by 15 to find the smallest class interval to be used. Divide the range by 15 to find the smallest class interval to be used. In this case, 69 ÷ 12 = 5.75, and 69 ÷ 15 = 4.60. Since it is impractical to use any class interval except a whole number, the larger number, 5.75, is "rounded down" to 5 and the 4.60 "rounded up" to 5, even though a class interval of 6 would yield 12 score classes for these 38 scores. Odd-numbered interval widths such as 5, which have whole-number midpoints when the score-class limits are fractional (end in .5), are usually preferred to even-numbered interval widths, which have fractional midpoints when the class limits are fractional. The midpoint of the score class 110-114, which contains the 5 scores 110, 111, 112, 113,

and 114, is 112 (that is,  $110 + [(114 - 110) \div 2] = 110 + (4/2) = 110 + 2 = 112$ ). (Another way to determine the midpoint of an interval is simply to average the reported limits of the interval:  $(110 + 114) \div 2 = 112$ .) If a class size of 6 were used, with score limits of 108 - 113, for example, the midpoint of this even-numbered group would be 110.5, which might result in more complex computations. Hence, a class interval of 5 is preferable to 6 when the class limits are fractional.

- 3. Determine the limits of the classes. There must, of course, be enough classes to include the highest score and the lowest score. To facilitate tabulation, start each class with a multiple of the class interval. If the lowest class starts with 40, which is a multiple of 5, it will accommodate the lowest score, 44, whereas a class beginning with 45 will not. Each succeeding whole-number class lower limit will be 5 points above the one just below it. The next class will start at 45, the next at 50, and so on, until the highest score, 112, is included in the class 110-114.
- 4. Make the tabulation. A tally is made for each score opposite the class in which it falls. To make the tabulation it is not necessary to have the scores arranged in order, for this process may require more time than the tabulation itself. In the original alphabetical list, the first score is 90. In the tabulation column opposite the class which begins with 90, a tally line is drawn to indicate the score. The next score is 66. This falls in the class which begins at 65, so a tally is made there. In the same way, a tally is placed in the column opposite the appropriate class for each of the other scores.

In the finished table, the steps by which it was made do not appear. Only two columns occur in the simplest form of a frequency distribution. The first shows the various classes, usually arranged in descending order from top to bottom, and the second shows the frequencies—the number of scores in each class.

To be sure that you understand Steps 3 and 4, above, stop at this point and construct a grouped frequency distribution of the 38 scores, using a class (grouping) interval of 6. Does the number of classes that result meet the 12-15 criterion suggested in Step 2, above?

When two or more groups of data are to be compared, it is usually best to include all the data in the same table. In that case there will be a column for the classes into which the scores are grouped and one for each of the schools or grades being compared. Table 3.4 shows a frequency table which combines the record of six schools on a certain text. The number of grouping intervals varies from 9 for School F to 17 for Schools A and D, although some intervals have no tallies.

31

TABLE 3.3 AN ILLUSTRATION OF THE PROCESS OF MAKING A GROUPED FREQUENCY DISTRIBUTION

Original scores			
(from Table 3.1)	Steps in	making the distribi	ution
90	Step 1. Determining the i	range.	
66	Highest score	112	
106	Lowest score	44	
84	Range = Differen	1 = 68 + 1	= 69
105			_ 0,
83			
104			
82	Step 2. Selecting the class	interval.	
97	$69 \div 12 = 5.75, 1$		al desirable.
97		Round down to 5.	
59	$69 \div 15 = 4.60$ , s		val desirable.
95	1	Round up to 5.	
78			
70			
47			
95	Steps 3 and 4. Determining	g the limits of the	e classes and making
100	the tabulati		
69			
44			
80	Whole-number limits		
75	of the 15 classes	Tally	Frequency (f)
75 -			
51	110-114	1	1
109	105-109	III	3
89	100-104	II .	2 4
58	95_99	1111	4
59	90-94	III	3
72	85-89	1	3 1 6
74	80-84	1111 /	6
75	75-79	1111	4
81	70-74	!!!!	4
71	65-69	!!!	,
68	60-64	<i>I</i> ,,,	,
112	55-59	iji	4 4 3 1 3
62	50-54 45-49	′,	i
91	45-49 40-44	;	i
93 84	40-14	,	$n = \frac{1}{38}$
04			n 38

#### The Form of the Table

A few words should be said about the pure mechanical makeup of the table as it often occurs in typed or printed form; Table 3.4 and the other tables in this book illustrate one format for the printed table. Each table bears an

TABLE 1.4 DISTRIBUTION OF READING READINESS SCORES FOR EACH OF SIX SCHOOLS IN A CERTAIN CITY

IN A CERT	School					All six	
Score	A	В	C	D	E	F	schools
				1			1
120-124				-			
115-119			1				1
110-114			3		2	2	7
105-109		_	,	2	5	3	15
100-104		3	2 4 3	4	4	5	23
95-99		6	4	5	6	10	31
90-94	5	2	3	4	4	î	18
85-89	4	4	1	6	4	8	29
18-03	2	3	6		ī	,	26
75-79	10	5	4	4 7	6	2	29
70-74	6	2 4 3 5 2 4	4	7	4	ĭ	21
65-69	9	4	3	_	i		13
60-64	4	5	1	2	1		13
55-59	1		3		1		5
50-54	1		1				2
45-49	1		1		_		
40-44			1	2	2		5 2 2 5 2 2
35-39	1	1					
30-34		2 1					1
25-29		i					
20-24							
15-19							_
10-14	1						1
N'	45	38	38	37	40	36	234

identifying number. Although either Roman or Arabic numerals may be used, Arabic numerals seem to be increasingly favored. The table number may be centered above the table title, or it may be given at the beginning of the title. The table often starts with a single or double horizontal line and usually ends with a single horizontal line. Another horizontal line separates the column headings from the table body, and other horizontal lines exparate any summarizing measures that are given under the table proper. Vertical lines may be used to separate the columns, but usually no lines are drawn along the margins of the page. It is considered good form to avoid abbreviations in the table whenever possible, and to make the title and headings complete enough to indicate the contents of the table clearly.

Wallis and Roberts (1965, Chap. 9) present an excellent chapter on the art of reading statistical tables. They remind the reader of the obvious precautions, e.g., read the title and headnotic carefully, and they demonstrate some subtle and sophisticated techniques for extracting hidden information to tables. Wallis and Roberts have managed to do all this and be entertaining as well.

#### 3.2 QUANTILES

One of the most efficient and useful methods of describing a group of observations is by means of quantiles. A quantile is a concept and percentiles, deciles, and quantiles are three examples of it. A quantile is a point on a number scale which is assumed to underlie a set of observations; the quantile divides the set of observations into two groups with known proportions in each group. For example, there are three quartiles  $(Q_1, Q_1, \text{ and } Q_2)$ ; they divide a group of observations into four quarters.  $Q_1$  is that point on the number scale such that one-fourth the observations lie below it; one-half the observations lie below  $Q_2$ , and three-fourths of the observations lie below  $Q_3$ . Thus the three quartiles divide a set of observations in each portion. The 99 possible percentiles  $(P_1, \dots, P_n)$  divide a set of observations. The nine deciles  $(D_1, \dots, D_n)$  divide a set of observations.

If 25% of the observations are always below  $P_{25}$ , the 25th percentile, and the same is true for  $Q_1$ , the first quartile, then  $P_{25}$  must equal  $Q_1$ . Figure 3.1




FIG. 3.1 Relationships among quantiles.

shows the relationships among the various quantiles defined so far and still another type called quintiles. (Quint for five.) The four quintiles divide a group into fifths. We shall denote them  $K_1, K_2, K_3$ , and  $K_4$  since the letter Q was used for quartiles.

Quantiles are very useful for summarizing data. Simply reporting that  $P_{\rm g}$  is 10.75 and  $P_{\rm 35}$  is 16.80 tells immediately that 5% of the observations are less than 10.75 and that 10% of then lie between 10.75 and 16.80. For some large groups of data that are familiar, the entire collection of observations can be pictured in a reader's mind if he knows only the values of three or four percentiles, for example. All too often, more complicated summary measures (such as those to be encountered soon) are used to describe data when certain quantiles are more easily calculated and more readily understood. It is regrettable that quantiles are not more widely used by persons attempting simply to describe a set of data.

# 3.3 DETERMINING PERCENTILES

Because of the relationships among various quantiles noted in Fig. 3.1, one need know only how to determine percentiles to find the values of any quantiles desired. (Typically, no one ever wishes to divide a group of observations into more than 100 quantiles.)

The definition of a percentile is simple: the Pth percentile is the point below which P percent of the scores lie. Calculating a given percentile is slightly more complicated than the definition might lead you to believe.

Before beginning the calculation of any percentile in a group of scores one must arrange the scores from smallest to largest. This may be a time-consuming operation in large groups of scores, and it may be more convenient to tabulate the scores in a grouped frequency distribution. The method we will present for finding a percentile point is general and applies to either a ranking or a grouped frequency distribution of the scores.

A teacher administered a 40-item achievement test to his 125 students. The test score was taken to be the number of questions answered correctly. An ungrouped frequency distribution of the 125 test scores appears in Table 3.5.

What is the 25th percentile in the group of 125 test scores, i.e., what is the value of  $P_{25}$ ?  $P_{25}$  is the point below which 25% of the 125 scores lie.

The calculation of any percentile will be facilitated if a cumulative frequency distribution is constructed. The cumulative frequency up to any given score is the total number of frequencies at or below that score. In the third column in Table 3.5 you will find the cumulative frequencies for the L2S achievement test scores. Notice, for example, that there are 106 persons with test scores of 33 or less. The cumulative frequency through the test score of 33 is 106.

The calculation of  $P_{25}$  can be accomplished in five steps:

Step I. Find (.25)n by dividing n by 4:

$$\frac{125}{4}$$
 = 31.25.

Step 2. Determine the lower real limit L of the score class containing the 31.25th person from the bottom.

Because 16 persons scored at or below a score of 28, and 34 persons scored at or below a score of 29, the 31.25th frequency lies on the score-class interval 28.5-29.5.

It will be assumed that the 18 frequencies at a score of 29 are evenly spread along the score class 28.5-29.5. Each frequency occupies (1/18)th

35

DETERMINATION OF P<sub>15</sub>, THE 25th PERCENTILE, IN A FREQUENCY DISTRIBUTION
WHERE THE SCORE-CLASS INTERVAL IS ONE

Test score	Frequency	Cumulative frequency	Calculations
38	1	125	п 125
37	1	124	Step 1. $0.25n = \frac{n}{4} = \frac{125}{4} = 31.25$ ,
36	3	123	Step 2. Find lower real limit of score class
35	5	120	containing 31,25th score:
34	9	115	$L \approx 28.5$ .
33	8	106	
32	17	98	Step 3. Subtract the cumulative frequency up to
31	23	81	L from 31.25:
30	24	58	31,25 = 16 = 15.25.
29	18	34	
28	10	16	Step 4. Divide the result of Step 3 by the
27	3	6	frequency f in the interval containing
26	1	3	the 31.25th score:
25	0	2	15.25
24	2	2	$\frac{15.25}{18} = 0.85.$
	$n = \overline{125}$		Step 5. Add the result of Step 4 to L:
			$P_{15} \approx 28.5 + 0.85 = 29.35$

of a unit. Determining how much of the interval the 31.25th score cuts off is a problem of interpolation within the interval. Steps 3 and 4 accomplish the interpolation.

Step 3. Subtract the cumulative frequency (cum.f) up to L from .25n. L is 28.5, and 16 frequencies have accumulated up to L. Hence,

$$.25n - (cum.f) = 31.25 - 16 = 15.25$$
.

In Step 3, one determines how many frequencies in the interval 28.5-29.5 lie below. 25n.

Step 4. Divide the result of Step 3 by the frequency f in the interval containing the .25nth frequency.

5-

$$\frac{15.25}{18} \approx 0.85$$
.

Step 4 is the determination of what fraction of the score-class interval lies below the .25nth frequency. There are 18 frequencies in the interval 28.5-29.5, and 15.25/18 = 0.85 of the interval is occupied by the first  $15\frac{1}{2}$  frequencies.

36

Step 5. Add the result of Step 4 to L. The sum is P25.

$$P_{\rm st} = 28.5 + 0.85 = 29.35.$$

Under the conventions we have adopted for the presentation of the scores,  $P_{25} = 29.35$ ; i.e., 25% of the 125 scores lie below 29.35. (Similarly, 75% of the 125 scores lie above 29.35.)

Steps 1 through 5 can be expressed in a single formula:

$$P_{25} = L + \frac{25n - (cum.f)}{f},$$
 (3.1)

where L is the lower real limit of the score interval of length 1 containing the .25nth frequency from the bottom of the distribution,

cum. f is the cumulative frequency up to L, and

f is the frequency within the score interval containing the .25nth frequency.

A more general form of Eq. (3.1) applies to the determination of any percentile in a frequency distribution whose score-class interval is 1. Suppose we wish to find the point which exceeds some proportion p of the frequencies.  $P_p$  represents the pth percentile.

$$P_{\bullet} = L + \frac{pn - (cum.f)}{f}, \tag{3.2}$$

where L is the lower real limit of the score interval containing the pnth frequency.

cum. f is the cumulative frequency up to L, and

f is frequency of scores in the interval containing the pnth score.

We shall illustrate the use of Eq. (3.2) by calculating  $P_{40}$  from the data in Table 3.5:

$$P_{\rm eq} = 30.5 + \frac{75 - 58}{23} = 31.24.$$

Calculation of any percentile point from a grouped frequency distribution is almost identical to the calculations for the ungrouped distribution. In fact, the formula for the grouped frequency distribution that will be developed includes Eq. (3.2) as a special case when the score interval is one unit wide.

The data in Table 3.6 are the ages to the nearest year of 1982 teachers who participated in special summer programs for the improvement of the teaching of selected high-school subjects.

The general formula for determining the pth percentile in a group of n scores is as follows:

$$P_{\bullet} = L + \frac{pn - (cum.f)}{f}(W), \tag{3.3}$$

where L is the lower real limit of the interval containing the pnth frequency from the bottom.

cum. f is the cumulative frequency up to L,

f is the frequency in the interval containing the pnth frequency, and W is the width of any score interval.

Note that Eq. (3.3) is exactly the same as Eq. (3.2) when W = 1, i.e., when the score interval is one score wide, which means that scores are not grouped more finely than originally in constructing the frequency distribution.

Application of Eq. (3.3) will now be illustrated by finding  $P_{20}$  for the data in Table 3.6. The pnth score—the 396th score—lies in the interval 24–27 that has a lower real limit of 23.5. The difference between pn and the cumulative frequency up to 23.5 is 396.4–135. Noting that the frequency in the interval including the 396th score is 295 and the interval width is 4 units, we obtain:

$$P_{20} = 23.5 + \frac{396.4 - 135}{205} \cdot 4 = 27.04$$

Under the conventions adopted for the calculation of percentiles, we can say that 20% of the teachers were younger than 27.04 years. We do not expect this statement to be exactly true. Errors have entered the process of determining the percentiles at two points. First, ages were reported to the

TABLE 3.6 ILLUSTRATION OF THE CALCULATION OF P<sub>80</sub> IN A GROUPED FREQUENCY DISTRIBUTION

Age interval	Frequency	Cumulative frequency	Calculations
64-67	4	1982	Step 1. $0.50n = 0.50(1982) = 991$ .
60-63 56-59 52-55 48-51	38 82 120 125	1978 1940 1858 1738	Step 2. Find lower real limit of score class containing 991st score: $L = 31.50$ .
44-47 40-43 36-39	160 221 204 307	1613 1453 1232 1028	Step 3. Subtract the cumulative frequency cum.f up to L from 991: 991 - 721 = 270,
32-35 28-31 24-27 20-23	291 295 135 n = 1982	721 430 135	Step 4. Divide the result of Step 3 by the frequency f in the interval containing the 991st score: $\frac{270}{307} = 0.88.$

Step 5. Multiply the result of Step 4 by the width W of the score class: (0.88)(4) = 3.52.

Step 6. Add the result of Step 5 to L:  $P_{ab} = 31.50 + 3.52 = 35.02$ .

nearest year instead of to the nearest month, day, hour, or minute. Second, it was assumed that the frequencies within each score interval were evenly distributed over the entire interval. Undoubtedly this assumption was false: at the younger ages frequencies are probably stacked up at the upper end of each score interval; at the older ages, frequencies are probably stacked up at the lower end of each interval. In making the assumption of evenly distributed frequencies, a compromise between computational labor and errors of approximation was made. The error made in approximating P20 or any other percentile is probably immaterial relative to the difficulties that would arise if we assumed that the frequencies were distributed in any manner other than evenly on each score interval.

That percentiles are fractional (e.g., 27.04) in measuring age is reasonable. We have no difficulty imagining that someone is precisely 27.04 years of age. What if the variable being measured is discrete? Suppose we build a frequency distribution of sizes of kindergarten classes in a large city school system. This variable, "size of class," can take only values such as 25, 26, 27, 28, etc. It is absurd to speak of a class with 27,31 students. Yet if we were to build a frequency distribution of class sizes and calculate percentile points by the methods in this chapter, we would most certainly obtain fractional values. Can such fractional percentile points be taken seriously? Isn't it utterly false that 81% of the classes can have a size of 32.41 while 89% have a size of 32.78? Of course, it is utterly false. The same percent of the classes have 32.41 or fewer students and 32.78 or fewer students, and this percent is exactly the number of classes with 32 or fewer students. Even though fractional percentiles in distributions of discrete variables cannot be reconciled with common sense, they are useful and widely employed. The alternative to using them is to abandon this convenient and helpful process of converting scores to percentiles and adopt some more cumbersome procedure. No one seems willing to do this just because a class with 32.50 students seems a little ridiculous

#### 3.4 GRAPHING DATA

There can be little doubt that the graphical representation of educational data is a valuable supplement to statistical analysis and summarization. A graph or chart tends to attract the reader's attention. The average casual reader is likely to give scant attention to the ordinary printed matter in a research report and to be unimpressed by the mass of tabular data often piled up at the end. However, his eye is likely to be arrested by any picture or chars that may happen to be included, and this may lead him to read the entire discussion.

A graph is often an effective method of clarifying a point. One small graph will often make a point more clearly than a dozen tables or paragraphs.





FIG. 3.2(a) Projected school enrollments, 1965-1980. Data refer to the civilian noninstitutional population five to 34 years old. Copyrighted; reproduced with permission of the National Industrial Conference Board, 845 Third Asenue, New York, N.Y. 10022.

# THE CURVE CHART WHICH CHART TO USE? Thousands THE DATA 100 Motor Buses in Operation in United States THE GROUPED COLUMN Thousands of Buses 100-PREVENUE ŧΩ 25



REVENUE MOTOR BUSES in Operation


1945

1949


80 60


> 40 20

> > 1941



1941 1943 1945 1947






# 1941 1945 1948 1949


THE PICTOGRAM


FIG. 3.2(b) The same data plotted in 15 different ways. From Mary E. Spear, Charting Statistics (New York: McGraw-Hill Book Company, 1952). enspapers, by permusion of the publisher. 6-1952 McGraw-Hill Book

#### THE INDEX CHART




## THE SUBDIVIDED COLUMN







## THE PICTORIAL SURFACE




#### THE LOGARITHMIC CHART



#### THE PAIRED BAR Thousands of Buses



#### THE COLUMN and CURVE



#### THE PIE CHART



It is sometimes said that the facts speak for themselves. In reality, statistics often stand speechless and silent, tables are sometimes tongue-tied, and only the graph cries aloud its message. Ordinary numerical data are quite abstract; they convey their meaning vaguely and with effort to the average mind. The picture or graph is a more concrete representation.

A wide variety of graphs and charts are shown in Fig. 3.2(a) and (b). In Fig. 3.2(b) basic information concerning motor buses in operation in the United States is given first in tabular form, followed by 15 different black and white graphs. Each graph conveys a unique point in an impressive manner.

Spear's description of the function of graphs is a fitting conclusion to this introduction.

In the present day, when visual education in all aspects has become, not only an aid to, but also a vital basis of learning, our attention is called more than ever before to the almost limitless possibilities in this field. The eye absorbs written statistics, but only slowly does the brain receive the message hidden behind written words and numbers. The correct graph, however, reveals that message briefly and simply. Its purposes, which follow, are clear from its context:

- 1. Better comprehension of data than is possible with textual matter
- 2. More penetrating analysis of subject than is possible in written text
- This triple purpose of the chart can be carried out through careful planning and familiarity with the functions of all types of graphs and media. The following six steps are fundamental to the development of graphic presentation that will describe statistical data with clarity and dramatic
  - 1. Determine the significant message in the data,
  - 2. Be familiar with all types of charts and make the correct selection. 3. Meet the audience on its own level; know and use all appropriate
  - 4. Give detailed and intelligible instructions to the drafting room.
  - 5. Know the equipment and skills of the drafting room.
  - Recognize effective results.\*

#### 3.5 REPRESENTING A FREQUENCY DISTRIBUTION GRAPHICALLY

The ordinary frequency distribution does not give a very clear picture of the situation. There are three common methods of representing a distri-

<sup>\*</sup> From Mary E. Spear, Chaeting Statistics (New York: McGraw-Hill Book Company. 1952), pp. 3-4, by permission of the publishers. C. 1952 by the McGraw-Hill Book

bution of scores graphically: the histogram or column diagram, the frequency polygon, and the smooth curve.

#### The Histogram or Column Diagram

The histogram is a series of columns, each having as its base one class interval and as its height the number of cases, or frequency, in that class. Figure 3.3




FIG. 3.3 A histogram, or column diagram, representing the percentage values assigned to an arithmetic paper by 42 scorers.

represents a histogram showing the distribution of percentage values assigned to an arithmetic paper by 42 scorers. Since the greatest frequency is 9, in the 59,5-64,5 class, it is not necessary to extend the vertical or frequency scale at the left above 9. And since the scores range from the 29,5-34.5 class to the 74,5-79,5 class, it is necessary to represent the horizontal scale only through that distance. For clarity, however, it is customary to extend




FIG. 3.4 A histogram, or column diagram, representing the distribution of the 83 IQ's in a small junior high school.

the scale one class interval above and below that range. In order to avoid having the figure be too flat or too steep, it is usually well to arrange the scales so that the width of the histogram itself is about one and two-thirds times its height—that is, the ratio of height to width should be approximately 3:5. A column is centered around the midpoint of the score-class interval. In actual practice it is customary to represent the histogram in outline form, rather than to show the full length of the columns. Figure 3.4 illustrates the shaded outline form of the histogram.

#### The Frequency Polygon

The process of constructing a frequency polygon is very much like that of constructing the histogram. In the histogram, the top of each column is indicated by a horizontal line, the length of one class interval, placed at the proper height to represent the frequency in that class. But in the polygon a point is located above the midpoint of each class interval and at the proper height to represent the frequency in that class. These points are then joined by straight lines. As the frequency is zero at the classes above and below those in the distribution, the polygon is completed by connecting the points that represent the highest and lowest classes with the base line at the midpoints of the class intervals next above and below. Figure 3.5 shows a polygon for the same data represented by a histogram in Fig. 3.4.




FIG. 3.5 A frequency polygon representing the percentage values assigned to an arithmetic paper by 42 scorers.

#### The Smooth Curve

Sometimes a smooth curve is drawn instead of a histogram or frequency polygon. The only difference is that for the former a smooth line is drawn as close as possible to the points, and for the latter two figures an angular or jagged line is used.

A smooth curve widely used in representing test scores is the percentile curve or ogive. Figure 3.6 shows a percentile curve used to represent the percentage data already used to illustrate the histogram and the polygon. The points that determine the percentile curve are located on the horizontal line at the upper limit of each class, at the position that indicates on the horizontal scale the percentage of scores up to and including that class. Notice, also, that two columns have been added to the ordinary frequency table. The cumulative frequency column indicates the number of scores up to and including each class. For example, there is one score in the 30-34 class, and there are two in the 35-39 class, making a cumulative frequency of 3 in the two lowest classes. The cumulative percent column shows what percentage each of these cumulative frequencies is of the total. In the




FIG. 3.6 A percentile curve representing the percentage values assigned to an arithmetic paper by 42 scorers.

illustration, the total, n, is 42. The first entry in this column is, of course, 100; the second is 98, because 41 is 98% of 42; the third is 95, because 40 is 95% of 42; and so on for the others. Each value in the cumulative percent column is represented as a point on the upper limit of that class interval (the horizontal line separating that class from the class above it), since it includes the percentage of scores up through that class. These points determine the curve. As a rule, especially in small groups where irregularities are most likely to occur, it is best to miss some of the points in order to obtain a smooth and regular curve; but care should be taken in order to leave about as many points on one side of the line as on the other. In this way the ogive will fit the trend of the points as closely as possible.

10	<u>f</u>	Tally
145-149	1	x
140-144	2	XX
135-139	2	XX
130-134	5	XXXXX
125-129	8	XXXXXXXXX
120-124	5	XXXXX
115-119	16	XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX
110-114	12	XXXXXXXXXXXXXX
105-109	10	XXXXXXXXXXXXX
100-104	8	XXXXXXXXXX
95-99	6	XXXXXXX
90-94	8	XXXXXXXXXX
85-89	6	XXXXXX
80-84	1	X
75-79	_1	x

FIG. 3.7 Bar graph made on a typewriter, showing the distribution of 91 IQ's in a junior high school.




FIG. 3.8 Bar graph made on a typewriter, showing the percentage of pupils of each age group who were graduated from high school and the percentage who entered high school at that age but did not graduate.

#### Typewriter Graphs

A satisfactory bar graph can be made on the typewriter. Figures 3.7, 3.8, and 3.9 illustrate this type of graph. Other graphs, such as the circle or pie graph, and various picture graphs, or pictographs, are occasionally used in educational measurement; these are discussed especially by Snear (1952).

Score	Frequency for Grade			Bar Graph for School Grade				
Scote	7	8	9	Seventh	Eighth	Ninth		
200-219			3	[		999		
180-199	1	4	5	7	8888	99999		
160-179	3	3	7	777	888	9999999		
140-159	4	9	7	7777	888888888	9999999		
120-139	11	7	11	71777777777	8888888	9999999999		
100-119	1,	7	2	7777	8888888	99		
80- 99	4	2	1	7777	88	9		
60- 79	1	3		7	888	1		
40- 59		1			8	1		
20- 39	1	1	ľ		8			

FIG. 3 9 Graph made on a typewriter, showing the overlapping of grades seven, eight, and nine in reading comprehension.

#### Which Graph is Best?

As we might expect, no one type of graph is equally good for all purposes. The histogram is the easiest of all to understand and therefore is usually best if no more than one distribution is being fepresented. But if two or more distributions are to be compared, frequency polygons (or relative frequency polygons) are usually better, since so many lines coincide when histograms are superimposed that the picture is likely to be confusing. The percentile curves have many advantages not possessed by other curves. An important one is that from them it is possible to estimate with a high degree of accuracy the quartiles, medians, and other similar points. As we will see in the next section, by means of percentile curves several groups can be presented, for convenient comparison, on a single graph. The main value of bar graphs, circle graphs, and picture graphs lies probably in school publicity and in the motivation of learning. "A successful graph," as the prominent educator Douglas Scates pointed out long ago, "depends far more on careful thought and judgment than on techniques" (Scates, 1942).

# Representing Two or More Distributions Graphically

It is often desirable to compare two or more distributions. For example, school administrators may wish to compare the entire distribution of verbal ability test scores of the pupils in one school with that of pupils in another school. Also, the overlapping of scores among the various grades within a single building is a striking way to present the need for individualized instruction and varied materials within the same grade.

### Representing Entire Distributions

When it is important to compare two or more entire distributions, as would be the case in a study of the status of a school or school system, the choice will usually be between the frequency polygon and the percentile curve. We have already pointed out the difficulty of superimposing two or more histograms. A series of polygons may be drawn on the same sheet one above the other, or alongside each other. In Fig. 3.9, a method of showing overlapping by using bar graphs made on the typewriter is illustrated. (Perhaps the scores there are grouped too coarsely. According to the conventional rule it would be better to have 12 to 15 score classes, instead of the 7, 9, and 7 actually used in Fig. 3.9.)

#### The Use of Palygons

The distinct advantage of polygons over histograms for representing a series of distributions is that polygons can be superimposed on each other with less crossing of lunes. In this form, comparisons among distributions are more easily made. Figure 3.10 illustrates this possibility with the distribution of reading comprehension scores for the 100 seventh, 100 eighth, and 100 ninth grades of a certain school. (For some purposes it could be quite misleading to graph overlapping frequency polygons for different sizes of groups. In these instances, frequencies should be converted to relative frequencies—proportions—before graphing.) The great overlapping in reading ability of the three grades stands out clearly. But even with only three distributions, the lines cross and recross so many times that it becomes difficult to make any accurate comparison of one grade with another. More than three classes can hardly be represented in the same graph by frequency polygons without considerable confusion.

#### The Use of Percentile Curves

For the graphic comparison of two or more distributions the percentile curve has some distinct advantages. Since the frequencies are reduced to




FIG. 3.10 Frequency polygons representing the distribution of reading comprehension scores on a reading test for the seventh, eighth, and muth grades of a certain school (100 students in each grade).

		event	_		Eighth			Ninth		Percentile scale
Test		Cum		f	Cum		f	Cum	Cum % (	
score	<u>,</u>	-1-	-/0	—	<del>-</del>	_	3	36	100	
200			<u>ا</u>	4	37	100	5	33	92	- of A
180	_1_	28	100		33	89	7	28	78	- John Silver
160	3	27	96	3		81	7	21	59	
140	4	24	86	_9_	30	-	H	14	39	
120	11	20	71	7	21	57		3	8	0.98
100	4	9	32	<u>L7</u> _	14	38	2	۲÷	3	K-95/91
80	4	5	18	2	7	19	1	+-	۲	10/
60	1	1	4	3	5	14	!	<b>⊢</b> –	<del> </del>	15
40	<del>  -</del> -	+-		1	2	5		1_	<b>↓</b> _	ρίω <sup>6</sup>
	<del> </del> -	┼~	-	1	1	3_	L_	L_	<b>!</b>	<u> </u>
20	ļ	<del> </del>	<del> </del>	-	T-	1	Γ	<u> </u>		0, 0, 0,
	1									O <sub>1</sub> O <sub>2</sub> O <sub>3</sub>

FIG. 3.11 Comprehension scores on a reading test for the seventh, eighth, and ninth grades.

percents, it is possible to compare groups of unequal size. Another important advantage is that several distributions can be represented in a single graph without difficulty or confusion. Figure 3.11 shows the distribution of reading

comprehension scores, for the same grades as in Figs. 3.9 and 3.10, in the form of percentile curves.

From these percentile curves we can observe several relationships that were not apparent in the polygons. It is clear that although the seventh and eighth grades have almost exactly the same average scores, the eighth grade has greater variability. This is evident since the upper half of the eighth grade exceeds the upper half of the seventh grade, while the lower half of the seventh.

Furthermore, although the ninth grade runs consistently above the other two grades, about 15 percent of the ninth-grade pupils fall below the median of the seventh and eighth grades.

#### 3.6 MISLEADING GRAPHS

So far in this chapter we have stressed correct procedures in the construction of graphs and have presented many examples of good graphs. The unwary researcher or the propagandist can construct graphs which seriously mislead the reader, however. Perhaps the best protection against committing these errors is to study several examples of poor graphs.




FIG 3.12 Percent of drop-outs at the end of grade 11 for the years 1960-1964

Figure 3.12 is a graph of the percent of high school juniors dropping out of school in a particular district at year's end for the period 1960 through 1964. The percent of drop-outs shows a steady rise from 9% in 1960 to 14% in 1964, but has not even doubled over the five-year span. However, because the vertical scale of the graph does not start at 0%, as it should, the as it was in 1960. The true ratios of drop-outs is six times as great in 1964 es it was in 1960. The true ratios of percents of drop-outs by years are percented when the percent drop-outs scale extends from 0% at its base up to 14%.

In Fig. 3.13, the school bonds for three mythical school districts are graphed. The diameter of each circle represents the amount of the bond in

- It is advisable not to show any more coordinate lines than necessary to guide the eye in reading the diagram.
- The curve lines of a diagram should be sharply distinguished from the ruling.
- 11. In curves representing a series of observations, it is advisable, whenever possible, to indicate clearly on the diagram all the curves representing the separate observations.
- The horizontal scale for curves should usually read from left to right and the vertical scale from bottom to top.
- 13. Figures for the scales of a diagram should be placed at the left and at the bottom or along the respective axes.
- 14. It is often desirable to include in the diagram the numerical data or formulae represented.
- 15. If numerical data are not included in the diagram, it is desirable
- to give the data in tabular form accompanying the diagram.

  16. All lettering and all figures on a diagram should be placed so as to be easily read from the base as the bottom, or from the right-hand
- edge of the diagram as the bottom.

  17. The title of a diagram should be made as clear and complete as possible. Subtitles or descriptions should be added if necessary

Also see Arkin and Colton (1936), Walker and Durost (1936), and Kelley (1947) for more on graphing and tabulating data.

## PROBLEMS AND EXERCISES


 This problem is an exercise in constructing a grouped frequency distribution. The following data are Stanford-Binet intelligence test scores for 75 adults:

	minoru-D	met intelli	gence test :	cores f
141	104	101	130	
92	87	115	91	148
100	133	124		96
132	118	98	92	123
97	124	118	101	107
110	111		146	107
106	135	138	121	129
107	110	97	108	108
105		101	129	105
83	110	116	113	123
	127	112	114	105
127	114	113	106	
95	105	95	105	139
109	102	102	102	106
108	92	131		89
104	94	121	86	134
	- •	421	107	103

- a. Determine the inclusive range of the above set of scores.
- b. Divide the inclusive range [found in (a)] by 12 and by 15, separately.
- c. Choose the class size to be the whole number which lies between the two quotients found in (b) above.
  - d. Using the class size found in (c) above, construct a grouped frequency for the 75 scores:

Interval	Score interval	Tally	Frequency	Cumulative freq.
14	145–149			75
13				
12				
11				
10				
9				
8				
7				
6				
5				
4				
3				
2				
1	80-84			
				$n = \overline{75}$

- 2. Find the 50th percentile,  $P_{80}$ , for the 75 intelligence-test scores from the grouped frequency distribution constructed in Prob. 1.
- Complete the frequency polygon started below for the grouped frequency distribution in Prob. 1. Be sure to label the horizontal axis with the midpoints of the 14 score intervals.



4. Plot two relative frequency polygons on the same graph from the following grouped frequency distributions of Scholastic Apittude Test Verbal scores for the 903 men and the 547 women in the freshman class of a large Eastern university:

	M	en	Wo	тел	
SAT-V score interval	Frequency	Relative frequency	Frequency	Relaine frequency	
750-800	1	.001	4	.007	
700-749	27	.030	28	.051	
650-699	63	.070	56	.102	
600-649	138	.153	85	.155	
550-599	174	.193	117	.214	
500-549	202	.226	128	.234	
450-499	171	.189	86	.157	
400-449	96	.106	32	.059	
350-399	25	.028	9	.016	
300-349	4	.004	ī	.002	
250-299	1	.001	i	.002	
200-249	1	.001	ò	.000	
	n <sub>m</sub> = 903	1.00	n = 547	1.00	

4

# MEASURES OF CENTRAL TENDENCY

#### 4.1 INTRODUCTION

We saw in Chapter 3 how the properties of a collection of scores can be depicted graphically or in tabular form. Frequently a graph or table of data tells us more than we want or need to know, and the message it conveys may be time-consuming to communicate. Usually we single out for description just two or three properties of a set of scores. These properties (e.g., the typical "size" of the scores and their spread) may be describable by indexes known as summary statistics. In this chapter we shall study summary statisties which describe the typical "size" of a set of scores.

The descriptive indexes that will be developed in this chapter would be used to answer a question like "How tall is the typical male graduate student in this university?" If the scores in a set are thought of as positioned along a number line, the property of the set that we now want to describe is where along that line the scores tend to lie. Are they bunched around a score of 71, or do they center about a score of 67? Different measures of the central tendency of a set of scores imply different definitions of a "central score." There are only a few summary measures of central tendency in common use, and in the following sections we shall study them in detail.

The most easily obtained measure of central tendency is the mode. The mode is the score in a set of scores that occurs most frequently. Not every set of scores has a single mode by a strict interpretation of this definition, so the working definition of the mode contains some qualifications and conventions that will be discussed after an illustration.

In the set of scores (2, 6, 6, 8, 9, 9, 9, 10) the mode is 9 because it occurs more often than any other score. Note that the mode is the most frequent score (9 in this example) and not the frequency of that score (3 in this example).

#### 4.3 CONVENTIONAL USE OF THE MODE

- When all of the scores in a group occur with the same frequency, it is customary to say that the group of scores has no mode. Thus there is no mode in the group (0.5, 0.5, 1.6, 1.6, 3.9, 3.9).
- When two adjacent scotes have the same frequency and this common frequency is greater than that for any other score, the mode is the average of the two adjacent scores. Thus, the mode of the group of scores (0, 1, 1, 2, 2, 2, 3, 3, 3, 4) is 2.5.
- 3. If in a group of scores two nonadjacent scores have the same frequency and this common frequency is greater than that for ny other scores, two moder exist. In the group of scores (10, 11, 11, 12, 13, 14, 14, 17), both 11 and 14 are modes; the group of scores is said to be bimodal. Large sets of scores are often referred to as bimodal when they present a frequency polygon that looks like a Bactrian (two-humped) camel's back even though the frequencies at the two peaks are not strictly equal. This slight twisting of the definition is allowed because the term bimodal is so convenient and nicely descriptive. A convenient distinction can be made be-




FIG. 4.1 A frequency distribution of test scores with a major mode at 6 and minor modes at 3.5 and 10.

tween major and minor modes. In a group of scores the major mode is the single score which satisfies the definition of the mode. However, several minor modes might exist at points throughout the group of scores. These minor modes are essentially local peaks on the frequency distribution. For example, in Fig. 4.1 the major mode is at 6 and minor modes exist at 3 and 10.

#### 4.4 THE MEDIAN

You have already encountered the median (though it wasn't called by that name) and learned how to find it in Sec. 3.2 on quantiles.

Definition: The median, Md, is the 50th percentile in a group of scores. It is the score that divides the ranked scores into halves, such that half of the scores are larger than the median, and the other half are smaller.

#### 4.5 CALCULATION OF THE MEDIAN

- If the data are an odd number of untied scores, e.g., 11, 13, 18, 19, 20, the median is the middle score when they are ranked; Md = 18.
- If the data are an even number of untied scores, e.g., 4, 9, 13, 14, the median is the point halfway between the two central values when the scores are ranked: Md = (9 + 13)/2 = 11.
- 3. If tied scores occur in the data, particularly at or near the median, a frequency tabulation of the scores will probably be necessary. Interpolation within a score class will often be necessary in such instances. For example, suppose that 36 scores ranging from 7.0 to 10.5 have the following ungrouped frequency distribution:

Score	Frequency	Cumulativ frequency
10.5	2	36
10.0	3	34
9.5	2	31
9.0	6	29
8.5	10 = 5 + 5	23
8.0	8)	13
7.5	4 13	5
7.0	i) ·	1
,	n = 36	

The mean is no more than the familiar "average." People often use the word "average" for any one of the three above-mentioned "measures of central tendency." The name "mean," being less ambiguous, is preferred.

You may begin to question this proliferation of measures of central tendency, especially when someone is holding you responsible for learning

them. However, each measure has characteristics that make it uniquely valuable.

The mode is easiest to calculate—it can be found literally at a plance. Moreover, in very large groups of scores it is a fairly stable measure of the center of the distribution. In many distributions of large numbers of measures taken in educational and psychological research, the mode is close to two other measures of central tendency, the median and the mean.

The median stands between the mode and the mean in computational effort, if the calculation is done by hand. This measure is obtained almost entirely by counting and can be easily calculated once the data have been arranged in numerical order. For a large number of scores, the data can first be arranged in a grouped frequency distribution (which is considerably easier than ranking the data), and then the median can be easily obtained. For purely descriptive purposes, classroom teachers and others working with small samples will find that percentile measures of central tendency and variability will serve them well enough.

The mean of a set of data involves the most arithmetic computation. The value of the mean is affected by the individual values of all of the scores in the set of data. The median and mode may not be affected by all of the values. For example, observe what happens to the values of the mean, median, and mode when the largest score in the following set is doubled:

	Mean	Median	Mode
Set 1: 1, 3, 3, 5, 6, 7, 8	3,2	5	3
Set 2: 1 3 3 5 6 7 16	42	5	3

The value of the mean is especially affected by what might be called "outliers," i.e., scores which lie far from the center of the group of scores. Whether this is an advantage depends upon the particular questions you are asking of the data.

# CALCULATION OF THE MEAN

The definition of the mean is so well known and the calculations by which it is found are so simple that you may wonder why its calculation deserves special attention. Part of the reason is that, traditionally, statistical calculations were performed by hand without the aid of machines. Under these circumstances, various methods of simplifying the operations in finding  $\vec{X}$  group of scores is that they have the group frequency distribution shown in the left portion of Table 4.2.

TABLE 4.2 ILLUSTRATION OF CALCULATION OF THE MEAN FROM A GROUPED FREQUENCY DISTRIBUTION

Score interval	f.	Midpoint of score interval	$f_i \cdot midpoint$
70-74	1	72	72
65-69	0	67	0
60-64	3	62	186
55-59	2	57	114
50-54	6	52	312
45-49	10	47	470
40-44	8	42	336
35-39	8 )	37	296
30-34	4	32	128
25-29	2 [	27	54
20-24	4	22	88
15-19	1	17	17
10-14	_1	12	12
	$n = \sum_{i=1}^{13} f_i = 50$	$\sum f_i$ (mi	idpoint) = 2085

Lacking better information and for the sake of simplicity, we assume that the scores in any score interval are evenly distributed along the interval. This assumption is usually false and for this reason the value we shall calculate is only an approximation (a rather close one) to the mean of the ungrouped data. [The degree of error introduced into the calculation of many statistics by this assumption was the subject of some theoretical work by Sheppard (e.g., see Keeping, 1962, p. 107) to which you may wish to refer at some point later in your statistical training.] Under the assumption, the sum of the f, scores in any score interval equals the product of f, and the midpoint of the interval. These products appear as the last column in Table 4.2. The grand sum of all of the scores in the group is approximately equal to the sum of these products. Thus, the mean of the ungrouped scores is approximately equal to this grand sum divided by n, i.e.,

$$\vec{X} \cong \frac{\sum f_i(\text{midpoint})}{n},$$
 (4.3)

and the summation extends over all of the score intervals.

For Table 4.2, the following approximation to the mean of the scores is obtained:

$$\bar{X} \simeq \frac{2085}{50} = 41.70.$$

If each score in a set whose mean is  $\bar{X}$  is multipled by a constant c, the mean of the resulting scores is  $c\bar{X}$ , because  $\sum_{i=1}^{n} cX_i/n = c\sum_{i=1}^{n} X_i/n = c\bar{X}$ .

A fourth property of the mean concerns the n deviation scores. The sum of the squared deviations of scores from their arithmetic mean is less than the sum of the squared deviations around any point other than X.

That is,  $(X_1 - \bar{X})^2 + (X_2 - \bar{X})^2 + \dots + (X_s - \bar{X})^2$  is smaller in value than  $(X_1 - b)^2 + (X_2 - b)^2 + \dots + (X_n - b)^2$ , where b is any number other than  $\bar{X}$ , the mean.\*

For example, the sum of the squared deviations of 0, 1, 1, 3, 5 around 2, their mean, is  $(0-2)^2+(1-2)^2+(1-2)^2+(3-2)^2+(5-2)^2=(-2)^2+(-1)^2+(-1)^2+(1)^2+(3)^2=16$ . The sum of the squared deviations of 0, 1, 1, 3, 5 around 1 is equal to 21, which is greater than 16.

#### 4.9 MEAN, MEDIAN, AND MODE OF COMBINED GROUPS

We might know the means, medians, and modes of three separate classrooms in a school and wish to find the same measures for all three classrooms combined. This will be a simple matter in the case of the mean, but for the median and mode it will be necessary to go back to the original data and make new calculations. The ease with which the mean of the combined groups is found reveals one of the advantages of definite summary statistics in terms of simple algebraic operations, such as adding and dividing, and having every score in a group exert an influence on the measure of central tendency. The median and mode are found by the operations of ranking and inspecting the data, respectively.

The means and frequencies for classrooms A, B, and C are as follows:

$$\overline{X}_A = 11.9$$
  $n_A = 24.$   $\overline{X}_B = 14.2$   $n_B = 30.$   $\overline{X}_C = 10.8$   $n_C = 28.$ 

\* Proof:

$$\begin{split} \sum_{i=1}^{n} \left[ (X_{i} - (\bar{X}_{i} + c))^{2} \right] &= \sum_{c=1}^{n} \left[ (X_{i} - \bar{X}_{i}) - c \right]^{2} \\ &= \sum_{c=1}^{n} (X_{i} - \bar{X}_{i})^{2} - 2c \sum_{i=1}^{n} (X_{i} - \bar{X}_{i}) + nc^{2} \approx \sum_{i=1}^{n} (X_{i} - \bar{X}_{i})^{2} + c^{2}, \\ \text{because } \sum_{i=1}^{n} (X_{i} - \bar{X}_{i}) = 0, c^{2} \geq 0, \text{ so} \\ &\sum_{c=1}^{n} (X_{i} - \bar{X}_{i})^{2} \leq \sum_{c=1}^{n} [X_{i} - (\bar{X}_{i} + c)]^{2} = 0. \end{split}$$

The total n of all three classes combined is equal to  $n = n_A + n_B + n_C = 82$ . The mean of the combined groups is simply the sum of all 82 scores divided by 82. The sum of the 24 scores in class A is simply  $n_A \cdot \tilde{X}_A = 24(11.9) = 285.6$ , because the mean is the sum of the scores divided by the number of scores, i.e.,  $\Sigma X = n\tilde{X}$ . Similarly for groups B and C, the sums of the scores are 30(14.2) = 426.0 and 28(10.8) = 302.4, respectively. 285.6 + 426.0 + 302.4 = 1014.0. If we combine all 82 scores and sum them, we will also obtain 1014.0. Thus, the mean of all three classes combined is  $(1014.0)[82 \approx 12.4$ . Symbolically, the mean of the combined groups is:

$$\overline{X}_{c} = \frac{n_{A}\overline{X}_{A} + n_{B}\overline{X}_{B} + n_{C}\overline{X}_{C}}{n_{A} + n_{B} + n_{C}}.$$
(4.4)

You should now be able to write the formula for the mean of four combined groups when you are given only the four means and numbers of scores per group.

Notice that if each group is based on the same number of frequencies,  $n_A = n_B = n_C = n$ , then Eq. (4.4) becomes

$$\bar{X} = \frac{n(\bar{X}_A + \bar{X}_B + \bar{X}_C)}{3n} = \frac{\bar{X}_A + \bar{X}_B + \bar{X}_C}{3}.$$
 (4.5)

This shows that if the three groups are the same size, the mean of the combined group is the same as the unweighted average of the three means. Of course, this is true for combining any number of means of equal-size groups.

Attempting to find the median or the mode of a combination of groups is a different matter, however. Suppose you know that group A has six scores and the mode is 17 and that group B also has six scores and a mode of 19. What is the mode of groups A and B combined? Would you guess that it is 18 = (17 + 19)/2? If you did you might be wrong, because groups A and B might look like this:


А	В	
15	15	
15	15	
17	17	
17	19	
17	19	
18	19	

When A and B are combined, the score of 15 occurs four times and is the mode of the combined groups. There was no way of knowing that this might happen when you were told only the modes for A and B individually. For both the mode and the median, you must have the original data in hand before you can find these measures of central tendency on combined groups.

#### 4.10 INTERPRETATION OF MODE, MEDIAN, AND MEAN

Each of the measures of central tendency we have presented has an interesting interpretation in terms of errors which result when a single statistic takes the place of each score in the group. The sense in which the mode is the most representative score or the score which best "takes the place of all of the scores" is fairly obvious. If we were forced to select one score to stand for every score in a group, the selected score would equal the score for which it stands the greatest number of times if it (the selected score) is the mode of the group. Or similarly, if we were being paid one dollar for each time we correctly guessed which score in a group would be selected by chance, we should make most money in the long run by always guessing the mode.

One interpretation of the median of a group is not so obvious. Suppose that the scores in a group are placed along the real-number line. (1, 3, 6, 7, 8) appear on the line below:



The Md indicates the median of the group, 6. The distance between 6 and 1 is 5 units; between 6 and 3, 3 units; between 6 and 6, 0 units; between 6 and 7, 1 unit; between 6 and 8, 2 units. The sum of these distances, 5+3+0+1+2=11, is smaller than would be the sum of these distances of the five points from any other point on the line. (Try it and see for yourself.) The median of a group of scores is that point on the number line such that the sum of the absolute (i.e., unsigned) distances of all scores in the group to that point is smaller than the sum of distances to any other point.\* If the median is taken in place of every score in the group, the least error results—provided "error" is defined as the sum of the absolute distances of each score to the score which will take its place.

The interpretation of the mean has already been noted. The mean of a group of scores is that point on the number line such that the sum of the squared distances of all scores to that point is smaller than the sum of the squared distances to any other point. If the mean is taken in place of every score in the group, the least error results—provided "error" is defined as the sum of the squared distances of each score to the score that will take its place.

<sup>\*</sup> For a proof of this property of the median, see Horst (1931).

# 4.11 CHOICE OF A MEASURE OF CENTRAL TENDENCY

Calculation of the mode, median, or mean is purely mechanical. Machines do it with far greater accuracy and speed than humans. However, the choice among these three measures and their interpretation may sometimes require judicious thought. Here are some considerations to keep in mind when you are faced with the choice:

- In small groups the mode can be quite unstable. The mode of the group (1, 1, 1, 3, 5, 7, 7, 8) is 1; but if one of the 1's is changed to 0 and the other to 2, the mode becomes 7.
- The median is not affected by the size of the "large" and "small" scores above or below it. For example, in a group of 50 scores the median will not change when the largest score is tripled.
- 3. The mean is influenced by the size of every score in the group. If any one score is changed by c units, X, will be changed in the same direction by cfu units. For example, if 100 is added to the third largest score in a group of 10, the mean of the group is increased by 10 units.
- 4. Some groups of scores simply do not "tend centrally" in any meaningful way, and it is often misleading to calculate one measure of central tendency. This is particularly true of groups of scores that have more than one mode. For example, an acquaintance of the authors' who is a researcher on curriculum development maintains that he can build achievement tests composed of eight multiple-choice items that can separate a group of students into those who have acquired the concept of adding two-digit numbers and those who have not. The "haves" will make scores of 6, 7, and 8; the "have-noxi" will make scores of 0, 1, and 2. Suppose a typical group of students produced scores yielding the histogram shown in Fig. 4.2.

The mean of the scores depicted in Fig. 4.2 is approximately 3.85, even though no person scored 3.85 or within two units of that




FIG. 4.2 Hatogram of scores on a 10-tem test of adding two-digit numbers (n = 51).




FIG. 4.3 A symmetrical, unimodal group of scores,

score. The median of the group is approximately 2.17, even though the score that is two ranks above the middle score is 61 Neither the mean nor the median depicts this group of scores well. Perhaps the best simple summary of the size of the scores is the statement that "the histogram is bimodal and U-shaped with one mode at 0 and the other at 8."

- 5. The central tendency of groups of scores which contain extreme values is probably best measured by the median if the scores are unimodal. As pointed out previously, each score in a group influences the mean. Thus, one extreme value can move the mean of the group far away from what would generally be regarded as the central region. For example, if nine persons have incomes ranging from \$4500 to \$5200, with an average of \$4900, and the tenth person's income is \$20,000, the average income of the group of 10 is \$6410. This figure does not do justice to either group, though it would be an impressive figure for the president of a small company (whose salary was \$20,000) to report as the average salary on his payroll. The median would be preferred as a measure of central tendency in this instance. Demographers, economists, and journalists often choose to report "median income" because they wish to avoid the problem just illustrated.
- 6. In unimodal groups of scores that are symmetrical (i.e., the half of the histogram below the mode is the mirror image of the half above the mode), the mean, median, and mode are equal. For an example, see Fig. 4.3. The frequency polygon shows that the mean, median, and mode are all 40.

Lack of perfect symmetry in the frequency polygon or histogram of a group of scores generally has a regular effect on the relationships among the mean, median, and mode. Suppose that a preponderance of the scores in some group lie above a peak in the frequency polygon as in Fig. 4.4.

In Fig. 4.4 the mode (Mo) equals 100, the median (Md) equals 104.6 and the mean (X) equals 105.98. If a preponderance of the scores lie lower than the peak of a frequency polygon, it would




FIG. 4.4 An asymmetrical frequency polygon illustrating the relationships among the mean, median, and mode.

generally be true that the mean would be the smallest number, the median would be next largest, and the mode would be the largest number.

7. An additional consideration which bears on one's choice of a measure of central tendency can be discussed only superficially with the concepts developed thus far. When the group of scores in hand is considered to be a sample from a much larger symmetrical group, the mean of the sample is probably closer to the center of the large group of scores than is either the median or the mode of the scores in hand. We shall return to this important point in Sec. 11.4.

The following anecdote summarizes many of the problems that arise in the use of measures of central tendency:

An improbable anecdote should make this problem of heterogeneity (i.e., that no measure of central tendency is an adequate description of all of the scores in a group) plain. Five men once sat near each other on a park bench. Two were vagrants, each with total worldly assets of 25 cents. The third was a workman whose bank account and other assets totaled \$2000. The fourth man had \$15,000 in various forms. The fifth was a multimillionaire with a net worth of \$5,000,000. Therefore, the modal assets of the group were 25 cents. This figure describes [the financial assets of] two of the persons perfectly, but is grossly inaccurate for the other three. The median figure of \$2000 does little justice to anyone except the workman. The mean, \$1,003,400.10, is not very satisfactory even for the multimultionaire. If we had to choose one measure of central tendency, perhaps it would be the mode, which describes 40 percent of this group accurately. But if told that "the modal assets of five persons sitting on a park bench are 25 cents," we would be likely to conclude that the total assets of the group are approximately \$1.25, which is more than five million dollars lower than the correct figure. Obviously, no measure of central tendency whatsoever is adequate for these "strange benchfellows," who simply do not "tend centrally.""

Quoted from page 73 of Julian C. Stanley, Measurement in Today's Schools, 4th Ed.,
 C 1964 Reprinted by permission of Premise-Hall, Inc., Englewood Cliffs, N.J.

Another escape from the contradictions in this anecdote is to point out that no summary statistic is needed for a group of five scores.

### 4.12 OTHER MEASURES OF CENTRAL TENDENCY

Numerous other ways of locating a "central value" in a group of scores exist. A few of these measures will be presented here despite the rarity of their appearance in the literature of educational research.

### The Geometric Mean

Undoubtedly you recall that  $\sqrt{4} = 2$  and that, more generally, if  $b = \sqrt{a}$ , then  $b^2 = a$ . Perhaps your memory is somewhat more vague on the general question of roots of numbers.  $\sqrt[3]{a}$  is read "the cube root" of the number a and is that number which when raised to the third power equals a, i.e., if  $\sqrt[3]{a} = b$ , then  $b \cdot b \cdot b = b^3 = a$ . The nth root of a, denoted  $\sqrt[3]{a}$ , is that number which when raised to the nth power (i.e., multiplied by itself n times) equals a. For example,  $\sqrt[4]{16} = 2$  because  $2 \cdot 2 \cdot 2 \cdot 2 = 2^4 = 16$ .

**Definition:** The geometric mean of the n positive numbers  $X_1, \ldots, X_n$  is given by

$$gm = \sqrt[n]{X_1 \cdot X_2 \cdot \ldots \cdot X_n}, \qquad (4.6)$$

i.e., the geometric mean of  $X_1, \ldots, X_n$  is the *n*th root of the grand product of all n of the X's.

The definitional formula for gm is not computationally convenient. If you find that you need to calculate gm for four, five, or more scores, refer to Senders (1958, p. 316).

It is necessary to restrict the definition of gm to positive numbers; if any single  $X_1$  were zero, the grand product of  $X_1, \ldots, X_n$  would also be zero and hence gm would equal zero even if zero was (by any other definition) far from a central value of the group.

The geometric mean is useful for describing the central location of rate-of-change scores (see Ferguson, 1959, p. 50).

### The Harmonic Mean

This measure of central tendency is sometimes applied when a group of

Definition: The harmonic mean of the positive scores  $X_1, \ldots, X_n$  is given by

$$hm = \frac{1}{[(1/X_1) + (1/X_2) + \dots + (1/X_n)]/n}$$
$$= \frac{n}{\sum_{i=1}^{n} 1/X_i}.$$

It is thus their reciprocals (where  $1/X_t$  is the reciprocal of  $X_t$ ). To find  $hm_t$  each score is divided into 1, the resulting reciprocals are summed, and this sum is divided into n.

The harmonic mean has very limited application. A contraharmonic mean is thought to have broader application than either the geometric or harmonic mean (see Senders, 1958, pp. 317-18); however, it is doubtful that any of these three measures is sufficiently familiar to most readers that the use of them in written reports would pass without considerable confusion.

### The Ratio of Means and the Mean of Ratios

An IQ score, as computed from most of the earlier intelligence tests, is 100 times the ratio of a person's mental age to his chronological age:  $\Omega=100(MA/CA)$ . At first sight, it might appear that the average IQ of a group of persons could be found by dividing the average mental age of the group by the average chronological age and multiplying by 100, i.e., that  $\overline{\Omega}=100(\overline{MA/CA})$ . However, this is not generally true.

Only under very special conditions will it be true that

$$\frac{1}{n}\sum_{i}\frac{X_{i}}{Y_{i}}=\frac{\overline{X}_{i}}{\overline{Y}_{i}},$$

i.e., that the mean of the ratios is the ratio of means (see Stanley, 1957b). This seems like such a simple fact, but in a slightly more complicated form the error of assuming that the mean of ratios is the ratio of means sneaks into statistical literature (e.g., see Winer, 1962, pp. 61 and 119; Ferguson, 1959, p. 288).

### PROBLEMS AND EXERCISES

Find the mean, median, and mode of the following set of scores: 1.2, 1.5, 1.6, 2.1, 2.4, 2.4, 2.7, 2.8, 3.0, 3.0, 3.0, 3.1, 3.1, 3.1, 3.4.


- 2. Suppose the number 0.5 is added to each of the 15 scores in Prob. 1 above. What will be the value of the mean and median of these 15 augmented scores?
- Find the mean and median of the 100 scores in the following grouped frequency distribution:

Score interval	Frequency
20-22	8
17-19	14
14-16	41
11-13	26
8-10	7
5-7	4
	n = 100

- 4. Suppose that each of the 100 scores in the grouped frequency distribution in Prob. 3 is multiplied by 3. What would be the values of the mean and median of the 100 resulting scores?
- 5. Find the mean and median in the following grouped frequency distribution:

Score interval	Frequency
20-24	2
15-19	11
10-14	17
5-9	13
0-4	9
(~5)-(-1)	7
, , , .	n = 59

- 6. Group A contains 10 scores, the mean and median of which are 14.5 and 13, respectively. Group B contains 20 scores, the mean and median of which are 12.7 and 10, respectively. What are the mean and median of the 30 scores obtained from combining Groups A and B?
- The seven members of the Sunday Afternoon Picnic Society (SAPS) live along a straight stretch of Highway 101. Their homes are positioned along the highway as follows:



The cost of gas—3.5 cents per mile—for the travel of all members to the Sunday outing is taken out of the club treasury. Since any point along Highway 101 is a fine place for a picnic, where along the road should the members hold their picnic so as to spend the minimum amount of money for travel?

8. Find the mean SAT-V score from the n = 903 scores for men in Prob. 4 at the end of Chapter 3. (Hint: a quick way to find the mean is to multiply the midpoint of each score interval by the relative frequency of the scores in that interval, and then add up these products for all 12 intervals.)

### MFASURES OF **VARIABILITY**

### 5.1 INTRODUCTION

Measures of central tendency tell us about the concentration of a group of scores on a number scale. A particular measure of central tendency gives a score that "represents" in one of several senses all of the scores in the group. This process disregards the differences that exist among separate scores. Other descriptive statistics are required to measure the variation of scores within a group. In this chapter, several statistics that measure, in different ways, the variability (heterogeneity, dispersion, scatter) in a group of scores will be presented and discussed.

Later in this text, you should begin to see that some of the most important functions of statistics relate to procedures by which variability, which is in a sense uncertainty, can be reduced, explained, or accounted for. The whole scientific enterprise is concerned with notions of variability. When much unexplained variability exists, predictions can never be very accurate. But when explanations can be fashioned for why people or things differ, uncertainty can be reduced and portions of variability can be removed.

MEASURES OF VARIABILITY

For example, if nothing is known about why people differ in intelligence, one is faced with great uncertainty when attempting to predict intelligence; some people would appear "bright" and others "dull" and no one would know why. However, if it is known that heredity and environment produce quantifiable influences on measured IQ, then knowing a child's ancestry and early upbringing would permit a more accurate prediction of his adult intelligence. In other words, the variability of IQ's for persons with like heredity and environment is less than it is for people in general. But before we can continue with these loftier matters, we must learn about conventional indices of variability.

#### 5.2 THE RANGE

The range simply measures the full distance along the number scale over which the scores vary. Because slightly different definitions of the range have been given in the past, it will be necessary to distinguish two types of range: the inclusive and the exclusive range.

Definition: The exclusive range is the difference between the largest and smallest scores in a group.

For example, the exclusive range of the scores 0, 2, 3, 3, 5, and 8 is 8-0=8. The scores -0.2, 0.4, 0.8, and 1.6 have an exclusive range of 1.6 -(-0.2)=1.8.

Definition: The inclusive range is the difference between the upper real limit of the interval containing the largest score and the lower real limit of the interval containing the smallest score.

For example, the heights of six boys are measured to the nearest inch. height of the shortest boy is somewhere between 58.5°, and 66°. The actual real limit being 38.5°. The upper real limit of the interval containing the largest score is 66.5°. Thus, the inclusive range equals 66.5° — 58.5° = 8°, which is 1 larger than 66 — 59 (see Fig. 5.1).




FIG. 5.1 Illustration of the inclusive and the exclusive range.

The exclusive range is the distance between the smallest and largest reported scores in a group, and is thus likely to exclude an actual score that lies above the largest or below the smallest reported score (see Sec. 2.3 for definitions of actual and reported scores).

The inclusive range is sufficiently large to include all actual scores as well as all reported scores.

In the future, if we refer to "the range" without specifying inclusive or exclusive, the point being made will be equally valid for either. Although the meaning of the range as a measure of variability is quite clear, it has certain drawbacks. Because the range is determined by just two scores in the group, it ignores the spread of all scores except the largest and smallest. For example, if 100 scores are spread evenly from 1 through 10, the inclusive range is 10.5 – 0.5 = 10. But if one score is at 1 and one score is at 10 and the remaining 98 scores are at 5, the inclusive range is still equal to 10. For almost any purpose, these two types of heterogeneity (one in which 100 scores are uniformly distributed over a ten-unit interval and the other in which all but two of the 100 scores are clustered at the same point) have different meaning; but they cannot be distinguished by merely inspecting the range. The range is by far the crudest measure of variability commonly employed.

### 5.3 D, THE 90TH-TO-10TH PERCENTILE RANGE

A second measure of variability is D, the range between the 10th and 90th percentiles in a group of scores. It was defined by an educational statistician, Truman Kelley (1921), as

$$D = P_{90} - P_{10}, (5.1)$$

D is somewhat more stable than the range because it is directly affected by a greater number of scores. It is easier to compute than other measures of variability we shall meet later in this chapter. Neither of these advantages has been great enough to make D a popular measure of variability. It is seldom used.

#### 5.4 THE SEMI-INTERQUARTILE RANGE

In Sec. 3.2 we considered the three quartiles of a distribution of scores:  $Q_1$ , the point on the scale below which 25% of the scores lie,  $Q_2$  (the median), and  $Q_3$ , the point above which 25% of the scores lie. The distance between

the first and third quartiles of a group of scores, i.e.,  $Q_3 - Q_1$ , is called the

interquartile range. The semi-interquartile range is half this distance.

Using Q to denote the semi-interquartile range, we have the following:

Definition: The semi-interquartile range Q is half the distance between the third and first quartiles, i.e.,

$$Q = \frac{Q_3 - Q_1}{2}.$$
 (5.2)

Q is an easily obtained and useful measure of variability. For deequive purposes it is superior to the range on any criterion except computational simplicity. If two groups of scores have the same value of Q, they are much more likely to possess similar patterns of heterogeneity than are two groups with the same range.

In distributions that are nearly symmetrical around the mean or median, Q can be used to reconstruct the score limits between which approximately 50% of the scores are contained. If it is known that 250 scores which are approximately symmetrically distributed around a median of 63 have a semi-interquartile range of 11, then approximately 50% (125) of the scores lie between

$$Md - Q = 63 - 11 = 52$$

and

$$Md + Q = 63 + 11 = 74$$
.

If the distribution of scores is very asymmetrical around the median, then as many as 70% of the scores might lie within the range Md-Q to Md+Q. The symmetric and asymmetric cases and the use of Q are illustrated in Fig. 5.2.




FIG. 5.2 Interpretation of Q. Exactly 50% of the scores lie between Md-Q and Md+Q in the symmetric distribution on the left. 52.5% of the scores he between Md-Q and Md+Q in the somewhat asymmetric distribution on the right.

VEIV . /

The range, the semi-interquartile range Q, and  $D = P_{90} - P_{10}$  are three measures of dispersion, scatter, heterogeneity, or variation met thus far. Each increases in value as the set of scores on which it is computed shows greater dispersion—less homogeneity. Note that, as with the mode and median, calculation of these three measures does not involve every individual score in a group of scores. We shall now encounter a fourth measure in the calculation of which, like the mean, every score is utilized.

Deviation scores, scores of the form  $X_i - \bar{X}_i$ , reflect something about the variation in a set of scores. A set of scores with great heterogeneity will have some large deviation scores. What would the deviation scores be if all the scores in the set were 9? The mean would be 9, hence every deviation score would be 9 - 9 = 0. In the most homogeneous set of scores that it is possible to achieve, all the deviation scores equal zero. Some combination of the deviation scores might be a useful measure of variation.

If we were to sum all of the deviation scores, would that sum reflect the variation in the original scores? No, since this sum is always exactly zero:  $\sum_{i=1}^{n} (X_i - \bar{X}) = 0$ . To overcome this we could square each deviation score and sum the resulting squared scores. Hence, for a given set of scores a measure of the form

$$\sum_{i=1}^{n} (X_i - \bar{X})^2 = (X_1 - \bar{X})^2 + \ldots + (X_n - \bar{X})^2$$

will be large when the scores are heterogeneous and small when they are homogeneous. We would not have had to square the deviation scores to get rid of their signs; we could have simply regarded all of them as positive (i.e., taken their absolute value). This method would have led to a different measure of variation called the mean deviation, which you will encounter in Sec. 5.9. The value of the above expression also depends upon how many scores are considered. The larger n, the larger the sum tends to be. This is a limitation if one wishes to compare the variability in two sets that differ in numbers of scores. This limitation is overcome by dividing the expression by n-1. The resulting measure of variability is called the variance (denoted by  $s_2^n$ ) and has the formula

$$s_x^2 = \frac{\sum_{i=1}^{n} (X_i - \bar{X})^2}{n-1}.$$
 (5.3)

Why did we divide by n-1 instead of simply n? A satisfactory answer cannot be given at this point because as yet in this chapter we have not developed the concepts necessary to give the explanation much meaning. You will learn in Sec. 12.4 why n-1 was chosen instead of n.

We shall use the following data to illustrate the calculations of the variance of a set of six scores:

Score	Score - Mean	(Score - Mean)2
	1-2=-1	1
3	3-2=1	1
3	3 - 2 = 1	1
Ō	0-2=-2	4
4	4 - 2 = 2	4
1	1 - 2 = -1	1
12	ō	12
	$s_z^2 = \frac{12}{6-1} = \frac{12}{5}$	= 2.4

Suppose we had a second, more heterogeneous set of six scores with the same mean, 2. These data, and the calculations, are as follows:

In the above examples,  $s_{s}^{2}$  was easily calculated because each score and the mean were whole numbers. Computation of  $s_{s}^{2}$  as above would be tedious, however, if the mean were 17.697, for example. For this reason we seek, by algebraic manipulation, an expression for  $s_{s}^{2}$  that is computationally simpler in such instances.

$$s_{2}^{2} = \frac{\sum_{i=1}^{n} (X_{i} - \widetilde{X})^{2}}{n-1} = \frac{\sum_{i=1}^{n} (X_{i}^{2} - 2\widetilde{X}X_{i} + \widetilde{X}_{i}^{2})}{n-1} = \frac{\sum_{i=1}^{n} X_{i}^{2} - 2\widetilde{X}\sum_{i=1}^{n} X_{i} + \sum_{i=1}^{n} \widetilde{X}_{i}^{2}}{n-1}.$$

If we recall that  $\sum X = n\dot{X}$ , we can write the above expression as follows:

$$\frac{\sum_{i=1}^{n} X_{i}^{2} - 2n\overline{X}_{i}^{2} + n\overline{X}_{i}^{2}}{n-1} = \frac{\sum_{i=1}^{n} X_{i}^{2} - n\overline{X}_{i}^{2}}{n-1}.$$

Since the square of the mean is the square of the sum of all the scores divided by the square of n, we have

$$s_x^2 = \frac{\sum_{i=1}^{n} X_i^2 - \left| \left( \sum_{i=1}^{n} X_i \right)^2 / n}{n-1}.$$
 (5.4)

We can also multiply both sides of Eq. (5.4) by a fancy form of the number I, namely n/n, to obtain another formula for  $s_2^n$  that does not contain the mean:

$$s_{z}^{2} = \frac{n \sum_{i=1}^{n} X_{i}^{2} - \left(\sum_{i=1}^{n} X_{i}\right)^{2}}{n(n-1)}.$$

### 5.6 CALCULATION OF THE VARIANCE s<sup>2</sup>

The calculation of  $s_2^2$  by means of Eq. (5.4) will be illustrated on the six scores used in the first example above: 1, 3, 3, 0, 4, 1.

X	$X^2$	Final calculations
$ \begin{array}{c} 1\\ 3\\ 3\\ 0\\ 4\\ 1\\ \sum_{i=1}^{6} X_i = \overline{12} \end{array} $	$ \begin{array}{c} 1 \\ 9 \\ 9 \\ 0 \\ 16 \\ \frac{1}{1} \end{array} $	$\sum_{1}^{n} X^{2} - \frac{\left(\sum_{1}^{n} X\right)^{2}}{n} = 36 - \frac{12^{2}}{6} = 36 - \frac{144}{6}$ $= 36 - 24 = 12$ $s_{x}^{2} = \frac{12}{6 - 1} = \frac{12}{5} = 2.4$

When one or more of the possible values a variable can assume occurs more than once in a group of scores, a simplification of the calculation of  $x^2$  is possible, as was the case in the calculation of  $X^2$  (see Sec. 4.7). In the above illustration, the number 3 appeared twice. The amount contributed to the sum of the scores  $(\sum X)$  by the two threes was 3 + 3 = 2(3). The amount

contributed to the sum of the squared scores  $(\sum X^2)$  by the two threes was 9+9=2(9). If the value  $X_i$  has a frequency in the group of  $f_i$ , then this value will contibute  $f_iX_i$  to the sum of the scores and  $f_iX_i^2$  to the sum of the squared scores. Consequently, in finding  $\sum X^2$ , for example, it is not necessary to square the value  $X_i$  each time it occurs. Instead,  $X_i^2$  is found once and multiplied by the number of times  $X_i$  occurs in the group.

Using the same data upon which the calculation of  $\bar{X}$  was illustrated

in Sec. 4.7, we shall find  $s_x^2$  by this shortened method, where

$$s_z^2 = \frac{\sum_{i=1}^k f_i X_i^2 - \left[ \left( \sum_{i=1}^k f_i X_i^2 \right) / n \right]}{n-1},$$

k is the number of different scores, and  $\sum_{i=1}^{k} f_i = n$ .

	count	Intermediate calculations	
Original scores	$X X_i f$	fX fX <sup>1</sup>	Final calculations
2 6 10 3 6 10 3 6 11 5 8 11 5 8 11 5 9 15 5 9 18	2 4 1 3 9 2 5 25 4 6 36 3 8 64 2 9 81 2 10 100 2 11 121 3 15 225 1 18 324 1	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	$\sum_{i=1}^{3} X_i = \sum_{i=1}^{3n} f_i X_i = 166.$ $\sum X_i^4 = \sum f_i X_i^2 = 1632.$ $\sum X_i^4 = \frac{(\sum X_i)^3}{n}$ $= 1632 - 1312.19 = 319.81.$ $z_n^4 = \frac{319.81}{20} = 15.995.$
		2 july 100 2 july = 103	

### 5.7 THE STANDARD DEVIATION :

A measure of variability closely related to the variance is the standard deviation. The standard deviation, denoted by s, is defined as the positive square root of the variance. To find s, one first finds s<sup>2</sup> and then finds the square root of s<sup>2</sup>.

$$s_2 = \sqrt{\frac{\sum X^2 - [(\sum X)^2/n]}{n-1}}$$
(5.5)

If the variance is 16, what is the standard deviation?

$$s_z = \sqrt{s_z^2} = \sqrt{16} = 4$$
.

The standard deviation is often a useful measure of variation because in many distributions of scores we know approximately what percent of the scores lie within one, two, three, or more standard deviations of the mean. For example, we may know that 70% of the scores lie between X-s, and X+s.

### 5.8 SOME PROPERTIES OF THE VARIANCE

Suppose we added a constant number to every score in a set of scores. How would the variance of the scores be affected? In Sec. 5.6 we found that the scores 1, 3, 3, 0, 4, 1 have variance equal to 2.4. Let's add 2 to each score and then calculate st:

(Original score + 2) - Mean		(Deviation score)
3	-4 = -1	ı
5	-4 = 1	1
5	-4 = 1	1
2	-4 = -2	4
6	-4 = 2	4
3	-4 = -1	1
$Sum \approx \overline{24}$	Sum of deviations $= \overline{0}$	12
Mean = 4		
	$s_x^2 = \frac{12}{6-1} = \frac{12}{5} = 2.4$	

Adding 2 to each score did not change the value of  $s_s^2$ . In general, adding a constant c to each score in a group will not change the variance (nor the standard deviation) of the scores:

$$\sum_{i=1}^{n} \frac{\left[ (X_i + c) - \left[ \sum_{i=1}^{n} (X_i + c) / n \right] \right]^2}{n-1} = \sum_{i=1}^{n} \frac{\left[ X_i + c - (\sum X_i / n) - (nc/n) \right]^2}{n-1}$$

$$= \frac{\sum_{i=1}^{n} (X_i + c - \overline{X}_i - c)^2}{n-1}$$

$$= \sum_{i=1}^{n} (X_i - \overline{X}_i)^2$$

$$= \sum_{i=1}^{n} (X_i - \overline{X}_i)^2$$

$$= \sum_{i=1}^{n} (X_i - \overline{X}_i)^2$$

What would happen to  $s_2^2$  if each score were multiplied by a constant, say 2?

Note also 9.6 equals  $(2^a) \cdot (2.4)$  In general, multiplying each score by a constant c makes the variance of the resulting scores equal to  $c^2s_a^2$ :

$$\sum_{i=1}^{n} \left[ \frac{cX_i - \left(\sum_{i=1}^{n} cX_i / n\right)}{n-1} \right]^2 = \sum_{i=1}^{n} \left[ \frac{cX_i - \left(c\sum_{i=1}^{n} X_i / n\right)}{n-1} \right]^2 = \sum_{i=1}^{n} \left[ \frac{c(X_i - X_i)^2}{n-1} \right]^2 = \frac{\sum_{i=1}^{n} \left[ \frac{c(X_i - X_i)^2}{n-1} \right]^2}{n-1}$$

 $\sum_{i=1}^{n} c^{2}(X_{i} - \overline{X})^{2} = c^{2}\sum_{i=1}^{n}(X_{i} - \overline{X})^{2} = c^{2}\sum_{$ 

two separate groups was found to be a simple weighted average of the means of the two groups (see Sec. 4.9). The comparable situation for variances is more complicated. It will be seen that the variance of the set of scores formed by pooling scores from groups a and b depends on both the variances and means of the two groups. Notice that if group a comprises the scores 3, 3, 3 and 3 and group b is 6, 6, 6, 6, then the variance of groups a and b combined (3, 3, 3, 3, 6, 6, 6) is not zero even though a = a = a = 0.

Suppose that a and b denote two separate sets of scores:

	Group a	Group b
Group size Mean	n <sub>a</sub>	n,
Variance	53 5	X.,

The variance of the group of  $n_a + n_b$  scores formed by combining groups a and b is

$$s^{2} = \frac{\left[ (n_{a} - 1)s_{a}^{2} + (n_{b} - 1)s_{b}^{2} + n_{b}(\overline{X}_{a} - \overline{X}_{a})^{2} + n_{b}(\overline{X}_{b} - \overline{X}_{a})^{2} \right]}{n_{a} + n_{b} - 1}, (5.6)$$

Ġ

where

$$\overline{X}_{..} = \frac{n_a \overline{X}_{.a} + n_b \overline{X}_{.b}}{n_. + n_b}$$

### 5.9 THE MEAN DEVIATION

An additional measure of variability, the mean (or average) deviation is somewhat easier to calculate than the standard deviation but is less useful. The deviation of each score in a group from the mean is denoted by  $\vec{X}_t - \vec{X}$ . The collection of all n of these deviations is descriptive of the amount of variability in the original scores. As we saw in Sec. 4.8, however, the sum of these positive and negative deviations is not in the least descriptive of the total amount of variability in the group of scores because it is always precisely zero. If the deviations are regarded as distances of the scores from  $\vec{X}$  without regard to sign, the sum of these distances is descriptive of the variability in the scores.

The distance of each  $X_i$  from  $\overline{X}_i$  is found by a process known as taking the absolute value of 4.65 is denoted [4.65] and equals 4.65. In fact the absolute value of any positive number is that positive number itself. Thus |2| = 2, and |105| = 105. The absolute value of any negative number is found by changing the minus sign to a plus sign: |-3| = 3; |-1.69| = 1.69. Finally, |01 = 0.

$$|5| \approx 5$$
,  
 $|0| \approx 0$ ,  
 $|-5| \approx 5$ .

Another way to think of the process of taking absolute values—though one would be foolish to do it this way in calculations—is that  $|a| = \sqrt{a^2}$ , i.e., the absolute value of a number is the positive square root of the square of the number!

The distance of the score  $X_i$  from the mean is given by  $|X_i - \overline{X}_i|$ . The average of the n distances of the scores from their mean is called the mean deviation, MD. (Do not confuse MD, the mean deviation, with Md, the median.)

$$MD = \frac{\sum_{i=1}^{n} |X_i - \bar{X}_i|}{n}.$$

There is no simpler expression for the mean deviation that might be used for purposes of calculation. An illustration of the calculation of MD appears in Table 5.2. The mean deviation has not often been used as a measure of

TABLE 52 ILLUSTRATION OF THE CALCULATION OF THE MEAN DEVIATION

Scores	X - X	X - X	Final calculations
10	10 - 12 = -2	2	$MD = \frac{\sum  X_i - \bar{X} }{2}$
12	12 - 12 = 0	ō	$MD = \frac{2 \cdot 10^{-11}}{2}$
13	13 12 1	i	. "
10	$10 - 12 \Rightarrow -2$	•	= = 1.60
15	15 - 12 = 3	3	5
$\sum_{i=1}^{n} X_i = 60$ $X = 12$		$\sum_{i=1}^{n}  X_i - \overline{X}  = 8$	
X = 12		<u></u> 1	

variability, even though it is easily calculated and has a logical simplicity. One reason for this is that the mean deviation does not have a theoretical undergirding comparable to that of the variance, for example. The mathematical statistician finds that the process of "taking absolute values" presents special difficulties for certain mathematical derivations. The mean deviation has been defined and discussed at this point, however, because of the role it will play later in this text in a novel approach to testing hypotheses about variabilities in populations.

### 5.10 STANDARD SCORES

It is frequently desirable to describe the position of a score in a set of scores by measuring its deviation from the mean of all scores in standard deviation units. For example, a particular set of 100 scores has a mean of 18.75 and a standard deviation of 2.60. If you know only that a certain one of these 100 persons has a score of 20, his relative position in the collection of 100 scores is not immediately apparent. It can be determined by a series of calculations that 20 lies 1.25 units (20 – 18.75 = 1.25) above the mean and that this distance is 1.25/2.60 = 0.48 standard deviation units. Thus

For any arbitrary score  $X_i$  in the set of 100 scores, the deviation of the score from the mean measured in standard deviation units is given by

$$\frac{X_1 - 18.75}{2.60}$$
.

All 100 such original scores could be so transformed. What would be the mean and standard deviation of the resulting 100 transformed scores?

The mean of the 100 values of Y = 10.75;

The mean of the 100 values of X-18.75 is equal to the mean of the score subtracting X scores minus 18.75, since subtracting a constant from each X is 18.75, the mean of X-18.75 must be zero. If X-18.75 has a mean

Aside from being a convenient means of communicating the position of a person's score relative to the mean and measured in standard deviation units, z scores are a step toward transforming a set of X scores to an arbitrary scale with a convenient mean and standard deviation. For some purposes, for example, the z scores themselves may not be convenient. The negative scores might be bothersome, and a collection of z scores will undoubtedly be full of decimal numbers. A transformation of the z scores an eliminate these minor problems.

Since the nz scores have mean zero and standard deviation 1, we know that cz—formed by multiplying each z score by the constant c—will have a standard deviation of |c|, and cz + d will have a mean of

$$cz + d = c(0) + d = d.$$

For example, a set of 250 X scores has mean 79.65 and standard deviation 5.71. Suppose we wish to transform these scores so that their mean is 50 and their standard deviation is 10. We could first transform them to x scores by forming the 250 values of z = (X - 79.53)5.71. Next, we could calculate 10z + 50 for each of the 250 z scores. The resulting set of 250 transformed scores has a mean of 50 and a standard deviation of 10. A person with a transformed score of 59 lies (59 - 50)/10 = 9/10 of a standard deviation above the mean of the 250 scores; furthermore, we know that this same person's X score is approximately 79.65 + 0.9(5.71) = 85. The transformed score of 59 is more immediately informative than the X score of 85.

There are many scales of measurement (arbitrary means and standard scores can be placed along any such scale, i.e., they can be given a desired mean and standard deviation, merely by letting c be the desired standard deviation and d the desired mean in the expression cz + d. Intelligence-test scores are often transformed to a scale with mean 100 and standard deviation 15 or 16. T scores formed by 10z + 50 find wide application. These and other popular scales are shown in Fig. 6.5 in Chapter 6.

#### 5.11 SKEWNESS

The degree of asymmetry of a frequency distribution is one of its more important properties. Exactly symmetrical frequency polygons and histograms almost never occur with real data. The degree to which the frequency distribution of a group of scores is asymmetrical is its skewness. The nature and extent of asymmetry will be apparent if a frequency polygon or histogram is observed, but such observation is not always possible or convenient. Consequently, various summary statistics measuring the type and degree of asymmetry of a group of scores have been devised.

The best measure of the skewness of a set of scores has the formula

skewness = 
$$\frac{\sum_{i=1}^{n} (X_i - \bar{X}_i)^2 / n}{s_*^2}.$$
 (5.7)

As was seen in Sec. 5.10, it is standard practice to denote the directed distance which a score lies from the mean of the group it is in, measured in units of the standard deviation of that group, by z.; i.e.,

$$z_{z} = \frac{X_{i} - \bar{X}_{i}}{s_{-}}.$$

If we recognize that

$$\frac{(X_i - \overline{X})^3}{s_x^2} = \left[\frac{(X_i - \overline{X})}{s_x}\right]^3 \approx z_x^3$$

then the measure of skewness in (5.7) becomes

skewness 
$$=\frac{\sum_{i=1}^{n} z_{2i}^{3}}{n} = \overline{z_{i}^{3}}.$$
 (5.8)

Thus, one measure of skewness is simply the average of the z scores which have been raised to the third power. (In mathematical statistical work this measure of skewness is denoted by  $\sqrt{\beta_1}$ . The measure is due to Karl Pearson and its properties have been widely studied.)

Suppose the skewness of the two distributions in Fig. 5.3 is being measured. The mean of the scores for distribution A in Fig. 5.3 is about 16.




FIG. 5.3 Two skewed frequency distributions.

When the z scores for A are formed there will be some very large positive ones (because the largest X score is 22, which is 6 units above the mean) that will be larger in absolute value than any of the negative ones (the smallest

X score, 13, lies only 3 units below the mean). Now the algebraic sign of a number remains the same when it is cubed;  $(-2)^3 = -8$ . Hence, for distribution A, the contribution of the cubed negative z scores to  $\sum z_1^2/n$  will be less than the contribution of the cubes of those large positive z scores. Consequently, the value of  $\sum z_n^3/n$  in Eq. (5.8) will be large and positive. We say that distribution A is positively skewed because its measure of skewness is positive. A positively skewed frequency distribution has scores that extend further above the mean than the small scores extend below it.

Distribution B in Fig. 5.3 is negatively skewed. The value of  $\sum z_x^3/n$ for distribution B is negative. Try to convince yourself that this is true. Distribution A is more markedly positively skewed than distribution B is negatively skewed.

In a symmetric distribution the value of the measure of skewness in Eq. (5.8) is zero. This is true because exact symmetry implies that every negative z score can be paired with a positive z score of equal size. Since the cubes of negative scores are still negative, the sum of the cubes of each pair of positive and negative z scores is zero.

Unfortunately, the value of  $\sum z_1^2/n$  is laborious to obtain, even on a desk calculator, for any sizable group of scores. A quick method for measuring the skewness of a distribution grows out of the fact that the mean is more affected by extreme scores than the median. In the chapter on measures of central tendency we saw that in unimodal positively skewed distributions the mean is larger than the median, which is in turn larger than the mode. In negatively skewed distributions, the mean is smaller than the median, which is in turn smaller than the mode. This suggests that the position of the mean relative to the median tells something about the skewness of the distribution. This is indeed so for reasonably large groups of scores, say n of 50 or greater. A simple measure of skewness that makes use of these facts has the following definition:

skewness = 
$$\frac{3(\bar{X} - Md)}{s_*}$$
. (5.9)

In words, the skewness of a distribution can be measured by taking three times the difference between the mean and median, divided by the standard deviation. The values of Eq. (5.9) will generally range between -3 and +3. When a distribution is symmetric, Eq. (5.9) will be zero. The measure of skewness in Eq. (5.9) can be used to compare the skewnesses of different distributions because division by  $s_x$  has made the measure independent of the variability of the distribution.

### 5.12 KURTOSIS

We have seen how statisticians describe three properties or features of groups of scores. These three properties are central tendency, variability, and FIG. 5.4 "Peaked," "flat," and "mesokurtie" curves (A, B, and C, respectively).



symmetry. A fourth property completes the set of features of distributions of scores that are generally of interest in analyzing data. One may wish to know something about how peaked or flat a frequency polygon or histogram is. Kurtosis is a Greek word and refers to the quality of "peakedness" of a curve. (Karl Pearson is credited with formalizing the concept of kurtosis in statistics and proposing a method of measuring it.)

In Fig. 5.4 three curves differing in "peakedness" or kurtosis appear. The first (A) is quite peaked; such a curve is called leptokurite. (The prefix "lepto" means "slender" or "narrow.") The second curve (B) is relatively flat; such curves are called platykurite. (The prefix "platy" means "flat" or "broad.") The "peakedness" or degree of kurtosis of the third curve (C) is a standard against which the kurtosis of other curves is measured. The third curve in Fig. 5.4 is the normal curve, which will be discussed at length in Chapter 6, and is said to be mesokurite, "meso" meaning "intermediate."

We shall now learn how the statistician measures the kurtosis of a curve. First it is necessary to point out, however, that the concept of kurtosis applies only to unimodal distributions and concerns the steepness of the curve in the vicinity of the single mode. (If a distribution has two modes, it would be acceptable to talk about the kurtosis of the curve in the vicinity of each mode.)

The customary measure of kurtosis has the following definitional formula:

kurtosis = 
$$\frac{\sum_{i=1}^{n} (X_i - \bar{X}_i)^4 / n}{s_i^4}.$$
 (5.10)

If we recognize that  $(X_t - \bar{X}_t)^4/s_x^4$  is simply

$$\left\lceil \frac{(X_i - \bar{X}_i)}{5} \right\rceil^4 = z_{rr}^4$$

then we see that a measure of kurtosis is given by the formula

kurtosis = 
$$\frac{\sum_{i=1}^{n} z_{x_i}^4}{z_i^4} = \frac{z_i^4}{z_i^4}$$
. (5.11)

That is, kurtosis is measured by taking the average of the fourth powers of the x scores. The relationships between the size of the kurtosis statistic and the "peakedness" of the distribution on which it is calculated are recorded in Table 5.3.

TABLE 5.3 RELATIONSHIP OF THE VALUE OF THE KURTOSIS STATISTIC TO

Nature of distribution	Description of "peakedness"	Value of kurtosis statistic (Eq. 5.11)
Normal, e.g., curve C in Fig. 5.4	Mesokurtic	3
Peaked, e.g., curve A in Fig. 5.4	Leptokurtic	Greater than 3 (can become very large)
Flat, e.g., curve B in Fig. 5.4	Platykurtic	Less than 3 (must be zero or greater)

#### PROBLEMS AND EXERCISES

Calculate the inclusive range, variance, standard deviation, and mean deviation
of the following set of scores:

102	112	116
106	114	119
111	115	120
112	115	122

(Hint: To simplify calculations, first subtract 100 from all scores; this will not change the values of any of the measures of variability.)

Find the semi-interquartile range for the IQ scores in the following grouped frequency distribution;

Score interval	Frequency	Cumulative frequent
150-159	5	100
140-149	7	95
130-139	9	88
120-129	12	79
110-119	17	67
100-109	21	
90-99	12	50
80-89	8	29
70-79	6	17
60-69	1	9
50-59	2	3
	_	2

The median of the above set of scores is 109.5. Determine the proportion of scores that lie in the range determined by subtracting and adding Q to the median, i.e., what proportion of the scores lie between  $109.5 \pm Q$ ?

 Find s<sub>2</sub> and s<sub>2</sub> for the following grade-placement scores arranged in an ungrouped frequency distribution:

Score	Frequenc
6.9	2
6.8	4
6.7	5
6.6	9
6.5	14
6.4	10
6.3	6
6.2	3
6.1	2
60	

(Hint: Subtract 6.0 from each of the ten score values to simplify calculations.)

- Indicate whether each of the following distributions of measures is probably negatively or positively skewed:
  - a, ages of students in U.S. universities;
  - b. numbers of children in U.S. families;
  - c. population of cities in the United States;
  - d. ages at death of females in the United States.

5.	Group A	Group
	13	28
	11	26
	10	25
	0	24

The variances of both groups A and B equal 5. Will the variance of the ten scores formed by pooling together groups A and B be less than, greater than, or equal to 5?

 For the 290 students taking a 50 item social studies achievement test the mean is 32.50 and the standard deviation is 4.80. Find the z scores corresponding to the following:

	Test scores X	z score
a.	28	-0.94
b.	36	
c.	45	
d.	20	

Two vocabulary tests were administered to the 40 students in a beginning French
class. The means, the standard deviations on the two tests, and the raw scores
obtained by students 4 and B were as follows:

	Х.	S <sub>z</sub>	A's scores	B's scores
Test 1	54.10	14.28	45	60
Test 2	21.25	3.52	30	21

- a. Which student has the larger total raw score on tests 1 and 2?
- b. Calculate the z scores for both students on each test.
- c. Which student has the larger total of his two z scores?
- d. Which student has a better knowledge of French vocabulary? (This question has no easy answer; attempt to raise the issues involved.)
- 8. Prove that  $\sum_{i=1}^{n} z_i^2 = n 1$ .

# THE NORMAL DISTRIBUTION

### 6.1 INTRODUCTION

This chapter is an interruption, but a very necessary one. It must stand alone and fail to be immediately integrated into the statistics we have developed to this point because we are attempting here to gain knowledge of techniques that rest on essential concepts we shall not present in detail because of their complexity. This is the price that all of us pay for attempting to master the upper layer of statistics before establishing a foundation in bedrock. Obviously, we feel that the present venture is worth the price.

## 6.2 HISTORY OF THE NORMAL DISTRIBUTION

The brief history of the discovery and study of the normal distribution given here does not do the topic justice. Even the mathematically naïve student will find Helen Walker's account of the history of the normal distribution informative and rewarding reading (Walker, 1929, Chap. II).

THE NORMAL DISTRIBUTION

In Europe in the seventeenth century, a handful of mathematicians were pursuing small, private researches which would one day be incorporated into the theory of probability. (See Chapter 10.) These studies, by such men as Blaise Pascal (1623-1662) and Pierre de Fermat (1601-1665), were undertaken at the request of Chevalier de Mere, a gambler to whom the nature of chance must have been particularly urgent.

One of the single greatest events in the early history of probability was the publication in 1713 of Ars Conjectandi by the Swiss mathematician Jacob Bernoulli (1654-1705). A central issue in probability theory during its infancy was determining the probability that an event would occur some number of times if it were given several independent opportunities to occur. For example, if a fair coin is flipped 20 times, what is the probability that 15 "heads" will occur? Or, if a die is cast 10 times, what is the probability that the face of the die on which six dots appear will turn up exactly twice? The solutions to these problems were known at the time Ars Conjectandi was published (we will develop the solutions in Chapter 10), and the formula properties of these solutions were the main concern of this great work. However, the computations involved in obtaining the solutions for large problems were enormous. No reasonable man would attempt to calculate directly the probability that 10,000 tosses of a coin will result in 8000 or more "heads," for example. Although it would be clear to him what calculations are necessary, they would be too laborious to carry out.

Some activity in the early eighteenth century was directed toward finding convenient mathematical approximations to many of the calculations that problems in probability involved. In 1730, James Stirling published a formula that gave an approximation to the grand product of the first nintegers, i.e.,  $(1)(2)(3) \dots (n-1)(n)$ , a term that appears often in probability theory. (Whether Stirling actually derived the approximation himself is currently being disputed by mathematicians.) With Stirling's approximation, the stage was then set for the solution of the grandest problem of all. How does one approximate the probability that n independent trials of an event with probability P of producing one (a "success") of its two outcomes will yield r "successes"? The man equal to the task at the time was Abraham De Moivre (1667-1754).

Before returning to De Moivre's work, let us take a closer look at the problem he sought to solve. Suppose that a coin is to be flipped 10 times. Assume that the coin is just as likely to come up "heads" as "tails." We can ask, what is the probability that 0 "heads" will result, or what is the probability that 1 "head" will result, ..., or what is the probability that 10 "heads" will result from the 10 flips? The exact answers to all eleven of these questions can be calculated, even though for just 10 tosses the calculations are already becoming arduous. (For as many as 1000 flips of a coin the calculations border on being impossible.) The probabilities that

 $0, 1, 2, \ldots, 9$ , or 10 "heads" will result in 10 flips of a fair coin are graphed in Fig. 6.1.

The problem De Moivre sought to solve was how to find a mathematical curve that would approximate closely the curve obtained by connecting the tops of the columns in Fig. 6.1. If such a curve could be found, the nearly impossible calculations of the probability problem could be replaced by simply reading off points on mathematical curves or looking up numbers in a mathematical table.




FIG. 6.1 Graph of the distribution of probabilities of obtaining certain numbers of "heads" in 10 flips of a fair coin.

De Moivre was able to show that a mathematical curve which came very close to the curve that connects the tops of the lines in Fig. 6.1 (and the curve for almost any other such problem) has the following formula:

$$u = \frac{1}{\sqrt{2\pi} \sigma} e^{-(X-\mu)^{\frac{3}{2}/2\sigma^{\frac{3}{2}}}},$$
 (6.1)

where u is the height of the curve directly above any given value of X in the plotted frequency distribution,

 $\pi$  is the ratio of the circumference of any circle to its diameter, which is approximately 3.142,

e is the base of the system of natural logarithms, approximately 2.718, and

 $\mu$  and  $\sigma$  are numbers that locate the curve along the number line and control its spread.

Equation (6.1) is the formula for the normal distribution. It is certainly ominous looking, but do not let that frighten you. We shall have little to do with the formula as such.

The graph of Eq. (6.1) yields the familiar, symmetric, bell-shaped curve known as the normal curve. We speak of a normal curve, because Eq. (6.1) imparts a characteristic shape to the graph. However, by changing the values of u and  $\sigma$  we can move the normal curve up and down the scale and change its spread. The value  $\mu$  corresponds to the mean of a large frequency distribution that would look like the normal curve, and  $\sigma$  corresponds to the standard deviation of the distribution. In Fig. 6.2, the graph of the normal distribution for  $\mu=0$  and  $\sigma=1$  appears. We shall use the letter z as the symbol for a normally distributed variable with  $\mu = 0$  and  $\sigma = 1$ .

The curve in Fig. 6.2 does not meet the z-axis at the points 3 and -3; in fact, although it gets closer and closer to the z-axis as  $\hat{x}$  gets larger than 3, it never touches the axis. The highest point on the curve is above the z value of 0; at this point u is approximately .3989. Notice that the curve is symmetric around the vertical line drawn through  $z=\mu=0$ . The normal curve will always be symmetric around  $\mu$ . The area of the space under the curve and above the z-axis is exactly 1. The skewness of the normal curve is zero, because the curve is perfectly symmetric around the central value  $\mu$ . The kurtosis of the normal curve, i.e., the average of the fourth powers of the z-scores, is 3. As we noted earlier, distributions with a measure of kurtosis greater than 3 are more peaked than the normal distribution and are said to be leptokurtic. Distributions which are flatter than the normal have a measure of kurtosis less than 3 and are called platykurtic. The normal distribution is said to be mesokurtic (see Sec. 5.11).

Another feature of the normal curve is the special way in which it bends. On either side of  $\mu$ , which is 0 in Fig. 6.2, the curve ceases to bow away from the X-axis and begins to bow toward it, i.e., the curve has a point of inflection, at exactly a distance of one  $\sigma$  from  $\mu$ , which is one unit in Fig. 6.2.

The normal curve in Fig. 6.2 is a special one because it has been chosen as a standard. It is called the unit normal curve because the area under the curve is 1. Its mean and standard deviation are convenient ( $\mu = 0$ ,  $\sigma = 1$ ), and any other normal curve can be moved along the number scale and stretched or compressed by a simple transformation (i.e., subtract  $\mu$  and divide by  $\sigma$ ) so that it coincides with the unit normal curve.

It will often be necessary to find the ordinate u (the height of the curve above the z-axis) for any value of z on the unit normal curve or the area under the curve between any two values of z. Solving Eq. (6.1) for u when z is given is far too inconvenient; and even though we know that the area under the curve from  $z=-\infty$  to  $z=\infty$  is 1, the area between any other two values of z is difficult to find. Statistical tables are a necessity at this point. Table B in the Appendix gives the area under the unit normal curve to the




FIG. 6.2 The normal distribution for  $\mu = 0$  and  $\sigma = 1$ .

left of any point on the z-axis from -3.00 to +3.00. Also given in Table B is the ordinate u of the unit normal curve for values of z from -3.00 to **1.3 00** 

A few examples will illustrate the use of Table B. Suppose one desires to find the area under the unit normal curve to the left of z=-2.50. The value -2.50 is found in the first column of Table B. To the right of this entry in the second column, titled "Area," the number .0062 is found. Thus. only 62 ten-thousandths of the area under the unit normal curve is contained to the left of z = -2.50. The height of the unit normal curve at the point z = -2.50 is found in the "Ordinate" column to the right of the "Area" column. For z = -2.50, u = .0175.

Check and see if you can look up these areas and ordinates correctly in Table B in the Appendix.

Ordinate at 7

Value of z	Area to the left of z	Ordinate at	
0.00	.5000	.3989	
0.50	.6915 .1020	.3521 .1781	
-1.27 1.96	.9750	.0584	

Since the total area under the curve is 1, the above areas (but not the ordinates) can be read as proportions or percents of the total. 97.5% of the area under the unit normal curve lies to the left of 1.96.

Table B is also used to find the area under the unit normal curve between any two values of z. For example, the area to the left of z = -1.27 is .1020 and the area to the left of z = 0.50 is .6915. Therefore, the area between and the area between -1.27 and 0.50 is .6915 -1.1020 = .5895. In other words, about 59% of the area lies between these two points. This is illustrated in Fig. 6.3.




FIG. 6.3 Determining the area under the unit normal curve between two values of z.

6.4 THE FAMILY OF NORMAL CURVES

Actually there are infinitely many normal curves, a different one for each different pair of values for  $\mu$  and  $\sigma$ . The curve in Fig. 6.4 has a mean  $\mu$  of 20 and a standard deviation  $\sigma$  of 5, but it is a normal curve nonetheless.

What do all of these normal curves have in common? For our purposes, their most important common property is the amount of area under the curve between any two points expressed in standard deviations. For example, in any normal distribution approximately

- 1. 68% of the area under the curve lies within one  $\sigma$  of the mean either way (i.e.,  $\mu \pm 1\sigma$ ),
- 2. 95% of the area under the curve lies within two σ's of the mean, and
- 3. 99.7% of the area under the curve lies within three  $\sigma$ 's of the mean  $\mu$ .

You can check these relationships and obtain the exact areas by finding the areas under the unit normal curve between -1 and +1, -2 and +2, and -3 and +3 in Table B of the Appendix.

The normal curve and its relationship to various transformed scales which are widely used in educational and psychological measurement appear in Fig. 6.5.




FIG. 6.4 The normal curve for  $\mu = 20$  and  $\sigma = 5$ .

## 6.5 THE UNIT NORMAL DISTRIBUTION AS A STANDARD

The value of X for the unit normal distribution locates a point X units from the mean. It will be most useful in the future if all references to scores in normal distributions are in terms of deviations from the mean  $\mu$ , in standard deviation  $\sigma$  units. For almost any application of the normal curve, we shall want to know how many standard deviations a score lies above or below the mean. Knowing this, questions about the area between points on any normal curve or heights of the curve above any point can be answered by reference to the unit normal curve. The deviation of a score from its mean is  $X - \mu$ ; the number of standard deviations X lies from its mean is  $(X - \mu)/\sigma$  is called the unit normal deviate. If X has a normal distribution with mean  $\mu$  and standard deviation  $\sigma$ ,  $(X - \mu)/\sigma$  has the unit normal distribution, but not otherwise.

The shape of the normal curve does not change when we subtract  $\mu$  and divide by  $\sigma$ . If we would like to know what proportion of the area lies to the left of a score of 20 in a normal distribution with mean 25 and standard deviation 5, we can translate this question into "What proportion of the area lies to the left of (20  $\sim$  25)5 = -1 in the unit normal distribution ?"

These points are summarized in the following statement:

If X has a normal distribution with mean  $\mu$  and standard deviation  $\sigma$ , then  $z = (X - \mu) \sigma$  has a normal distribution with mean 0 and standard deviation 1, i.e.,  $z = (X - \mu) / \sigma$  has the unit normal distribution.

The area between  $X_1$  and  $X_2$  in the normal distribution with mean  $\mu$  and standard deviation  $\sigma$  is the same as the area between  $z_1 = (X_1 - \mu)_j \sigma$  and  $z_2 = (X_2 - \mu)_j \sigma$  in the unit normal distribution.

#### 6.6 USES OF THE NORMAL CURVE

De Moivre invented the normal curve for a particular use; namely, to provide an easy, approximate solution to applications of probability theory. Surely he was never aware that his discovery would find applications in practically every corner of science that now exists. Indeed, the wide application and occurrence of the normal distribution are a wonder.

The normal distribution plays an important role in both descriptive and inferential statistics. We must defer any discussion of the normal distribution in inferential statistics until the later chapters in this text.




FIG. 6.6 Frequency polygon for heights of 8585 adult men born in Great Britain during the nineteenth century.

The normal curve is an excellent approximation to the frequency distributions of large numbers of observations taken on a variety of variables. The frequency polygons of heights of adult males and adult females both look quite like a normal curve. The frequency polygon in Fig. 6.6 shows the distribution of heights of 8585 adult men born in Great Britain during the nineteenth century.


The frequency polygon in Fig. 6.6 is based on the frequency distribution shown in Table 6.1 (see Rugg, 1917):

TABLE 6.1

### FREQUENCY DISTRIBUTION FOR FIG. 66

	ENCY DISTRIBU	Height	Frequency	Height	Frequency
Height	Frequency 4	64*	669	70°	646 392
58" 59" 60"	14 41	65* 66*	990 1223	72* 73*	202 79
61° 62″	83 169	67°	1329 1230	74* 75*	32 16
63"	394	69*	1067	76*	

Psychometric tests of general and special mental abilities often yield distributions of scores that conform closely to the normal distribution. It is fairly well known that IQ's from the Stanford-Binet Intelligence Test are approximately normally distributed with a mean (a) of 100 and standard deviation (a) of 16 for people in general (see Fig. 6.7). Tests of educational



and M. A. Merrill, Measuring Intelligence (Boston: Houghton Mifflin, 1937).

attainment which are constructed in accord with the same psychometric principles upon which ability tests rest usually produce frequency polygons that resemble the normal curve. Measures of human ability in general are frequently nearly normally distributed. Whether or not some ties can be drawn between this fact and the relationship of the normal curve to the probability distribution of groups of binomial trials for chance events is a task we leave gladly to the more philosophically inclined.

Somehow the misapprehension arises in the minds of many students that there is a necessary link between the normal distribution—an idealized description of some frequency distributions—and practically any data they might collect. The normal curve is a mathematician's invention that is a reasonably good description of the frequency polygon of measurements on several different variables. A collection of scores that are exactly normally distributed has never been gathered and never will be. But much is gained if we can tolerate the slight error in the statement and claim from time to time that scores on a variable are "normally distributed." (On this point, see Borine, 1920, and Kelley, 1923.)

The mathematical statistician has greatly contributed to the eminence of the normal distribution. Although many different mathematical curves would fit empirical frequency polygons tolerably well, special mathematical advantages are gained when the normal curve can be assumed to "fit the data." Certain mathematical properties of the normal curve, Eq. (6.1), produce simple and elegant proofs to many inferential statistical problems. Without the normal curve, mathematical statisticians would have to labor under extreme complications that arise when other mathematical curves are used to represent data.

### 6.7 THE BIVARIATE NORMAL DISTRIBUTION

The theory of correlation, which will be the subject of Chapter 7, has close historical ties with the normal distribution and the bicariate normal distribution. One preoccupation of statistics since its inception as a formal discipline has been the description of the way variables are related. Do tall fathers tend to have tall sons? Will a plot of land produce a higher yield of corn if we increase the amount of nutrients in the soil? Are bright children less athletic than less intellectually able children? Each of these questions can be studied abstractly as the problem of describing the way in which scores on a variable X are paired with scores on a second variable Y for the same persons. Thus, these questions concern bicariate relationships, i.e., relationships between two variables.

If we measure a large group of persons in two ways, e.g., we might measure each person's IQ(X) and physical strength (Y), the data can be represented in a bivariate frequency distribution. For each person there is a pair of scores, his score on X and his score on Y. A bivariate frequency distribution is a picture of the frequency with which different pairs of X and Y scores occur in a group of persons. Figure 6.8 is a bivariate frequency distribution for a group of persons measured on IQ(X) and physical strength

(Y).

From Fig. 6.8, we can see that approximately 20 persons scored 125 on the IQ variable X and 30 on the physical strength variable Y. The height of the line at the intersection of 125 and 30 must be measured against the vertical scale of frequencies.




FIG. 6.8 Bivariate frequency distribution for a large group of persons measured on IQ (X) and physical strength (Y).

A large number of bivariate frequency distributions built from data gathered in educational and psychological settings show a characteristic shape. A surface drawn through the end points of the columns which represent the frequencies in a bivariate frequency distribution often looks like a bell—in three dimensions—that has been stretched in the X and Y directions and rotated around its center in the X-Y plane. Much was to be gained if the mathematical statistican could find a set of mathematical curves that gave a good description of many bivariate frequency distributions. The mathematical surface which was found to accomplish this objective is called the bicariate normal distribution. This smooth, continuous, bell-shaped surface provides a mathematically convenient and satisfactory representation of numerous bivariate frequency distributions.




FIG. 69 One of the family of bivariate normal distributions (a cross section).

As was true of the normal distribution, the bivariate normal distribution is a family of three-dimensional surfaces. One member of this family appears in Fig. 6.9. All bivariate normal distributions have the following characteristics:

- The distribution of the X scores, completely disregarding the Y scores with which they are paired, is a normal distribution.
- The distribution of the Y scores, completely disregarding the X scores with which they are paired, is a normal distribution.
- 3. For each single score on X, say  $X_1$ , the Y scores of persons with a score on X of  $X_1$  are normally distributed with variance  $\sigma_{x,x}^2$ .
- 4. For each single score on Y of  $Y_1$ , the X scores of persons with a score of  $Y_1$  are normally distributed with variance  $\sigma_1^2$ .
- The means of the Y scores for each separate score on X fall on a straight line.

We shall refer to the bivariate normal distribution on several occasions, but we shall not explore further properties of it. For a more advanced discussion see Walker and Lev (1953, pp. 248-49).

### PROBLEMS AND EXERCISES

- Let z stand for the unit normal variable, i.e., the normally distributed variables with mean 0 and standard deviation 1. Find the area under the unit normal curve which lies:
  - a. above z = 1.00;
  - b. below z = 2.00; c. above z = 1.64;
  - c. above z = 1.04
  - d. below z = -1.96;
  - e. between z = 0 and z = 3.00;
  - f. above z = -0.50;
  - g. between z = -1.50 and z = 1.50.
- Find the ordinates of the unit normal distribution above each of the following z scores:
  - a. z = 1.00;
  - b. z = -1.00;
  - c. z = 2.25:
  - d. z = -0.15.
- Find the z scores which are exceeded by the following proportions of the area under the unit normal distribution;

	Proportion of area above z score	z scor
a.	0.50	0
6.	0.16	+1.00
c.	0,84	
đ.	0.05	
e.	0.005	
f.	0.995	
g.	0.10	

4. If in the general population of children Stanford-Binet IQ's have a nearly normal distribution with mean 100 and standard deviation 16, find the percentile equivalent of each of the following IQ's:

	I.Q.	Percentile equivalen
a,	100	
ъ.	120	
c.	75	
đ.	95	
e.	140	

108 THE NORMAL DISTRIBUTION

> 5. X and Y have a bivariate normal distribution. Both X and Y have means of zero and variances of 1. Suppose that for X = 1.25 the variance of the associated Y variable is 0.50. What is the variance of the Y variable associated with a value of 1.50 for the X variable?

#### MEASURES OF RELATIONSHIP

#### 7.1

In this chapter we shall begin the study of the description of relationships between variables. Before we leave this general topic at the end of Chapter 9, we hope that the major portion of the description of relationships between variables which you will find useful will have been covered. A complete discussion of the measurement of relationship or correlation, a subject that has undergone almost a century of research, would easily require a book ten times as long as this chapter. The topics we shall purposely overlook in this text can be found in references cited in Sec. 7.9 at the end of this chapter.

# 7.2 THE PEARSON PRODUCT-MOMENT CORRELATION COEFFICIENT

Researchers are often concerned with the way in which two variables relate to each other for a given group of persons (classrooms, schools, nations,

etc.). For example, do students who tend to read earlier than others also tend to have higher achievement in science in the sixth grade? Do large classrooms show lesser gains in knowledge over a semester than smaller classrooms? Is the average length of employment of teachers in a school directly related to the average salary for teachers? Obviously, to answer such questions we must make observations on each variable for a group of units (typically persons, but they might be classrooms, schools, counties, etc.). The data gathered to answer one such question might take the following form:

Student no.	Stanford-Binet IQ score (X)	Raw score on chemistry achievement test (Y		
1	120	31		
2	112	25		
3	110	19		
4 5	120	24		
	103	17		
6	126	28		
7	113	18		
8	114	20		
9	106	16		
10	108	15		
11	128	27		
12	109	19		

In this example, the variables observed on 12 students were IQ as determined by the Stanford-Binet Intelligence Scale in grade 6 and achievement in high-school chemistry as measured by a 35-item teacher-made test.




FIG. 7.1 Scatter diagram showing relationship of IQ (X) to achievement in chemistry (Y) for 12 students.

The relationship between the two variables can be depicted graphically in a presentation of the data called a scatter diagram. The scatter diagram for the above data appears as Fig. 7.1.

Each unit (person in this example) is represented by a point on the scatter diagram. A dot or mark is placed for each person at the point of intersection of straight lines drawn through his 10, score perpendicular to the X axis and through his chemistry-test score perpendicular to the Y axis. The scatter diagram in Fig. 7.1 shows a moderate positive relationship between X and Y. As yet, however, we have no summary measure of this relationship.

The general question of "relationship" must be given a somewhat more precise meaning. Is high relative standing on X of a person with respect to a group paired with high or low relative standing of that person on Y, or is there no systematic pairing off of high and low relative standings?

The standing of a person relative to others in a group on X and Y relative to the means of two distributions is reflected in the size and algebraic sign of the deviation scores  $(X_i - \bar{X})$  and  $(Y_i - \bar{Y})$ , respectively. If a person is high on both variables, as is student II above, the product of  $(X_i - \bar{X})$  and  $(Y_i - \bar{Y})$  will be large and positive for him. Similarly, if a person is relatively low on both X and Y,  $(X_i - \bar{X})(Y_i - \bar{Y})$  will be large and positive for him also (because the product of two negative numbers is a positive number). Now if X and Y are directly related (high paired with high and low with low) substantially, most of the products  $(X_i - \bar{X})(Y_i - \bar{Y})$  will be positive; consequently, the sum of these products for all persons

 $\left[\text{i.e., }\sum_{i=1}^{n}(X_{i}-\vec{X}_{i})(Y_{i}-\vec{Y}_{i})\right]$  should be large and positive.

If X and Y bear an *inverse* relationship to each other (high X paired with low Y and vice versa), many persons with positive  $(X_t - X_t)$  scores will tend to have negative  $(Y_t - \overline{Y})$  scores, and negative  $(X_t - \overline{X})$  scores will tend to be paired with positive  $(Y_t - \overline{Y})$  scores. In this case, the products  $(X_t - \overline{X})(Y_t - \overline{Y})$  will generally be negative. Hence,

$$\sum_{i=1}^{n} (X_i - \vec{X})(Y_i - \vec{Y})$$

will be negative when X and Y are inversely related.

If X and Y bear no systematic relationship to each other (high X's are as likely to be paired with low Y's as with high Y's, and the same is true of low X's), then of the people with large positive  $(X_t - \bar{X})$  scores some will have positive  $(Y_t - \bar{Y})$  scores and others will have negative  $(Y_t - \bar{Y})$  scores. When the products  $(X_t - \bar{X})(Y_t - \bar{Y})$  are formed, some will be positive and others will be negative. The sum of the products,

$$\sum_{i=1}^{n} (X_i - \bar{X}_i)(Y_i - \bar{Y}_i),$$

should contain about an even balance of positive and negative terms of the same size and should therefore be relatively close to zero.

Thus we have the quantity  $\sum_{i=1}^{n} (X_i - \bar{X}_i)(Y_i - \bar{Y}_i)$  being large and positive

when X and Y are strongly related directly, being near zero when X and Y are unrelated, and being large and negative when X and Y are strongly related inversely. However, this sum of the products of deviation scores is still not an adequate summary measure of relationship. For one thing, its size depends on how many pairs of scores are included in its calculation. Since we may wish to compare the degree of relationship between X and Y in two groups of different size, we shall want to make the measure of relationship independent of the size of the group on which it is calculated. A simple averaging procedure will accomplish this. Two means calculated on different size groups are comparable in terms of locating central scores, but the simple sums of the two groups of scores are not. This is why we take an average if we want a statistic to be independent of group size. However, for the same reason that \$\frac{1}{2}\$ was defined by dividing the sum of squared deviations by \$n - 1\$ instead of \$n\$, we should divide

$$\sum_{i=1}^{n} (X_i - \overline{X}_i)(Y_i - \overline{Y}_i) \text{ by } n-1.$$

The quantity  $\sum_{i=1}^{n} (X_i - \bar{X}_i)(Y_i - \bar{Y}_i)/(n-1)$  is a measure of the relationship between X and Y and is called the *covariance* of X and Y. The covariance of X and Y is denoted by  $s_{n-1}$ :

$$s_{xy} = \frac{\sum_{i=1}^{n} (X_i - \bar{X})(Y_i - \bar{Y})}{n-1}.$$
 (7.1)

Notice that the covariance of X with itself is simply the variance of X:

$$s_{xx} = \frac{\sum_{i=1}^{n} (X_i - \bar{X})(X_i - \bar{X})}{n-1} = \frac{\sum_{i=1}^{n} (X_i - \bar{X})^2}{n-1} = s_x^2$$

The covariance is a perfectly good measure of association in many problems in the physical sciences and engineering. (In fact, physicists call the hallowed "correlation coefficient" of the behavioral sciences—which we shall encounter presently—the "dimensionless covariance.") And it is an adequate measure as long as the scale (mean and variance) of the variables are not arbitrary and contain some meaning. Many variables with which whatever anyone wishes since we are usually interested only in relative positions in a group. This is particularly true of psychological and educational test data.

The process of deviating both the X and Y scores around their respective means has made  $s_{x_i}$  independent of the means of the scores. To make the desired measure of relationship independent of the standard deviations of

the two groups of scores, one need only divide  $s_{xy}$  by  $s_x$  and  $s_y$ . The result is the desired measure of relationship between X and Y. It is called the Pearson product-moment correlation coefficient and is denoted by  $r_{xy}$ :

$$r_{xy} = \frac{s_{xy}}{s_x s_y} \tag{7.2}$$

The designation r comes from the word regression. In the early applications of the coefficient by Francis Galton and Karl Pearson (1857–1936), it played an important role in the study of association of physical characteristics in humans, a study that first pointed up the regressive nature of physical measurements from one generation to the next. Although Pearson played the most important role in establishing the mathematical properties of  $r_{xy}$ , the notion of a correlation coefficient equal to  $s_{xy}/(s_x s_y)$  can be traced through the writings of Galton back to an article published in 1846 by the Frenchman Bravais.

#### 7.3 A COMPUTATIONAL FORMULA FOR r.,

Equation (7.2) is definitional and not convenient for computing r<sub>av</sub>. We shall now derive a form more convenient for calculating r<sub>av</sub> on a desk calculator, given the raw scores X and Y. Begin with

$$r_{xy} = \frac{s_{xy}}{s_x s_y} = \frac{\sum_{i=1}^{n} (X_i - \bar{X})(Y_i - \bar{Y})/(n-1)}{\sqrt{\sum_{i=1}^{n} (X_i - \bar{X})^2/(n-1)\sqrt{\sum_{i=1}^{n} (Y_i - \bar{Y})^2/(n-1)}}.$$
 (7.3)

Note that 1/(n-1) can be factored out of the two terms  $(1/\sqrt{n-1})$  from each term) in the denominator of Eq. (7.3), and it cancels the 1/(n-1) in the numerator of Eq. (7.3). Also recall that since  $\sqrt{a}\sqrt{b} = \sqrt{ab}$ , the terms in the denominator of Eq. (7.3) can be combined under the radical.

$$r_{av} = \frac{\sum_{i=1}^{n} (X_i - \bar{X})(Y_i - \bar{Y})}{\sqrt{\left[\sum_{j=1}^{n} (X_i - \bar{X})^2\right] \left[\sum_{j=1}^{n} (Y_i - \bar{Y})^2\right]}}.$$
 (7.4)

Consider just the numerator of (7.4):

$$\sum_{i=1}^{n} (X_i - \bar{X})(Y_i - \bar{Y}) = \sum_{i=1}^{n} X_i Y_i - \bar{X} \sum_{i=1}^{n} Y_i - \bar{Y} \sum_{i=1}^{n} X_i + n \bar{X} \bar{Y}. \quad (7.5)$$

Several steps were involved in moving from Eq. (7.4) to Eq. (7.5): expanding a binomial, moving constants (e.g., X) outside of summation signs, and summing constants. Try to fill in the details.

Recalling that  $\sum Y = n\overline{Y}$ , and  $\sum X = n\overline{X}$ , we can write the right side of Eq. (7.5) as follows:

$$\sum_{i=1}^{n} X_{i} Y_{i} - n \overline{X}_{i} \overline{Y}_{i} - n \overline{Y}_{i} \overline{X}_{i} + n \overline{X}_{i} \overline{Y}_{i} = \sum_{i=1}^{n} X_{i} Y_{i} - n \overline{X}_{i} \overline{Y}_{i}.$$
 (7.6)

If we had chosen to replace  $\bar{X}_{\cdot}$  by  $\sum X/n$  and similarly for Y, we would have obtained

$$\sum_{i=1}^{n} (X_{i} - \bar{X})(Y_{i} - \bar{Y}) = \sum_{i=1}^{n} X_{i} Y_{i} - \frac{(\sum X_{i})(\sum Y_{i})}{n}.$$
 (7.7)

Either Eq. (7.6) or Eq. (7.7) provides a simple formula for the numerator of  $r_{xy}$ . We already know a simple way to compute the denominator of  $r_{xy}$ .

$$\sum (X_i - \bar{X})^2 = \sum X_i^2 - \frac{(\sum X_i)^2}{n}$$
 (7.8)

$$\sum (Y_i - \bar{Y})^2 = \sum Y_i^2 - \frac{(\sum Y_i)^2}{2}. \tag{7.9}$$

Combining Eqs. (7.7), (7.8), and (7.9) produces the following formula for  $r_{xy}$ ,

$$r_{xx} = \frac{\sum X_i Y_i - (\sum X_i)(\sum Y_i)/n}{\sqrt{[\sum X_i^2 - (\sum X_i)^2/n][\sum Y_i^2 - (\sum Y_i)^2/n]}},$$
 (7.10)

which can be simplified further to become the computational formula

$$r_{xy} = \frac{n \sum X_i Y_i - (\sum X_i)(\sum Y_i)}{\sqrt{[n \sum X_i^2 - (\sum X_i)^2][n \sum Y_i^2 - (\sum Y_i)^2]}}.$$
 (7.11)

Equation (7.11) is more convenient than Eq. (7.10) for finding  $r_{\rm ev}$  on a deck calculator on which "negative multiplication" is possible. It will then be possible to calculate the numerator of Eq. (7.11) without divisions and without writing any numbers down on paper. This is also true for both terms in brackets in the denominator of Eq. (7.11).

7.4

### ILLUSTRATION OF THE

Some ability-test data will be used here to illustrate the calculation of  $r_{\rm ex}$  (7.10) and (7.11). A researcher is studying the relationship between two types of reasoning ability in high-school juniors: abstract reasoning and verbal reasoning. Two tests are constructed, one measuring abstract reasoning (X) and the other measuring verbal reasoning (Y). The two tests were administered to 40 high-school juniors from the junior class

of the only high school in an Illinois town of about 30,000 residents. The test scores for the 40 students appear in Table 7.1. Each test was 50 items

TABLE 7.1 RAW SCORES ON SO-ITEM TESTS OF ABSTRACT AND VERBAL REASONING ABILITY FOR 40 ILLINOIS HIGH-SCHOOL HUNDERS

Student	X Abstract reasoning	Y Verbal reasoning	Student	X Abstract reasoning	Y Verbal reasoning
Linda J.	19	17	Martin T.	38	30
Peggy Y.	32	7	Sharon L.	25	18
Deane L.	33	17	Julie E.	35	26
Constance L.	44	28	Natalie J.	22	17
William P.	28	27	Maryjean K.	40	17
Roger D.	35	31	Larry N.	42	26
Caroline E.	39	20	Michael B.	41	16
Trudy R.	39	17	Carleen M.	41	37
Peter A.	44	35	Scott C.	37	26
David E.	44	43	Sigrid K.	30	21
Cheryl G.	24	10	Jan W.	31	16
Georgia S.	37	28	Roger B.	41	37
Erma J.	29	13	Richard H.	42	37
Ronald L.	40	43	Bonita G.	24	14
Pamela J.	42	45	Rex N.	43	41
Edward B.	32	24	Richard S.	36	19
Rosa L.	48	45	Maurice D.	39	18
Karen M.	43	26	Warren W.	39	39
Roger W.	33	16	Jack G.	39	37
Richard T.	47	26	Stanley L.	48	47

<sup>\*</sup> We wish to express our appreciation to Dr. J. Thomas Hastings, Director of the Illinois Statewide Testing Program, who made these data available.

in length, and the test score was the number of correct answers given. A scatter diagram of the bivariate data in Table 7.1 appears in Fig. 7.2.

scatter diagram of the bivariate data in Table 1/1 appears in Fig. 1.2.

The intermediate and final calculations of Fig. for both formulas (7.10) and (7.10) appear in Table 7.2. All calculations were partformed as a desk calculator. (Without mechanical computation, the calculation of productmoment correlation coefficients is usually tedious.) Perhaps the only quantity

in Table 7.2 whose origin is doubtful in your mind is  $\sum_{i=1}^{6} X_i Y_i$ . This quantity is the sum over all persons of the *product* of each person's X and Y scores. For the first student in Table 7.1, Linda J.,  $X_1 = 19$  and  $Y_1 = 17$ ; for the second student, Pegy  $Y_1$ ,  $X_2 = 22$  and  $Y_2 = 7$ . The quantity

$$\sum_{i=1}^{40} X_i Y_i = (19 \cdot 17) + (32 \cdot 7) + \ldots + (48 \cdot 47) = 40,798.$$

The final calculations on the right-hand side of Table 7.2 show  $r_{ev}$  to

116

ILLUSTRATION OF THE CALCULATION OF I, FOR THE TABLE 7.2

DATA IN TABLE 7.1 Intermediate calculations

Final calculations

Equation (7.10):
$$n = 40$$

$$r_{er} = \frac{40,798 - (1465)(1057)/40}{\sqrt{(55,725 - (1465)/40)}}$$

$$\sum_{i=1}^{50} X_i = 1465$$

$$\sum_{i=1}^{40} Y_i^2 = 1057$$

$$\sum_{i=1}^{50} X_i^2 = 55,725$$

$$\sum_{i=1}^{50} Y_i^2 = 32,551$$

$$\sum_{i=1}^{51} X_i Y_i = 40,798$$

$$r_{rr} = \frac{40(40,798) - (1465)(1057)}{\sqrt{(40(55,725) - (1465))^2}}$$

$$r_{rr} = \frac{40(40,798) - (1465)(1057)}{\sqrt{(40(55,725) - (1465))^2}}$$

$$= \frac{83,415}{123,766,128} = 0.67.$$

be 0.67, by either Eq. (7.10) or Eq. (7.11). It will always be true that the two formulas will yield the same value, within rounding error. Thus, there appears to be a strong, direct relationship between abstract and verbal reasoning ability as measured by the two tests.




FIG. 7.2 Scatter diagram of the 40 pairs of test scores in Table 7.1.

### 7.5 RANGE OF VALUES OF rev

It is somewhat difficult to prove, but  $r_{ev}$  can never take on a value less than -1 nor a value greater than +1.\* (If you are dismayed by the apparent

TABLE 7.3 INTERPRETATION OF VALUES OF 120 Value of Description of linear T... relationship Scatter diagram +1.00Perfect, direct relationship About +.50 Moderate, direct relationship .00 No relationship (i.e., 0 covariation of X with Y) About --.50 Moderate, inverse relationship -1.00 Perfect, inverse relationship

$$\sum z_n^2 = \sum z_n^2 \approx n-1 \quad \text{and} \quad r_{nn} \approx \frac{\sum z_n z_n}{(n-1)}.$$

To show that  $r_{xy}$  cannot be less than -1, work with  $\sum (z_x + z_y)^2$ .

<sup>\*</sup> To prove that the value of  $r_{sv}$  cannot exceed +1, expand  $\sum_{1}^{n} (z_{s}-z_{s})^{s}$ , which is always greater than or equal to zero, and use the fact that

difficulty of the proof suggested in the footnote, you can at least take consolation in the fact that the simpler proof often given in elementary textbooks is fallacious.) Table 7.3 is a listing of various values of  $r_{av}$  with illustrations of the type of linear relationship that exists between X and Y for the given values of r...\*

In Table 7.4 some representative correlation coefficients are presented.

TABLE 74

TYPICAL VALUES O	CAL VALUES OF res			
Description:	of variables	Nature of	Typical value	
x	Y	subjects	of ras	
Iowa Test of Educational Development, Grade 9	Freshman grade-point average in college	Over 600 college students	.58	
The Stanford-Binet Intelligence Test IQ	The same test given one week later	Elementary-school pupils	.90	
Verbal reasoning ability (as measured by the Differential Aptitude Test)	Nonverbal reasoning ability	High-school juniors	,65	
Height	Achievement in college physics	Male college seniors	.00	

After a little experience with actual data, you will begin to develop a "feel" for the degree of relationship that is indicated by any particular value of r. You will probably learn to associate in your mind a scatter diagram of points and the approximate value of r that corresponds to it. We hesitate to apply descriptive adjectives to values of r, such as calling an r of .80 "high" or an r of .20 "low." Whether or not a particular r is "high," "low," or "moderate" depends upon how the two variables being correlated have been related in the past, what use one intends to make of the relationship between the variables, etc. Moreover, why use a vague and ambiguous adjectival description for a value of r when it is so simple merely to report the value?

Erlenmeyer-Kimling and Jarvik (1963) presented data that are enlightening illustrations of the use of r. They found typical values of r for large classes of studies published in the literature in which the intelligence-test scores of children were correlated with those of siblings, related children, or unrelated children. For example, the typical value of the correlation

<sup>•</sup> The sort of relationship  $r_{xy}$  is designed to measure is a special type, to be discussed in Chapter 8. For more on this point, see Sec. 7.7 on curvilinear relationships between X

coefficient between a child's IQ (X) and the IQ of his identical twin (Y) for a large set of identical-twin pairs was .88 when the twins were reared together. The typical correlation between IQ's of identical twins reared apart was .75. These and other data are reported in Table 7.5.

TABLE 7.5 CORRELATIONS BETWEEN IQ'S OF RELATED OR UNRELATED CHILDREN AS A FUNCTION OF GENETIC SIMILARITY AND SIMILARITY OF ENVIRONMENT

Nature of pairing	Typical value of rzs
Identical twins, reared together	.88
Identical twins, reared apart	.75
Fraternal twins of same sex	.53
Fraternal twins of opposite sex	,53
Siblings, reared together	.49
Siblings, reared apart	.46
Parent with own child	.52
Foster parent with child	.19
Unrelated, reared together	.16

### 7.6 THE EFFECT ON r<sub>zv</sub> OF TRANSFORMING SCORES

Often the mean and variance of the scores on X and Y are arbitrary. We can change them at will without consequence, it seems. But does the value of  $r_{xy}$  depend on the means and variances of X and Y? The answer is No. This answer was implicit in our development of the formula for  $r_{xy}$ ; now we wish to make it more explicitly.

The mean and variance of X (or Y) can be changed to any value we desire by multiplying X by a constant  $b \neq 0$ , and adding a constant a to the product, i.e., by forming bX + a. This process is called "taking a linear transformation of X." Suppose we take another (or perhaps the same) linear transformation of Y, dY + c, where  $d \neq 0$ . Is the correlation coefficient between X and Y the same as that between bX + a and dY + c?

The correlation of bX + a and dY + c is the covariance of the two divided by the product of the standard deviations. We know that adding a constant to a variable does not change its standard deviation and that multiplying a variable by a constant multiplies the standard deviation of that variable by the absolute value of the constant. Thus, the standard deviation of bX + a is  $|b| s_s$ , and dY + c has a standard deviation of  $|b| s_s$ .

$$s_{a,Y+a} = |b| s_x$$
, and  $s_{d,Y+s} = |d| s_y$ . (7.12)

The covariance of bX + a and dY + c is

$$s_{(bX+a)(dY+c)} = \frac{\sum\limits_{i=1}^n [bX_i + a - (b\tilde{X}_i + a)][dY_i + c - (d\tilde{Y}_i + c)]}{n-1} \ .$$

This expression reduces to the following:

$$\frac{\sum_{i=1}^{n} (bX_{i} - b\overline{X})(dY_{i} - d\overline{Y})}{n-1} = \frac{bd\sum_{i=1}^{n} (X_{i} - \overline{X})(Y_{i} - \overline{Y})}{n-1} = bds_{xy}.$$
 (7.13)

In words, the covariance of bX + a and dY + c equals bd times the covariance of X and Y. We can combine the results in Eqs. (7.12) and (7.13) into an expression for the correlation of bX + a and dY + c.

$$r_{bX+e,dY+e} = \frac{bds_{zy}}{|b||d|s_zs_y} = \frac{bd}{|b||d|}r_{zy}.$$
 (7.14)

In words, the correlation between bX + a and dY + c equals  $r_{xy}$  times the product of b and d over the product of the absolute values of b and d. As an example, suppose that X is transformed into 3X + 5 and Y into 2Y + 8.

$$r_{2X+5,2Y+5} = \frac{3 \cdot 2}{|3| \cdot |2|} r_{xy} = r_{xy}.$$

This particular transformation of X and Y had no effect on the correlation  $r_{xy}$ . In fact the ratio of bd to |b| |d| in Eq. (7.14) can never be anything but +1 or -1. Hence, no linear transformation of either X or Y (provided b or d is not zero) can change the size of the correlation between X and Y, though it may change the size of the correlation. If either b or d, but not both, is negative, the correlation of bX + a and dY + c will equal  $-r_{xy}$ . These results are summarized in Table 7.6.

TABLE 7.6 THE EFFECT OF LINEAR TRANSFORMATIONS OF X AND Y ON THE VALUE OF r<sub>s</sub> [SPECIAL CASES OF EQ. [7.14]]

	bX + a	dY + c	Value of rexestre	
<del></del>	b is positive b is negative b is positive b is negative	d is positive d is positive d is negative d is negative	f <sub>77</sub> —f <sub>87</sub> —f <sub>84</sub> f <sub>87</sub>	

### 7.7 INTERPRETING CORRELATION

#### A. Causation and Correlation

The presence of a correlation between two variables does not necessarily mean there exists a causal link between them. Even though concomitance (correlation) between events can be useful in identifying causal relationships when coupled with other methodological approaches, it is a dangerous and potentially misleading test for causation when used alone. First, even when one can presume that a causal relationship does exist between the two variables being correlated,  $r_{\rm ex}$  can tell nothing by itself about whether X causes Y or Y causes X. Second, often variables other than the two under consideration are responsible for the observed association. Third, the relationships that exist among variables in education and the social sciences are almost always too complex to be explained in terms of a single cause. Achievement in school is the resultant of numerous influences, in addition to being a complex concept itself which cannot be described adequately by any single measurement.

We shall examine some examples of the problems that arise in attempts to unearth causal relationships with correlational techniques. It is probably true that in the United States there is a positive correlation between the average salary of teachers in high schools and the percent of the school's graduates who enter college. Does this imply that a well-paid teaching staff cause better trained high-school graduates? Would the percent of high-school graduates entering college rise if we increased the pay of teachers? Certainly affirmative answers to these questions are not justified by the associational relationship alone. The relationship between the two factors is not simple, but one prominent variable not yet mentioned is the financial and economic condition of the community that largely determines its ability to pay both teachers' salaries and college tuitions. Moreover, the economic and financial condition of the community is in part dependent upon the intellectual powers of its citizens, another variable that contributes to both higher teachers' salaries and greater college attendance among the young people.

It has been found that the percent of "dropouts" in each of a number of high schools is negatively correlated with the number of books per pupil in the libraries of those schools. But common sense tells us that piling more books into the library will no more affect the dropout rate than hiring a better truant officer will bring about a magical increase in the holdings of the school library. If only common sense always served us so well!

Many researchers do not stop with one fallacious conclusion, i.e., that correlation is prima facie evidence for causation, but draw a second one as

...

relationships, not techniques to demonstrate noncausal ones. There are only relatively few of the former and they are valuable, but noncausal relationships exist in superabundance and the discovery of one is little cause for celebration. For further discussion, see Blalock (1961) and Campbell and Stanley (1963).


#### B. Presence of Identifiable Groups of Subjects with Different Means

A substantial correlation between two variables is a fact that can be explained differently in different situations. Some correlations result from measuring a cause and its effect, e.g., when X is food intake in a period of one month and Y is weight gain over the same period. Other correlations result from measuring two variables with a common cause or influence, e.g., when X is achievement in English and Y is achievement in social studies. Still other correlations result when two distinct groups of persons within which Y and Y are unrelated are pooled together.

Suppose that girls report greater anxiety than boys on an inventory such as Taylor's Manifest Anxiety Scale. It is well known that girls tend to score higher than boys on achievement tests in English, especially in the intermediate grades. The scatter diagram of the anxiety and English achievement scores for 15 boys and 15 girls might look like the one in Fig. 7.3.

The scatter diagram in Fig. 7.3 shows a moderately strong positive relationship between anxiety and English achievement when boys' and girls' scores are pooled. Does this mean that anxiety (tension) makes a student work harder and thus achieve more? Not at all. For if it did, why would one obtain no relationship between the two variables for boys and girls separately?

Figure 7.3 shows that nonzero correlations can result when distinct groups, e.g., boys and girls, with different average values on the two variables are pooled together. Either positive or negative relationships can result from this pooling. Sketch the scatter diagram that would result if one pools two groups for which X and Y are uncorrelated and for which group A





has a high mean on X and a low mean on Y and group B has a low mean on X and a high mean on Y. Does the resulting scatter diagram correspond to a zero, a positive, or a negative correlation between X and Y?

The identification of subgroups with differing means on X and Y does not negate the fact that X and Y correlate. However, it may provide a more rational explanation of why r is substantially different from zero.

#### C. Curvilinearity and the Shapes of Marginal Distributions

Of all the possible ways in which measurements on two variables can be related,  $r_{xy}$  measures only one type. The value of  $r_{xy}$  is a measure of the degree of linear relationship between X and Y. If X and Y are perfectly linearly related, the points in the scatter diagram will fall on a single straight line, as was illustrated in Table 7.3. If we scatter the points in such a scatter diagram above and below the line in a haphazard manner and about the same distance in each direction, we obtain various degrees of basically linear relationships between X and Y. If the points in a scatter diagram look as though they depart in a haphazard manner from a curved line, the relationship between X and Y may be basically curvilinear. To say that rev measures only the linear relationship between X and Y means that different sorts of curvilinear relationships between X and Y may produce values of rev that are misleadingly close to 0 if interpreted without reference to a scatter diagram of the data. X and Y can be closely related curvilinearly and yet  $r_{xy}$  can be zero. If we know that X and Y are generally linearly related, the meaning of rzy is fairly unequivocal. However, if X and Y have some sort of curvilinear relationship, values of re, near zero can be obtained even though X and Y are highly related. Figure 7.4 shows two different scatter diagrams, each of which has correlation coefficients of approximately zero.

However, even though the scatter diagrams A and B in Fig. 7.4




FIG. 7.4 Two instances of approximately zero product-moment correlation.




FIG. 7.5 The scatter diagram of scores for test A (which is too easy) and test B (which is too difficult for the group tested).

both have correlation coefficients of zero, there is considerable relationship present in B, while any systematic relationship between X and Y is lacking in A. The single illustration in Fig. A should be sufficient warning never to draw a rash conclusion that two variables are unrelated merely because  $r_{xy}$  is zero. Various degrees of curvilinearity of relationship between measures of variables are not uncommon. Educational and psychological test scores frequently show "ceiling" or "cellar" effects with atypical groups of persons, i.e., the tests may be too easy or too difficult with the result that many persons obtain the highest or the lowest test score. The scatter diagram of test scores for test A, which shows a "ceiling effect," and test B, which shows a "ceiling effect," might look like the one in Fig. 7.3.

The value of  $r_{AB}$  for the data in Fig. 7.5 is not large; it is probably only about 30. In the range for which both tests are of appropriate difficulty, the two tests appear to be more highly related. One suspects that if test A was made more difficult and Test B easier without radically affecting the content of either test, the value of  $r_{AB}$  for these persons would increase. The scatter diagram of the test scores for such altered tests would probably show less curvilinearity than now. (This example illustrates another important point: the degree of relationship obtained between any two variables—regardless of how the relationship is expressed—depends on the nature of the measurement of the variables. For example, we generally think of the characteristics weight and height as being rather closely related in adult human beings; but it is not difficult to conceive ways of measuring each variable that are so poor—e.g., measurement by the sober subjective judgments of four-year olds that weight and height scores would show almost no correlation.)

### 7.8 FURTHER REMARKS ON THE INTERPRETATION OF $r_{xy}$

Carroll (1961) presented a readable account of how the interpretation of reg is dependent upon the shapes of the distributions of X and Y and of their

joint distribution. Carroll's article is an excellent statement of many points touched only briefly here and parts of it will be clear to the student whose acquaintance with correlation does not extend beyond this and the next two chapters. He made the following observation on both the problem of interpreting  $r_{ax}$  and the statistical training of students:

"Students are not adequately informed that these limits [-1 to -1] and meanings "highly related," "moderately related," "not related" strictly have reference to certain statistical models. Two of the most frequently used models are the normal bivariate surface [see See, 6.6] and the linear regression model [see Chapter 8]... No assumptions are necessary for the computation of a Pearsonian coefficient, but the interpretation of its meaning certainly depends upon the extent to which the data conform to an appropriate statistical model for making this interpretation. As actual data depart from a fit to such a model [e.g., the bivariate normal surface], the limits of the correlation coefficient may contract, and the adjectival interpretations are less meaningful."

As an example of how the maximum value of  $r_{\pi \tau}$  might bound away former 1 when the frequency distributions of X and Y are skewed in different distributions: 0 on X and on Y with the following frequency distributions:

Score on X: 0 1 2 3 4 5 6 7 8 9 1
Frequency: 21 12 14 14 13 10 7 4 2 1
Score on Y: 0 1 2 3 4 5 6 7 0 0 1

Frequency: 0 1 2 3 4 5 6 7 8 9 10

Frequency: 0 0 1 1 2 2 4 5 6 7 71

X is highly positively skewed, and Y is highly negatively skewed. The maximum possible value of  $r_{r_0}$  is about .60. In other words, even if the maximum possible linear relationship between X and Y exists,  $r_{r_0}$  will be only about .60. This reflects no weakness in  $r_{r_0}$  as a descriptive measure; it cannot be blamed for not doing what it was not designed to do. In fact, be rather comforting. When X has so many values below its mean and Y has so many values above its mean, it is impossible for all positive deviations of the  $Y_i$  around  $\tilde{Y}$  to be associated with positive deviations of the  $X_i$  around  $X_i$ . In fact,  $r_{r_0}$  cannot attain the extreme values of +1 or -1 unless  $X_i$  and Y distributions have identical shapes.

This dependence of the maximum value of r<sub>s</sub>, on the distributions of X and Y places one in a difficult position. Suppose that an r<sub>s</sub> of .60 is obtained. What does it reflect? Moderate relationship between two variables whose joint frequency distribution looks like a bivariate normal surface, or the maximum possible relationship between a positively skewed X and a negatively

skewed Y? Earlier we saw how similar doubts surround a zero value for  $r_{ex}$ . Are X and Y actually unrelated or is the relationship between them non-linear? The most satisfactory resolution of all of these doubts can be made by looking at a scatter diagram of the X and Y scores. From such a diagram it can be seen immediately whether or not X and Y have a pronounced curvilinear relationship and whether X and Y are markedly skewed. Regrettably, researchers are too hesitant to construct scatter diagrams. One would think that one beneficial effect of electronic data processing would be increased plotting of scatter diagrams in correlational problems, either by hand or by computer. This has not yet happened. In the authors judgment, building and inspecting the scatter diagram so that  $r_{ex}$  can be interpreted more intelligently are well worth the little effort they take.

### 7.9 THE VARIANCE OF SUMS AND DIFFERENCES OF VARIABLES

Quite often in education and psychology one wishes to find the variance of a group of summed X and Y scores. Moreover, simply inspecting the formula that related  $s_{k+\nu}^2$ , the variance of the summed X and Y scores, to  $s_{k}^2$ ,  $s_{k}^2$ , and  $r_{\mu\nu}$  can illuminate the way in which influences combine to produce joint effects. In the history of mental test theory, a general expression for the variance of a sum of variables has played an important role (a total test score is the sum of scores on the individual items of the test).

The variance of X + Y, where each of the *n* sums is  $X_i + Y_i$ , has the following definition:

$$s_{x+y}^2 = \frac{\sum_{i=1}^{n} [X_i + Y_i - (\bar{X}_i + \bar{Y}_i)]^2}{n-1}.$$
 (7.15)

The terms inside the brackets in Eq. (7.15) can be rearranged to produce

$$s_{x+y}^2 = \frac{\sum_{i=1}^{n} \{(X_i - \bar{X}_i) + (Y_i - \bar{Y}_j)\}^2}{n-1}.$$
 (7.16)

If the bracketed expression in the numerator of Eq. (7.16) is expanded and the summation sign is distributed over the terms after expansion, one obtains

$$s_{x+y}^2 = \frac{\sum\limits_{i=1}^{n} (X_i - \bar{X})^2}{n-1} + \frac{2\sum\limits_{i=1}^{n} (X_i - \bar{X})(Y_i - \bar{Y})}{n-1} + \frac{\sum\limits_{i=1}^{n} (Y_i - \bar{Y})^2}{n-1}.$$
 (7.17)

You will recognize immediately that the first and last terms on the

right-hand side of Eq. (7.17) are  $s_x^2$  and  $s_y^2$ , respectively. The middle term is simply 2 times the covariance of X and Y,  $s_{xy}$ . Thus,

$$s_{x+y}^2 = s_x^2 + s_y^2 + 2s_{x+y} (7.18)$$

One way of denoting  $r_{xy}$  is by  $s_{xy}/(s_xs_y)$ . Obviously then,  $s_{xy} = r_{xy}s_xs_y$ . Therefore, replacing  $s_{xy}$  by an equivalent expression gives

$$s_{x+y}^2 = s_x^2 + s_y^2 + 2r_{xx}s_xs_y. (7.19)$$

Equations (7.18) and (7.19) relate the variance of the sum of two arrays of scores to the variance of each array and the covariance of the arrays.

An important special case of Eq. (7.19) is that in which X and Y are uncorrelated, i.e.,  $r_{xy} = 0$ . If this is true, then

$$s_{x+y}^2 = s_x^2 + s_y^2. (7.20)$$

What is an expression equivalent to  $s_{r-s}^2$ ?

$$s_{x_{-y}}^{2} = \frac{\sum_{i=1}^{n} \{(X_{i} - Y_{i}) - (\bar{Y}_{i} - \bar{Y}_{i})\}^{2}}{n - 1}$$

$$= \frac{\sum_{i=1}^{n} (X_{i} - \bar{X}_{i})^{2} + \sum_{i=1}^{n} (Y_{i} - \bar{Y}_{i})^{2} - 2\sum_{i=1}^{n} (X_{i} - \bar{X}_{i})(Y_{i} - \bar{Y}_{i})}{n - 1}$$

$$= s_{x_{i}}^{2} + s_{y_{i}}^{2} - 2s_{x_{i}} = s_{x_{i}}^{2} + s_{x_{i}}^{2} - 2r_{x_{i}, x_{i}, x_{i}}$$

This is an interesting development. The variance of the differences between a group of persons' scores on X and Y equals the variance of X plus the variance of Y minut where the covariance of X and Y (or twice  $r_{ey}$  times  $s_e$  and  $s_e$ ). Again, if X and Y are uncorrelated, then

$$s_{x-y}^2 = s_x^2 + s_y^2$$
. (7.21)

You may find it difficult to reconcile Eq. (7.20) with Eq. (7.21) and this is not surprising. Perhaps the following argument will help:

If X and Y are uncorrelated, then surely X and Y are uncorrelated, since Y is just a linear transformation of Y (see Sec. 7.6). If we let  $Y^*$  stand for Y, we know that  $z_{***}^* = z_*^* + z_*^*$  because X and  $Y^*$  are uncorrelated. The effect of multiplying Y by -1 to form  $Y^*$  gives  $Y^*$  a mean of -Y but leaves the variance at  $z_*^*$ . Therefore,  $z_{****}^* = z_*^* + z_*^*$ ; but of course,  $X + Y^*$  is just X - Y. Hence,  $z_*^* = z_*^* + z_*^* = z_*^* + z_*^*$ ; but of

The above results are not only useful in more advanced statistics—they can be illuminating also. For example, we know that

$$s_{x+y}^2 - s_x^2 - s_y^2 = s_x^2 + s_y^2 + 2r_{xy}s_xs_y - s_x^2 - s_y^2 = 2r_{xy}s_xs_y$$

Therefore, if we divide the above equation through by 25,5,, we find that

$$\frac{s_{x+y}^2 - s_x^2 - s_y^2}{2s_x} = r_{xy}.$$

Suppose one has three variables X, Y, and Z. What would be the variance of the sums formed by adding together a person's scores on these three variables to secure the sum X. Y, Y, Z.?

$$s_{z+y+z}^{2} = \frac{\sum_{i=1}^{n} [(X_{i} + Y_{i} + Z_{i}) - (\overline{X}_{i} + \overline{Y}_{i} + Z_{i})]^{2}}{n - 1}.$$
 (7.22)

The terms in brackets in Eq. (7.22) can be rearranged to yield

$$s_{z+y+z}^2 = \frac{\sum_{i=1}^{n} [(X_i - \bar{X}) + (Y_i - \bar{Y}) + (Z_i - Z)]^2}{n-1}.$$
 (7.23)

The numerator of Eq. (7.23) is a trinomial squared. You may recall from high-school algebra that  $(a+b+c)^2=a^2+b^2+c^2+2ab+2ac+2bc$ . Hence.

$$\begin{split} s_{z+y+z}^2 &= \frac{\sum (X_t - \bar{X})^2}{n-1} + \frac{\sum (Y_t - \bar{Y})^2}{n-1} \\ &+ \frac{\sum (Z_t - Z)^2}{n-1} + \frac{2\sum (X_t - \bar{X})(Y_t - \bar{Y})}{n-1} \\ &+ \frac{2\sum (X_t - \bar{X})(Z_t - Z)}{n-1} + \frac{2\sum (Y_t - \bar{Y})(Z_t - Z)}{n-1}. \end{split}$$

All of the terms to the right of the equal sign in the above equation are variances or covariances. The entire expression can be reduced to

$$s_{xy}^2 + s_y^2 + s_y^2 + s_z^2 + 2s_{xy} + 2s_{xz} + 2s_{yz}$$

which is the same as

$$s_{-tube}^2 = s_x^2 + s_y^2 + s_z^2 + 2r_{xy}s_xs_y + 2r_{xz}s_xs_z + 2r_{yz}s_ys_z$$

The problems of variances of sums of and differences between variables are very important in intermediate and advanced statistics. A thorough understanding of these concepts is requisite to most work in mental test theory, factor analysis, and many other areas which lie outside of statistics. You would do well to thoroughly understand the material in this section before moving on. If you need further instruction in this area, Edwards (1964, pp. 15–23) should be helpful.

#### 7.10 OTHER MATERIAL ON CORRELATION

This chapter does not contain the entire treatment of correlation in this text. Chapter 8 has as its subject the problem of least-squares estimation, which is closely related to correlation (see Sec. 8.4). In Chapter 9, correlation coefficients are presented for correlating scores that are nominal and ordinal. These three chapters by no means exhaust the subject of measuring relationships. Those aspects of the subject which have been slighted here can be covered adequately by consulting Ezekiel and Fox (1963), DuBois (1957), and Kruskal (1958).

#### PROBLEMS AND EXERCISES

- 1. Prove that the correlation of X and Y equals +1 when  $z_x = z_y$ .
  - (Hint: Start with the equation  $r_{xy} = \sum z_x z_y / (n-1)$ ). Since a person's z score on X is assumed to be identical to his z score on Y, substitute  $z_x$  for  $z_y$  in the formula for  $r_{xy}$ . Then prove that  $\sum_{i=1}^{n} z_{x}^{2}/(n-1) = +1$ . See Prob. 8 in Chapter
- 2. Brown calculated the covariance of height in inches X and running speed in seconds Y. He obtained a value of 27.60 on a sample of 50 students. Smith calculated the covariance of height in feet  $X(S_2^{1\prime}, S_{12}^{1\prime},$  etc.) and running speed Y from the same original data that Brown collected. Smith obtained a value of 2.30. Will Brown or Smith obtain a larger value for the correlation between X and Y with this one set of measures?
- 3. Indicate whether the covariance and, hence, the correlation of the two designated variables would probably be positive or negative in the population of all elementary-school pupils in the United States:
  - a. X, height in inches; Y, weight in pounds.
  - b. X, age in months between 6 and 16 years; Y, time in seconds required to
  - c. X, reading achievement in grade-placement units; Y, arithmetic achievement
  - d. X, I.Q. of student; Y, "citizenship" rating of student on a 10-point scale
  - e.  $\acute{X}$ , arithmetic achievement in grade-placement units; Y, number of days absent from school during the year.
- 4. For a particular set of data,  $s_x = 5$  and  $s_y = 4$ . What is the largest that  $s_{xy}$ could possibly be? [Hint:  $r_{xy}$  cannot be larger than +1;  $r_{xy} = s_{xy}[(s_x s_y)]$ .]
- 5. The correlation of X with Y is .60; the correlation of X with Z is -.80. Is X more closely linearly related to Y or to Z?

- 6. A researcher demonstrated a correlation of -.52 between average teacher salary X and the proportion of students who drop out of school before graduation Y, across 120 high schools in his state. He concluded that increasing teachers' salaries would reduce the "drop-out rate." Comment on his conclusion.
- 7. a. Find the value of the correlation coefficient r for the following data:

Person	X	}	
1	100	28	
2	90	25	
3	126	19	
4	112	24	
5	80	23	
6	115	21	
7	105	27	
8	110	25	
9	99	26	
10	97	25	
11	87	23	
12	76	18	
13	100	29	
14	80	20	
15	120	18	

- b. Plot a scatter diagram for the above data.
- c. Does the relationship between X and Y—if there is a nonzero one—appear to be predominantly linear or curvilinear?
- 8. Compute the r for data set a, below, and the r for data set b. Why do the r's differ in magnitude?

a.	Person	IQ		ь.	Person	Test score	Arith.
•	number	Test A	Test B		number	gen. vocab.	reasoning
		80	83		1	96	104
	ż	105	101		2	111	121
	3	121	117		3	89	84
	4	93	100		4	107	91
	5	99	96		5	102	114
	6	107	112		6	115	96
	7	119	123		7	98	109
	8	103	99		8	83	94
	9	102	110		9	104	116
	10	115	110		10	100	86
	11	87	81		11	117	101
	11	96	98		12	94	99

It has been shown that women tend to score much higher (i.e. have more "accepting" attitudes) than men on the Minnesota Teacher Attitude Inventory.

A researcher correlated the MTAI scores of a group of 100 experienced secondary teachers with the number of students each teacher failed in a year. He obtained an r of -3.9. He concluded that teachers tend to fail students because they do not have "accepting," attitudes toward students. Comment on the researcher's methods and conclusions.

$$\begin{aligned} 10. \ &\sum_{t=1}^{n} (X_{t} - \bar{X})(Y_{t} - \bar{Y}) = \sum_{t=1}^{n} [X_{t}(Y_{t} - \bar{Y}) - \bar{X}_{t}(Y_{t} - \bar{Y})] \\ &= \sum_{t=1}^{n} X_{t}(Y_{t} - \bar{Y}) - \bar{X}_{t} \sum_{t=1}^{n} (Y_{t} - \bar{Y}) = \sum_{t=1}^{n} X_{t}(Y_{t} - \bar{Y}) - \bar{X}_{t}(0). \end{aligned}$$

In words, the sum of the cross products of the X and Y scores, both in deviation score form, equals the sum of the cross products of X, not in deviation score form, and the Y's in deviation score form. This seems paradoxical. Has a mistake been made in the proof? Or is it truly a fact?

11. Is it true that 
$$\sum_{i=1}^{n} X_i(Y_i - \bar{Y}) = \sum_{i=1}^{n} Y_i(X_i - \bar{X})$$
?  
Hint: Show that  $\sum_{i=1}^{n} [X_i(Y_i - \bar{Y}) - Y_i(X_i - \bar{X})] = 0$ .

### PREDICTION AND ESTIMATION

#### 8.1 PRELIMINARIES

The concepts and procedures involved in the simplest form of statistical prediction are best illustrated by means of a few elementary notions in analytic geometry. We shall develop little more than the idea of a two-dimensional coordinate system and the equation for a straight line in this system.

A cartesian\* coordinate system is depicted in Fig. 8.1. The axes of the coordinate system—the mutually perpendicular X and Y lines—divide the plane (the flat, two-dimensional surface having length and width, but not depth) into four quadrants: I, II, III, and IV. This coordinate system is a means of marking off the plane in such a way that every point in the plane can be identified by a pair of numbers, (X, Y). The point (0, 0) is called the origin of the system and lies at the point where the X and Y lines cross, i.e., where the two axes intersect. The first number of any pair is the distance

Named after the French philosopher-mathematician René Descartes.





FIG. 8.1 Two-dimensional (X and Y) cartesian coordinate system.

one must travel horizontally from the origin (the X distance) to reach the point, and the second number is the distance the point lies retrically from the origin. The point A in Fig. 8.1 corresponds to the pair of numbers (2, 2). The first number is called the X-coordinate; the second number is called the X-coordinate. The point B corresponds to the pair of numbers  $\{-2, 1\}$ ; it lies two units to the left of the origin along the X axis and one unit above the origin along the Y axis. Points in quadrant I correspond to pairs of numbers both of which are positive; points in quadrant II correspond to pairs of numbers the first of which is negative, the second of which is positive. What are the signs of the numbers describing a selection of the point of the numbers describing a selection of the numbers describing a selection of the numbers describing a selection of the numbers described as the signs of the numbers descri






FIG 8.2 Graph of the straight line Y = 2X + 1.

quadrant? The fourth quadrant? What is the pair of numbers that determine the location of point C in Fig. 8.1?

You should realize that any point in a plane corresponds to a pair of real numbers and that any pair of real numbers identifies a point. Thus, (-100, 3.67) determines a point, as does (-41.65, 214.6).

Eventually, we shall arrive at a method of predicting a set of scores that uses a straight line in a plane to describe the set of predicted scores. It will be useful to know the manner in which any straight line in a plane can be completely described by a simple equation.

In Fig. 8.2 the straight line L crosses the Y axis at the point (0, 1), and the X axis at the point (-0.5, 0). For each unit on the X axis that the moving point which describes the line moves to the right, it rises two units on the Y axis. The following points lie on the line L: (-2, -4), (-1, -1), (0, 1), (1, 3), etc. The value of Y in the description of the point (X, Y) is systematically related to the value of X. For the line L in Fig. 8.2, the Y value of any point on the line equals twice the X value plus I, i.e., Y = 2X + 1. The equation Y = 2X + 1 is the equation for the straight line in Fig. 8.2. The number 1 is called the Y interrupt because it is the distance above the X axis at which the line intersects with the Y axis. The number 2 is the slope of the straight line. The slope is the number of units the line rises for each unit of movement to the right of the X axis, here 2:1.

The equation  $Y=b_1X+b_0$  is called the "general equation for a straight line." It says simply that the pairs of points (X,Y) that lie on any straight line are related in such a way that for any X value the Y value paired with it can be found by multiplying X by some number  $b_1$ , and adding a second number  $b_0$  to this product. This is a linear transformation of X to secure Y;  $b_1$  is a multiplicative constant, the same for every X, and  $b_0$  is an additive constant, the same for every X.

# 8.2 THE PROBLEM OF ESTIMATING Y FROM X (OR X FROM Y)

Given an individual's score on characteristic (variable) X, what information can be gained about his score on characteristic (variable) Y? Some examples of the estimation problem are:\*

 How, and how well, can we predict college English grades from high-school English grades? (The high-school grades precede the college grades, so we can predict the latter.)

 When X precedes the Y we wish to estimate from it, then we predict (i.e., tell in advance) Y from X, once we know the relationship between X and Y based on an earlier group.

- How well can we estimate Stanford-Binet IQ scores from California Test of Mental Maturity IQ scores? (No antecedent-consequent order is implied, so this is estimation to determine how nearly equivalent the z scores of each examinee are on the two tests.)
- 3. How well can we predict income at age 35 from rank in high-school graduating class?
- 4. How well can we estimate achievement from intelligence?

To derive a means of estimating the score of a person on one variable (which we shall denote by Y) from a different variable (X), we must know how X and Y are related. The variable we wish to estimate is called the dependent variable (Y), and the variable that will be used to estimate it is the independent variable (X). For example, we might wish to predict achievement in ninth-grade mathematics (dependent variable Y) from a group intelligence test given at the end of grade eight (independent variable X). Measures of Y might be scores on a 50-item achievement test in ninth-grade math. We must first gather data on some number n of students whose meticine we test in grade eight and whose achievement in ninth-grade math we measure. We next establish an equation that relates X and Y in this group; we would then use this equation in the future with students whose score on X we know and whose score on Y we would like to estimate.

Illustrative data can be tabulated as in Table 8.1, and graphed as in Fig. 8.3. It is apparent in Table 8.1 that five sums and the number of pairs (there, n=20) are required to determine  $b_1$ , which is the slope of the straight regression line needed to estimate Y (mathematics scores in grade 9) from X (IQ's in grade 8). The five statistics, shown at the upper right section of Table 8.1, are: the sum of the  $20 \ X$  scores—that is,  $X_1 + X_2 + \dots + X_n = 2165$ ; the sum of the  $20 \ Y$  scores (824); the sum of the  $30 \ Y$  scores (324); the sum of the  $30 \ Y$  scores (324); and the sum of the  $30 \ Y$  scores, which is  $30 \ Y$  scores, which is  $30 \ Y$  scores, which is  $30 \ Y$  scores, which is

$$X_1Y_1 + X_2Y_2 + \ldots + X_nY_n = 95(33) + 100(31) + \ldots + 118(48)$$
  
= 89,715.

By means of these six statistics one finds that  $b_1=.708$ , and that  $b_0=-35.441$ . Thus, the equation for estimating Y's from X's is .708X-35.441. The straight line that follows this formula is shown in Fig. 8.3. It has a slope of +.708, because for every increment of .708 in Y there is an increment of 1.000 in Y: .708/1.000=.708.

 $b_0 = -35.441$  means that if Fig. 8.3 were drawn with the X axis extending all the way back to X = 0 1Q, we would see that the regression line at the X of 0 crosses the Y axis at -35.441, the cartesian pair there being (0, -35.441).

137

TABLE 8.1 DATA FOR THE DETERMINATION OF A PREDICTION LINE

X Independent variable (IQ in grade 8)	Y Dependent variable (math score in grade 9)	Calculations
95	33	$\sum X = 2165 \qquad \sum Y = 824$
100	31	~
100	35	$\sum X^2 = 235,091$ $\sum Y^2 = 34,442$
102	38	21 = 200,001
103	41	$\sum XY \approx 89,715$
105	37	n=20
106	37	
106	39	$n \sum XY - \sum X \sum Y$
106	43	$b_1 = \frac{n \sum XY - \sum X \sum Y}{n \sum X^2 - (\sum X)^2}$
109	40	
		$\frac{20(89,715) - (2165)(824)}{20(235,091) - (2165)^2} = \frac{10,340}{14,595} = .708.$
110	41	20(235,091) (2165)2 14,595708
110	44	824 /2165\
111	40	$b_0 = P_1 - b_1 X_1 = \frac{824}{20}708 \left(\frac{2165}{20}\right) =$
112	45	20 (20)
112	48	-35.441
		Least-squares prediction line:
114	45	$\hat{Y} = .708X35.441$
114	49	- 1.004 92141
115	47	Other descriptive statistics:
117	43	•
118	48	$s_x = 6.198, s_y = 5.095, s_{xy} = 27.211$ $r_{xy} = .861$

What criteria led to the formulas for  $b_1$  and  $b_0$  used in Table 8.1.? This is an important question that deserves a detailed answer, as given below. Suppose we found an equation for predicting Y from X that had satisfied.

Suppose we found an equation for predicting I from X that has satisfactory properties. We would have two constants,  $b_1$  and  $b_0$ , such that multiplying X by  $b_1$  and adding  $b_0$  would give us an estimated value of Y. That is,

$$\hat{Y}_i = b_i X_i + b_0,$$

the predicted value of Y for the ith person, denoted by  $\hat{Y}_i$ , equals  $b_i$  times his X score plus  $b_o$ . Obviously,  $\hat{Y}_i$  will not always equal  $Y_i$ , i.e., even with the "best" straight-line prediction equation we shall usually make errors in predicting Y from X. We say, then, that  $Y_i = b_i X_i + b_0 + e_i$ , where  $e_i$  is an error of estimating Y from  $\hat{Y}$  for the ith person:

$$e_i = Y_i - \hat{Y}_i. \tag{8.1}$$

Another name for e<sub>i</sub> is the error of estimate. The nature of the error of estimate in the prediction problem is illustrated in Fig. 8.4 for a person




FIG. 8.3 Scatter diagram of X (IQ in grade 8) and Y (mathematics achievement in grade 9) for 20 students (see Table 8.1), with least-squares prediction line superimposed.

whose actual Y score is 4.36 units below the estimated Y for him that lies on the regression line.

How should we go about choosing  $b_1$  and  $b_0$ ? Usually,  $b_1$  and  $b_0$  are found by choosing them so that

$$\sum e_i^2 = e_1^2 + e_2^2 + \ldots + e_2^2$$

is as small as possible—that is, the sum of the squared errors of estimate is minimal.

This choice of a criterion for selecting  $b_0$  and  $b_1$  is arbitrary to some extent. It has historical precedent, and computationally it is convenient. Moreover, it is preferred on statistical inferential grounds, but just why it is preferred on these grounds cannot be known until you have learned considerably more statistics. However, the criterion of least squares (i.e., minimal sum of squared errors of estimately is not the only criterion possible. Although there has never been a serious contender with it for the role of criterion for fitting the regression line, other criteria exist. One is that  $b_0$  and  $b_1$  should be chosen so that the sum of the absolute values of errors made in prediction is as small as possible, i.e., so that  $|c_1| + \dots + |c_n|$  is minimized. This criterion leads to a "median-regression line." (You will recall that the median of a group of scores is the point around which the sum of the absolute deviations of the scores from that point is minimal.) Although the median-regression line is easily eacluated, it does not have the inferential theoretical superstructure that the least-squares regression line has

The use of the criterion of least squares for establishing a prediction line is over 150 years old. Karl Gauss (1777-1855), a famed German mathematician, physicist, and astronomer, is generally credited with having invented the criterion of least squares. In one form or another it underlies a large portion of theoretical and applied statistical work.

We noted earlier that e, is the difference between a person's actual Y score and the Y score we predict for him:

$$e_i = Y_i - \hat{Y}_i = Y_i - (b_1 X_i + b_0).$$
 (8.2)

We choose  $b_1$  and  $b_0$  so that

$$[Y_1 - (b_1X_1 + b_0)]^2 + [Y_2 - (b_1X_2 + b_0)]^2 + \ldots + [Y_n - (b_1X_n + b_0)]^2$$

is as small as it can possibly be. The exact manner in which one determines the values of  $b_1$  and  $b_0$  that minimize the above quantity is too complicated for us to detail. (See McNemar, 1962, pp 119–24.) We shall simply report the results here, and in Sec. 8.5 present a verification of the reported solutions.

b, is given by the following equation:

$$b_1 = \frac{n \sum_{i=1}^{n} X_i Y_i - \left(\sum_{i=1}^{n} X_i\right) \left(\sum_{i=1}^{n} Y_i\right)}{n \sum_{i=1}^{n} X_i^2 - \left(\sum_{i=1}^{n} X_i\right)^3},$$
 (8.3)




FIG. 8.4 Illustration of the error of estimate, e, for the person scoring 100 on variable X and 31 on variable Y.

bo is given by the equation

$$b_0 = \overline{Y}_1 - b_1 \overline{X}_1. \tag{8.4}$$

The values of  $b_1$  and  $b_0$  found in this way give the "best" prediction equation in the sense that the sum of the squared differences between the  $Y_i$  and the  $\hat{Y}_i = b_1 X_i + b_0$  is as small as it can possibly be for these data. In the last section in this chapter a simple algebraic verification that this is in fact true is presented.

Suppose that we seek the "best" equation for estimating Y from X by means of a straight line best fitting (in the least-squares sense) the dots of the scatter diagram and that we have the X and Y scores for 20 persons as given in Table 8.1.

Nothing has been said to this point about whether the dependent and independent variables are normally distributed or distributed in any other special way. No knowledge of the shapes of the frequency distributions of X and Y was needed to derive the least-squares regression coefficients  $b_0$  and  $b_1$ . Equations (8.3) and (8.4) for  $b_0$  and  $b_1$  produce the straight line that minimizes the sum of squared residuals regardless of the nature of the scatter diagram of the X and Y scores.

If we make some plausible assumptions about the distributions of large numbers of X and Y scores, we are likely to be rewarded by being able to perform a more penetrating prediction study. Our study does gain much, in fact, if we assume that X and Y have a bivariate normal distribution (see Sec. 6.7). It is assumed, then, that the n pairs of X and Y scores in hand form a random sample of a very large collection of X and Y scores that have a bivariate normal distribution.

The properties of the bivariate normal distribution now of importance are:

- The population means of the Y's for each separate value of X lie on a straight line;
- For any single value of X the associated Y scores are normally distributed;
- 3. For any single value of X the associated Y scores have variance  $\sigma_{YZ}^2$ , and this variance is the same for all of the X scores.

If we can be confident that the n pairs of X and Y scores are from a bivariate normal distribution, then property 1 above tells us that the use of a straight line for predicting Y from X was reasonable and cannot be improved upon by any curved line. Properties 2 and 3 above can be combined into a very useful technique that adds greatly to Y-from-X prediction and estimation problems. This technique will be developed next.

# 8.3 HOMOSCEDASTICITY AND THE STANDARD ERROR OF ESTIMATE

Property 3 of the bivariate normal distribution leads us to believe that, if we have, say, 19 persons with a score of 75 on X and 21 persons with a score of 80 on X, the variances of the two groups of associated Y scores should be about the same. This condition of equal variance of Y scores for each value of X is known as homoscedasticity (the roots of this word mean equal spread). The scatter diagram in Fig. 8.5 should help you gain an understanding of this condition.

It is important to note that homoscedasticity is a property of very large bodies of bivariate data. One should not expect equality of variances of Y scores for any two values of X when the  $n^2$ s are small, say of the order of 100 or less. For n's of 19 and 21 the variances of the Y scores for X = 75 and X = 80 in Fig. 8.5 are  $s_{n-2}^2 = 5.54$  and  $s_{n-2}^2 = 6.85$ . These two variances are not equal, but they are reasonably close. With such small numbers of persons we cannot ascertain very well whether the condition of homoscedasticity is satisfied, but at least it seems somewhat plausible for X equals 75 and 80 after inspection of the data of Fig. 8.5.

Obviously the sizes of the errors made in estimating Y from X are an indication of the accuracy of estimation. For the data in hand, i.e., the n pairs of X and Y scores, the differences between the actual Y scores and the predicted Y scores are measures of the errors that would result if X is used to estimate Y. These errors are called errors of estimate. The formula for the error of estimate for the thip person is

$$e_i = Y_i - \hat{Y}_i = Y_t - b_t X_t - b_\theta.$$
 (8.5)

Y
20

15

 $s_{p,75}^2 = 5.54$ 
 $s_{p,0}^2 = 6.65$ 

Traini for a X and 100

wariance

FIG. 8.5 Scatter diagram for 19 persons scoring 75 on X and 21 persons scoring 80 on X which exhibits nearly the same variance of Y scores for both values of X.

In words, the ratio of the variance of the predicted Y scores,  $\hat{Y}_i$  to the actual Y scores is equal to the square of the correlation coefficient between X and Y. Notice that this ratio tells nothing about the direction of the relationship; the ratio is never negative. For the data in Table 8.1,  $s_i^2 = 25.958$  and  $s_j^2 = 19.243$ . The ratio 19.243/125.958 equals .741, which corresponds to the square of the value of  $r_{sr}$  for these same data. It is often said in an attempt to explain the meaning of  $r_{sr}$ , that  $r_{sr}^2$  is the "amount of variance in Y explained by variance in X." This is ambiguous and nearly meaningless language without explicit definitions of what it means for a variable to have its "variance explained," and yet explicit definitions are usually lacking. Such explanations" of  $r_{sr}$  are attempts to verbalize Eq. (8.17) beyond the italicized sentence above.

The most significant formula for "understanding" what the correlation coefficient describes is Eq. (8.17). We shall try to interpret the meaning of  $r_{xy}^2$  in a slightly different way.

Suppose we wanted to predict each person's score on variable Y but no variable X was available. The predicted score that minimizes the sum of the squared errors of prediction in predicting Y happens to be Y for each person. That is, if no X variable is available and nothing is known about how individuals score on Y, the best prediction can be attained by predicting that everyone will score at the mean Y, on Y. A measure of the goodness of such prediction without an X variable is given by

$$\frac{\sum_{1}^{n}(Y_{i}-\overline{Y})^{2}}{n-1},$$

which happens to equal s2.

Now suppose that knowledge of a person's status on X is available for predicting his Y score. The sum of the squared errors of prediction made in predicting Y from X with the least-squares regression line is given by

$$\frac{\sum_{i=1}^{n}(Y_{i}-\hat{Y}_{i})^{2}}{n-1},$$

which is s2.

Now we know that  $s_0^2 + s_1^4 = s_2^2$ . The quantity  $s_2^2$  is the error made in prediction before knowledge of X is used;  $s_2^2$  is the error made when knowledge of X in used. The amount of error eliminated from  $s_2^2$  by knowledge of X must then equal  $s_2^2$ . Recall that

$$\frac{s_{\theta}^{2}}{s_{x}^{2}} + \frac{s_{x}^{2}}{s_{x}^{2}} = 1.$$

Noting that  $s_y^2/s_y^2 = r_{xy}^2$ , the above equation can be written as:

$$r_{xy}^2 = 1 - \frac{s_x^2}{s_y^2} = \frac{s_y^2 - s_x^2}{s_y^2} = \frac{s_y^2}{s_y^2}.$$
 (8.18)

The following interpretation of  $r_{xy}^2$  can be extracted from Eq. (8.18):

The value  $r_{ss}^2$  equals the proportion of the variance of squared errors,  $s_{ss}^2$ , made in predicting Y without knowledge of X that is eliminated when Y is predicted by the least-squares method from a knowledge of X.

For example, suppose  $r_{vv}^2 = .75$  and  $s_v^2 = 100$ . We know, then, that  $s_v^2 = 25$ . The total error that would be made in predicting Y without knowledge of X, in which case everyone is predicted to have a Y score of Y, is equal to

$$\frac{\sum_{1}^{n} (Y_i - \overline{Y})^2}{n-1} = s_y^2 = 100.$$

The total error made in predicting Y from X with the least-squares regression line is

$$\frac{\sum_{i=1}^{n} (Y_i - \hat{Y}_i)^2}{n-1} = s_e^2 = 25.$$

Therefore, 100 - 25 = 75 is the percent of error eliminated from  $s_y^2$  by knowledge of X. We note also that  $r_{xy}^2 = .75$ .

#### 8.5 VERIFICATION THAT THE LEAST-SQUARES CRITERION IS SATISFIED BY $b_1$ AND $b_0$

This algebraic derivation is placed in Appendix C because it will probably not be of interest to all students. Earlier we stated without proof or verification that Eqs. (8.3) and (8.4) give the values of  $b_1$  and  $b_0$  that minimize  $\sum \{Y_i - (b_0 + b_1 \lambda_0)\}^n$ , the sum of squared errors of estimate. Appendix C gives a relatively simple algebraic verification of the fact that Eqs. (8.3) and (8.4) for  $b_1$  and  $b_0$  produce the smallest possible sum of squared errors and (8.4) for  $b_1$  and  $b_0$  produce the smallest possible sym of squared errors while imparting a deeper understanding of the least-squares criterion to all. while imparting a deeper understanding of the least-squares criterion to all. We emphasize that the proof is not the means by which Eqs. (8.3) and (8.4)

were originally derived. Our proof is an after-the-fact verification, given the Eqs. (8.3) and (8.4) for  $b_0$  and  $b_1$ .

# 8.6 MEASURING NONLINEAR RELATIONSHIPS BETWEEN VARIABLES; THE CORRELATION RATIO n<sup>2</sup>

This section appears here for the sake of completeness and logical continuity. You may find it more comprehensible after having studied Chapter 15 on the one-factor analysis of variance.

Although we have repeatedly pointed out that the Pearson product-moment r measures only the degree of linear relationship between X and Y, we have yet to indicate a descriptive measure to use when the relationship between X and Y is predominantly nonlinear. As an example of a nonlinear relationship, consider the data in Fig. 8.7 relating age Y to performance X on the Digit Symbol subtest of the Wechsler Adult Intelligence Scale. The data depicted in Fig. 8.7 are tabulated in Table 8.2. It is obvious from

TABLE 8.2 WAIS DIGIT-SYMBOL-SUBTEST SCALED SCORES OF 28 PERSONS IN EIGHT EQUALLY SPACED AGE GROUPS

				0110013			
10	14	18	Age to 1 22	rearest year 26	30	34	38
7 8 9 9 10 Age-group	8 9 10 11	9 10 11 12	11 11 12 12	9 10 11	8 9 9	7 9 10	8
means: 8.60	9.50	10.50	11.50	10 00	9.00	8.67	8.00
		Grand	mean of all	$scores = \frac{20}{2}$	$\frac{59}{8} \approx 9.61$		

Fig. 8.7 that the scores on this subtest rise in straight-line fashion from age 10 to a peak at age 22 and then decline rather rapidly.

A measure of the linear or nonlinear relationship between X and Y is denoted by  $\eta^2$  (read "eta squared"), also called the correlation ratio. The correlation ratio has the following definitional form:

$$\eta_{x,x}^2 = 1 - \frac{SS_{withlin}}{SS_{total}},$$
(8.19)




FIG. 8.7 Relationship between age and performance of 28 persons on the digit symbol subtest of the WAIS.

where  $SS_{total} = \sum_{i=1}^{n} (Y_i - \overline{Y})^2$ , i.e., the sum of squared deviations of each Y scores, and Y scores, and

For the first value attained by X, the corresponding Y scores are deviated around their own mean and the sum of the squared deviations is calculated. For example, in Table 8.2 this first sum of squares is  $(7-8.60)^2+(8-8.60)^2+(9-8.60)^2+(9-8.60)^2+(9-8.60)^2+(10-8.60)^2$ . This process is repeated for each distinct value that X assumes. For example, for X=14, one calculates  $(8-9.50)^3+(9-9.50)^2+(10-9.50)^2+(11-9.50)^2$ . For the last group, X=38, the sum of the squared deviations of the Y "scores" around their mean is  $(8-8)^3=0$ , since there is only one score. Finally, these sums of squared deviations for the separate values of X are summed. The result is  $SS_w$ . (If you are reading this section after having read Chapter 15, it will help you to note that  $SS_{withs}$  is the "sum of squares within" for a non-factor analysis of variance with unequal n's.)

For the data in Table 8.2, the value of  $SS_{total}$  is 54.68 and the value of  $SS_{estbhn}$  is 24.87. Hence the value of  $\eta_{x}^{2}$  is

$$\eta_{y,x}^2 = 1 - \frac{24.87}{54.68} = 1 - .455 = .545.$$

The following considerations bear on the interpretation of  $\eta_{s,r}^a$ . The coefficient  $\eta_{s,r}^a$ —notice that Y precedes the comma and X follows it—is a measure of the extent to which Y is predictable from X by a "best-fitting" line that may be either straight or curved.

It is important to note that  $\eta_{n,x}^2$  and  $\eta_{n,x}^2$  will generally have different values. This is contrary to our experiences with r, for which  $r_{n,x} = r_{p,x}$ . We can give the fact that  $\eta_{n,x}^2$  may not equal  $\eta_{n,x}^2$  some intuitive appeal with the data in Table 8.2. If a person's age is 10, his Digit Symbol score can be

fairly confidently predicted to be about 8.60. However, if we know that a person's Y score is 8, his age X may either be low, around 10, or high, around 38. Hence, Y can be predicted from X reasonably well; but X cannot be predicted well from Y. These facts are reflected in the values of  $\eta_{y,x}^2 = .545$  and  $\eta_{x,y}^2$ , which we have not calculated but which is close to

The value of  $\eta_{x,x}^2$  should be compared with the value  $r_{xy}^2$  instead of  $r_{xy}$ . We saw that  $r_{xy}^2 = 1 - (s_x^2/s_y^2)$ , which is equal to

$$r_{xy}^2 = 1 - \frac{\sum_{i=1}^{n} (Y_i - \bar{Y}_i)^2}{\sum_{i=1}^{n} (Y_i - \bar{Y}_i)^2}.$$
 (8.20)

Equation (8.20) shows  $r_{2}^2$  to be (1 minus the sum of squared deviations of the Y scores around a straight prediction line) divided by  $(n-1)s_x^2$ . Equation (8.19) shows  $\eta_{u,x}^2$  to be (1 minus the sum of squared deviations of the Y scores around a curved prediction line that passes through the mean of the Y's for each value of X) divided by  $(n-1)s^2$ . The curvilinear prediction line for predicting Y from X appears in Fig. 8.7.

As with  $r_{xy}^2$ ,  $\eta_{y,x}^2$  must always be less than or equal to I and greater than or equal to 0. Furthermore,  $\eta_{y,x}^2 > r_{xy}^2$ . The difference,  $\eta_{y,x}^2 - r_{xy}^2$ , is a measure of the degree of nonlinearity of a best-fitting line for predicting Y from X. (See Glass and Hakstian, 1969.)

#### 8.7 ADDITIONAL READING ON PREDICTION

A comprehensive and excellent textbook for the student who wishes to pursue the study of statistical prediction beyond this text is William Rozeboom's Foundations of the Theory of Prediction (Homewood, Illinois: The Dorsey Press, 1966). The study of the correlation ratio and related matters can be pursued in E. A. Haggard's Intraclass Correlation and the Analysis of Variance (1958).

# PROBLEMS AND EXERCISES

- 1. Draw the graphs of the lines corresponding to each of the following equations:
  - b.  $Y = \frac{1}{2} 1/3X$ .
  - c. 2Y = 4 + 2X. (Hint: Divide both sides by 2.)
  - d. Y = 2(1 + 2X). (Hint: Multiply the right side out.)
- 2. What are the values of the Y intercept  $b_0$  and the slope  $b_1$  of the straight line that passes through the points (X = 2, Y = 2) and (X = 3, Y = 1)?


3. A particular high school has determined a prediction equation for predicting college grade-point-average at the state university from high-school grade-point-average. The equation is Ŷ = .76 + .62X. Predict the college grade-point-average which would be earned by students with the following high-school grade-point-averages:

4. The following are arithmetic test scores and final-exam scores for 12 students in an elementary statistics course:

nentary statistics c Student number	X, Arithmetic test	Y, Final exan
	33	65
1	36	51
2	39	53
3	29	42
4	41	50
5	38	53
6	42	64
7	35	54
8	23	50
9	37	45
10	28	63
11	25	50
12		equation V. =

Find  $b_0$  and  $b_1$  in the least-squares prediction equation  $\hat{Y}_t = b_1 X_t + b_0$ . Predict the final-exam score of person 13, whose arithmetic test score was 36. (For your curiosity, his final-exam score was 68. What was the error of estimate?)

- 5. Find the value of the standard error of estimate s, for the data in Prob. 4.
- 6. For the data in Table 8.1, calculate the error of estimate,  $Y \hat{Y}$ , for that person whose X score is 100 and whose Y score is 35.
- 7. In a particular estimation problem, \( \tilde{F} = 6.4X + 32.5, \) and the standard error of estimate equals 4.25. Assuming that \( X \) and \( Y \) have a bivariate normal distribution, determine the limits within which lie the middle 50% of the \( Y \) scores of a population of persons whose score on \( X \) is 10, i.e., determine \( Y\_1 \) and \( Y\_2 \) in the following figure:



Partial, part, and multiple correlation will be discussed briefly in the latter sections of this chapter.

# 9.2 AN OVERVIEW OF THE CHAPTER

Four types of measurement of variables will be distinguished:

- I. Nominal-dichotomous measurement. The mere presence or absence of something is noted. The data are 0's and 1's. The order of the scoring is generally arbitrary. Examples: Republican (1) Democrat (0); sibling in school (1) no sibling in school (0); male (1) female (0); married (1) not married (0).
- 2. Nominal-dichotomous measurement with underlying normal distribution. It is assumed that more sophisticated, extensive, or refined measuring techniques could produce an approximately normal distribution of measures, but the data in hand tell only whether a person stands above (1) or below (0) some point in this normal distribution. If in a large group of students it was known only whether a child's IQ exceeded a score of 120 (call this 1) or fell below (call this 0), the 1's and 0's would be dichotomized data with an underlying normal distribution. Of course, it usually would be inefficient to discard original scores on persons and record 1's and 0's instead, and this not normally done (although in the early history of factor analysis it was computationally convenient). Generally, one assumes that a more highly developed measuring device would produce normally distributed scores even though the device in hand allows only dichotomous observation. Examples: IQ above 100 (1), below 100 (0); above average height (1), below 100 (2);
- 3. Ordinal measurement. The data are the n consecutive and untied ranks 1, 2,...,n. These ranks may be converted measures from some other sort of observations (as when the rank scores 136, 124, 97 are ranked 1, 2, 3), or they may be the first translation of perceptions into numbers (as when a judge ranks 10 contestants from most proficient, 1, down to least proficient, 10). Example: the 94 members of a graduating class are ranked from I to 94 on the basis of their 94 grade-point averages.
- 4. Interval or ratio measurement. A unit of measurement exists, e.g., degree on the centigrade scale, inch, day, etc., and (in the case of ratio measurement) a zero point on a scale of measurement corresponds to the absence (i.e., zero amount) of the variable being measured. Any real number may result

from the act of measurement, and differences between scores reflect on the differences in amount of the characteristic possessed. In this chapter we shall generally regard the interval and ratio scores to be approximately normally distributed, though of course in some situations they may not be. Examples: height; intelligence-test scores; achievement-test scores; certain measures in psychological experiments.

If measurement can be accomplished at the interval- or ratio-scale level, the scores can be transformed into any of the other three levels above. For example, suppose that ten students earned scores on a test of verbal reasoning, which is believed to produce an approximately normal distribution of measurements roughly on an interval scale: see Table 9.1.

TABLE 9.1 CHANGING TEN INTERVAL-SCALE SCORES INTO RANKS (ORDINAL SCALE) AND 0, 1 (NOMINAL SCALE)

Student no.	Verbal reasoning scores	Rank scores	Dichotomized scoring— normal distribution underlying
1	17	3	1
2	10	7	0
3	29	1	1
4	16	4	1
5	3	10	0
6	14	5	1
ž	9	8	o
Ŕ	26	2	j ,
9	6	9	0
10	11	6	0

The first column gives an identification number for the student. The raw scores that are assumed to come from an approximately normal distribution appear in the second column. These ten scores were ranked to obtain the third column. In the fourth column, the top five raw scores were given a 1; the bottom five raw scores were given a 0. It does not seem proper in this example for nominal-dichotomous scores to be assigned (though they would be identical to the scores in the fourth column), because we have reason to believe that a normal distribution underlies the dichotomy, or at least we know that a ten-category distribution does.

Where there are two sets of scores,  $X_i$  and  $Y_i$ , for each of n individuals, either X or Y could be measured in any one of the four manners outlined above. Thus there are  $4 \times 4 = 16$  possible pairs of descriptions of the measurement of two variables that are to be correlated. These 16 possible pairs of conditions on X and Y can be represented as in Table 9.2.

TABLE 9.2 TEN DIFFERENT BIVARIATE SITUATIONS RESULTING FROM COMBINATIONS OF FOUR SCALES OF MEASUREMENT

	1	Measurement of variable	le X	
Measurement of variable Y	Dichotomous	Dichotomized, with underlying normal	Ordinal	Interval
Dichotomous	A	(B)	(C)	(D)
Dichotomized, with underlying normal	В	E	(F)	(G)
Ordinal	С	F	н	(1)
Interval or ratio	D	G	1	3

It is necessary to consider only 10 of the 16 possible pairs, since the designation of the two variables being correlated as X or Y is entirely arbitrary (because  $r_{xx} = r_{xy}$ ). In terms of correlational theory, the six cells with letters in parentheses in Table 9.2 are the same as the cells with the same letters not in parentheses. Table 9.2 will be the structure upon which the following discussion of several specific measures of relationship will be built. One appropriate correlational measure for two interval or ratio variables (1) was the subject of the previous chapter. This is an instance in which the Pearson product-moment coefficient  $r_{xx}$  is used. We shall consider some of the remaining nine cases (A–1) below.

# 9.3 MEASURES OF RELATIONSHIP

#### Case A

Both variables yield nominal-dichotomous measures: the phi coefficient,  $\phi$ . In this case, both X and Y have been measured dichotomously. The data can be thought of as arranged in two columns of 0's and 1's where each row corresponds to one person's two scores. For example, 12 students in academic trouble in their sophomore year of college might be observed on the variables marital status and "dropped out of college"; see Table 9.3. Arbitrarily 1 means married and 1 means dropped out, with 0 for not married and 0 for remaining in school. One measure of the relationship between X and Y is simply  $r_{so}$ , the Pearson product-moment coefficient. The Pearson product-moment coefficient and is denoted by  $\phi$ . The value of  $\phi$  for the data in Table 9.3 is .507; but this was not found with the usual computation formula for  $r_{so}$ . That formula can be replaced with a still simpler but algebraically identical formula when the data on X and Y are dichotomous.

TABLE 9.3 ILLUSTRATION OF THE CALCULATION OF THE PHI COEFFICIENT,  $\phi$ 

Student no.	X Marital status (married, 1; not married, 0)	Y Attrition (dropped out, 1; remained, 0)	Calculations
	0	0	$p_x = .4167$ $q_x = .5833$
,	1	1	$p_v = .5000$ $q_v = .5000$
3	0	1	
4	0	0 1	$p_{xy} = .3333$
5	1	1	.3333 (.4167)(.5000)
6	1	0	
7	0	0	$\varphi = \sqrt{(.4167)(.5833)(.5000)(.5000)}$
8	1	1	= .507
9	0	0	
10	0	1	
11	0	0	
12	1	1	

Let  $p_x$  be the proportion of people scoring 1 on X;  $q_x$ , the proportion scoring 0 on X, will be equal to  $1-p_x$ . The proportion scoring 1 on Y is denoted by  $p_x$ , and  $q_y = 1-p_x$ . One more definition is necessary:  $p_{xx}$  is the proportion of people scoring 1 on both X and Y. If we were to operate on the formula for  $r_{xy}$  with these new definitions, we would find that it simplifies algebraically to the following convenient form:

$$\phi = \frac{p_{xy} - p_x p_y}{\sqrt{p_x q_x p_y q_y}}.$$
(9.1)

Equation 9.1 is a convenient way to compute the phi coefficient. The following derivation shows that  $\phi$  is the Pearson product-moment correlation between two variables, each of which is scored 0, 1.

If both the numerator and the denominator of Eq. (7.10) on page 114 are divided by n, the expression for  $r_{xy}$  becomes

$$r_{xy} = \frac{(1/n)\sum X_t Y_t - \overline{X}.\overline{Y}_t}{\sqrt{[(1/n)\sum X^2 - \overline{X}^2][(1/n)\sum Y^2 - \overline{Y}^2]}}.$$
 (9.2)

If X and Y are measured dichotomously, X and  $\overline{Y}$  are simply the proportions of I's on each variable. For example,

$$\vec{X}_{\cdot} = \frac{(0+1+0+\ldots+0+1)}{12} = \frac{5}{12},$$

in the illustration in Table 9.3. Thus, we can replace  $X_1$  and  $Y_2$  with  $p_2$  and  $p_3$ , respectively. Now,  $X_1Y_1$  will be different from zero only when the nth person scores 1 on each variable, in which case  $X_1Y_2 = 1 \cdot 1 = 1$ . Surely,

then,  $\sum X_i Y_i$  is simply a count of the number of persons scoring 1 on both X and Y; hence  $(1/n) \sum XY = p_{xx}$ . Since X is 0 or 1 and because  $0^x = 0$  and  $1^x = 1$ .

$$\frac{\sum X^2}{n} = \frac{(0^2 + 1^2 + \ldots + 0^2 + 1^2)}{n} = p_x.$$

Therefore, substituting such expressions into Eq. (9.2) produces

$$r_{zz} = \frac{p_{zx} - p_z p_z}{\sqrt{(p_z - p_z^2)(p_z - p_z^2)}} = \frac{p_{zx} - p_z p_z}{\sqrt{p_z (1 - p_z)p_z (1 - p_z)}} = \frac{p_{zx} - p_z p_z}{\sqrt{p_z q_z p_z q_z}}.$$
 (9.3)

When one has no particular interest in the proportions  $p_a$  and  $p_a$  and finds it more convenient to tabulate dichotomous bivariate data in a contingency table (a table showing the joint occurrences of pairs of scores two variables in a group),  $\phi$  can be calculated with a convenient raw score formula. The data in Table 9.3 can be represented as in Fig. 9.1.

Attrition	Montal	Totals	
Airition	Not married (0)	Marned (1)	10:0/3
Dropped out (1)	2	4	6
Remained (0)	5	1	6
Totals	7	5	12

FIG. 9.1 Contingency table for the data in Table 9.3.

Figure 9.1 presents the frequencies of persons, showing the four possible pars of characteristics in Table 9.3. For example, five persons in Table 9.3 were not married and remained in school during their sophomore year. The marginal totals for rows show the numbers of persons at both levels of "attrition," irrespective of their marital status. What is the interpretation of the column totals?

Suppose that in each cell of a contingency table of the above sort we substitute a letter for the actual frequencies so that we can deal with the computation of \$\phi\$ more generally. See Fig. 9.2.

The number of persons scoring 0 on X and 1 on Y is denoted by a. The total number of persons scoring 0 on X is a+c. The total number of persons represented in the table is n. How many people scored 0 on X?

It can be shown by substituting such equivalences as  $p_a = (b+d)/n$ ,  $p_a - (a-b)/n$ , and  $p_{aa} = b/n$  into Eq. (9.1) that the phi coefficient for the

		Variable X		Totals
		0		
Se Y	1	a	b	a+b
Variable Y	0	С	d	c+d
Tota	ıle.	0+0	b+d	n

FIG. 9.2 General form of a 2 × 2 contingency table.

data arranged in a contingency table like that of Fig. 9.2 is

$$\phi = \frac{bc - ad}{\sqrt{(a+c)(b+d)(a+b)(c+d)}}.$$
 (9.4)

Equation (9.4) was first derived by Karl Pearson in 1901 in a paper, published in the Philosophical Transactions of the Royal Society of London, that dealt with the correlation between variables that cannot be measured quantitatively. To illustrate the calculation of  $\phi$  by Eq. (9.4), the data in Fig. 9.1 will be used.

$$\phi = \frac{20 - 2}{\sqrt{(7)(5)(6)(6)}} = \frac{18}{6\sqrt{35}} = \frac{3}{\sqrt{35}} = .507.$$

This value of  $\phi$  is exactly equal to the value found in Table 9.3 for the same data. This equivalence must always result, because Eqs. (9.1) and (9.4) are algebraically equivalent.

# Properties of φ

The meaning of the phi coefficient in a correlational sense is quite clear; it is simply the Pearson product-moment coefficient of correlation for dichotomous data. However, the interpretation of  $\phi$  can present special problems. In Chapter 7 it was seen that for the purpose of interpreting  $r_{xy}$ , certain assumptions, several of which can be checked by construction a scatter diagram of the data, must be made. The dichotomous data involved in the calculation of  $\phi$  are quite a departure from the two-variable normal surface to which the interpretation of  $r_{xy}$  has reference. Consequently, it is wrong to think that the interpretive meaning of a  $r_{xy}$  of .60 and a  $\phi$  of .60 is the same.

Especially pertinent here is the fact that  $\phi$  can assume the value +1 only when a+b and b+d are equal (and hence a=d) in the 2 × 2 contingency table, i.e., when there are the same proportions of 1's on both X and Y.\* Notice that in Fig. 9.3, where  $p_x=.2$  and  $p_y=.4$ , no higher

<sup>•</sup> Or, what is the same, when a+c=c+d, and therefore a=d. This is one necessary condition for  $\phi=+1$ . The other is that a=d=0. The 0, 1, and 1, 0 quadrants must be empty.




FIG. 9.3 The greatest possible positive association when  $p_x=.2$  and  $p_x=.4$ .

positive relationship between X and Y than is shown can be obtained. All of the ten I's on Y are paired with I's on X, but because there are 20 I's on X but only 10 1's on Y, there must be 10 1's on X that are paired with 0's on Y. Y cannot be predicted perfectly from X because knowledge that a person scores a 1 on X does not rule out the possibility that his score on Y is 0. The value of \$\phi\$ in Fig. 9.3 is .41. In other words, so long as 10 of the 25 scores on Y are 1's and 20 of the 25 scores on X are 1's (or vice versa), the value of  $\phi$  cannot exceed .41. This is regarded by some statisticians as a great disadvantage in the use of  $\phi$ . In an attempt to sidestep this undesirable property of φ, Cureton (1959) and others have divided the obtained value of  $\phi$  by the maximum value  $\phi$  can attain,  $\phi$  maximum, for the distribution of 1's and 0's obtained. This produces a coefficient,  $\phi/\phi_{max}$ , that can always range between -1 and +1, regardless of how discrepant the proportions of I's on X and Y are. Carroll (1961) discussed  $\phi/\phi_{max}$  at length and found it wanting. The basic principle involved here is that a product-moment r can attain the value +1 only if the distribution of the X's has the same shape as that of the Y's. The value -1 can be obtained only if the shape of the distribution of the X's is exactly the reverse of that of the Y's; for  $\phi$  this would mean that  $p_x = q_x$ . These conditions of identical or reversed distributions are necessary to attain the limits +1 or -1, but they are by no means sufficient. Even if X and Y are distributed identically, they can correlate 0

#### Case B

X is dichotomous; Y is dichotomous, with an underlying normal distribution. We know of no one satisfactory coefficient for correlating variables measured in these ways. Perhaps for most such situations, likely to be infrequent, one should fortgo the normality assumption and calculate \$\phi\$. Quilling (1969) has discussed this problem.

#### Care C

X is dichotomous and Y is ordinal. The only available coefficients for this situation are due to Cureton (1956) and Glass (1966). (Also see Stanley. 1968b.) Because these coefficients build upon rationales first proposed for correlating two ordinal variables, they can best be discussed after Case I later in this section

# Case D

One variable yields nominal-dichotomous measures, the other yields interval or ratio measures: the point-biserial correlation coefficient. In this case. one variable is measured dichotomously (e.g., sex, marital status), and the measurement of the other variable produces a collection of scores with interval or ratio properties. For example, in a group of high-school juniors. we might observe whether a student drops out of college (0) during the freshman year or remains in college (1), and also measure his intelligence. Observation of the two variables, intelligence (X) and attrition (Y), will produce two scores for each student.

One means of describing the relationship between X and Y is simply to calculate the Pearson product-moment coefficient on the data as they are. Such a coefficient is called a point-biserial correlation coefficient and is denoted by red. (The term biserial refers to the fact that there are two series of persons being observed on X: those who scored 0 on Y, and those who scored 1 on Y. Both the name and the rationale for this coefficient are due to Karl Pearson. The expression product-moment biserial is sometimes used instead of point-biserial.) A simplified formula for reb follows:

$$r_{pb} = \frac{\overline{X}_{.1} - \overline{X}_{.0}}{s_x} \sqrt{\frac{n_1 n_0}{n(n-1)}},$$
 (9.5)

where  $X_1$  is the mean on X of those who scored 1 on Y,  $\vec{X}_0$  is the mean on X of those who scored 0 on Y. s, is the standard deviation of all n scores on X. n, is the number of persons scoring 1 on Y,  $n_0$  is the number of persons scoring 0 on Y, and

 $n = n_1 + n_2$ 

Equation (9.5) represents an algebraic simplification of the Pearson product-moment correlation coefficient formula when Y is a dichotomous variable. Equation (9.5) is only one of several simplifications that could have resulted. Each of the following formulas is equivalent to Eq. (9.5), and one of them will probably be more convenient than the others for a particular problem.

$$r_{pb} = \frac{\overline{X}_{,1} - \overline{X}_{,}}{s_{,s}} \sqrt{\frac{n_1 n}{n_0 (n-1)}},$$
 (9.6)

where  $\bar{X}$  is the mean of all n scores on X.

$$r_{ph} = \frac{\overline{X}_{.} - \overline{X}_{.0}}{s_{x}} \sqrt{\frac{n_{0}n}{n_{1}(n-1)}}.$$
 (9.7)

As is particularly evident from consideration of Eq. (9.5),  $r_{gs}$  is a measure of the difference between the average scores on X of the persons scoring 1 on Y and the persons scoring 0 on Y. Because  $r_{gs}$  is nothing more than the product-moment correlation coefficient calculated on particular types of data, it must take on some value from -1 to +1, inclusive. When those scoring 1 on Y have the same average value on X as those scoring 0 on Y,  $r_{gs}$ , will be zero. Of course,  $r_{gs}$  is not defined if either  $r_{gs}$  or  $r_{gs}$  equals  $r_{gs}$  "covariation" cannot be studied if variation (on Y, in this case) does not exist.

The calculation of r, is illustrated in Table 9.4 on data gathered to

TABLE 9.4 ILLUSTRATION OF THE CALCULATION OF THE POINT-BISERIAL CORRELATION COEFFICIENT

	Y Sex	X Usiala	
Person	(1, male; 0, female)	fleight (in inches)	Calculating r,
1	1	59	
В С	0	67	
c	1	63	
D	1	65	
Ľ F	0	55	1. Equation (9.5);
F	1	72	l
G	0	62	64.25 61.14 / 8(7) 41
Ħ	0	60	$\frac{64.25 - 61.14}{3.91} \sqrt{\frac{8(7)}{15(14)}} = .41.$
1	1	64	•
3	1	66	2. Equation (9.6):
K	1	63	
L	0	61	$\frac{64.25 - 62.80}{3.91} \sqrt{\frac{8(15)}{7(14)}} = .41.$
41	1	62	3.91 7(14)
N	0	63	1
o	0	60	3. Equation (9.7):
A1 - 8	X - 64.25		62.80 - 61.14 77(15)
n 7	X - 61 14		$\frac{62.80 - 61.14}{3.91} \sqrt{\frac{7(15)}{8(14)}} = .41.$
n - 15	X ~ 62 80		
	s, e- 3 91		1
			1

relate set (the nominal-dichotomous variable) and height for 15 adolescents eight boys and seven girls. The boys are taller than the girls, on the average (64.25 versus 61.14 inches), but the relationship found between sex and height is only moderate (.41). Just one girl is taller than the average boy, but six of the seven pirks are taller than the shortest box.

#### Casa E

Both variables are dichotomous with underlying normal distributions: the tetrachoric correlation coefficient, r... In some instances, we think we know a great deal about the variable being measured, even though we can make only very crude measurements of it. For example, a test item is written to measure syllogistic reasoning power. The writer of the item believes that the ability to draw correct conclusions in a variety of syllogisms is a normally distributed trait, but the single test item will allow him to identify only a group of those who answer correctly (all of whom will be given a score of 1) and a group who do not (all of whom will be given a 0). As a second example. suppose that the heights of 1000 boys are approximately normally distributed. The researcher may choose to give a 1 to those taller than 5'2" and a 0 to those under that height as in Fig. 9.4. Surely he is discarding information. but he may gain a great amount of computational ease with tolerable loss of information, specially if his n is large. (This device was frequently used in psychometrics before the wide availability of mechanical computing equipment. Faced with the problem of computing several hundred correlation coefficients, it proved to be expedient to employ the methods discussed below as a short-cut approximation.)

Two variables X and Y are measured dichotomously on a group of persons, although it is believed that more costly and extensive operations could produce nearly normal distributions of measurements. Only 0 and 1 data are available, but the researcher's interest is in the correlation of X and Y he would have obtained if he had gathered the normally distributed measures. (The phi coefficient will usually underestimate this relationship.) If he feels that his understanding of the underlying variables (i.e., that they produce normally distributed measurements) entitles him to more information, informat




FIG. 9.4 Transforming normally distributed scores into dichotomous scores.

he will choose to calculate the tetrachoric coefficient of correlation, rest. between X and Y.

The observable data are dichotomous scores (0 or 1) for each person on X and Y. At this first stage, the data are in the same form as when a phi coefficient is calculated. All of the explicit information in the data is preserved when they are placed in a  $2 \times 2$  contingency table like that of Fig. 9.2.

The most exact formula for reet uses the frequencies a, b, c, and d in the  $2 \times 2$  contingency table to obtain an approximation to the value of  $r_{xy}$  that could be calculated if more sophisticated measurement of X and Y were possible. This formula was derived by Karl Pearson in 1900. Unfortunately, it is very complex, so we must settle for a more convenient though less exact approximation.


If you have some facility with trigonometry, you will have no trouble understanding the following formula for approximating rtet:

$$r_{trt} \doteq \operatorname{cosine} \frac{180^{\circ}}{1 + \sqrt{bc/ad}}.$$
 (9.8)

After calculating the term to the right of "cosine" in the above equation, you can refer to a table of trigonometric functions to find the cosine of the angle measured in degrees. Undoubtedly, you will find it more convenient to refer a function of the value of (ad)/(bc) to Table H in Appendix A, in which all of the operations in Eq. (9.8) have been carried out. It is entered with (bc)/(ad), provided that this ratio is greater than 1. If (bc)/(ad) is less than 1, the table is entered with (ad)|(bc), and the value of ren is negative.

Two examples will illustrate the use of Table H in finding rece

Two test items were administered to 100 persons. Each person was given a score of 1 if he answered an item correctly and a 0 if he answered it incorrectly or omitted it. Figure 9.5 contains a tabulation of the numbers of persons scoring each combination of correct and incorrect on the two items. The number of persons answering both items incorrectly was 64; five persons answered item 2 correctly and item 1 incorrectly. The value of (bc)/(ad) for the table (Fig. 9.5) is (64)(25)/(5)(6) = 53.33. Since this ratio is greater than I, it is referred directly to Table H in the Appendix. From Table H, we find that the approximate value of  $r_{tot}$  for a value of (bc)/(ad) equal to 53.33 is .93. This is an approximation to the value of the product-moment correlation coefficient between the normally distributed variables for which



Frequencies of pairs of

test items 1 and 2 provided only dichotomous measures. As a second example, suppose that for a certain  $2 \times 2$  contingency table, a = 31, b = 10, c = 5, and d = 14. If you draw this contingency table you will see a negative relationship between the two variables. Now (bc)/(ad) equals .115. Consequently, Table H is entered with (ad)/(bc), which equals 8.68, and the value of  $r_{te}$  is negative. From Table H we see that the value of  $r_{te}$  is -.70.

The reader must be warned that the use of Eq. (9.8) and, what is equivalent, of Table H can result in substantial errors (see Bouvier et al., 1954). Neither method should be used if (a + b) | n or (b + d) | n (i.e.,  $p_v$  or  $p_s$ ) departs greatly from .50. If both ratios equal .50, Table H will not be in error by more than .04 for values of  $r_{tet}$  between -.90 and +.90. When the error by more than .05, the tabled values are over-estimates of the true value of  $r_{tet}$ , when it is positive. If either (a + b) | n or (b + d) | n is greater than .70 or less than .30, the methods in this text for finding  $r_{tet}$  should not be used. Tables prepared by Jenkins (1955) should be used in such cases. The illustration in Fig. 9.5 of the use of Table H represents about the most text enterpresents about the most used.

An interesting development in the theory of correlation was the derivation of a "polychoric" correlation coefficient. Instead of transforming two normal distributions into dichotomies (i.e., just two scoring categories), which is the problem in the polychotomies having more than two categories. The problem is then one of estimating results between the normally distributed variables given only the contingency table, with several rows and distributed variables given only the contingency table, with several rows and distributed variables given only the contingency table, with several rows and is in the work of Lancaster and Hamdan (1964); and though it fills a theoretical gap, the computational labor involved requires the use of electronic computers.

The limits on the value of  $r_{tot}$  are -1 to +1. One particularly advantageous feature of this coefficient is that its maximum and minimum values are +1 and -1 regardless of how far (a + b)/n or (b + d)/n depart from equality. This property alone makes  $r_{tot}$  superior to the phi coefficient as a measure of relationship when normal distributions underlying the dichotomies can be assumed. (See Carroll, 1961.)

# Case F

93

X is dichotomous, with an underlying normal distribution, and Y is ordinal. We know of no coefficient appropriate for describing the relationship between wariables measured in this unorthodox combination of ways. If such measurements do arise, we suggest that you forego the assumption of a normal distribution underlying the dichotomy on X and proceed with the calculation of Cureton's rank-biserial correlation coefficient, which is discussed under Case H, below.

#### Case G

One variable yields dichotomous measures with an underlying normal distribution, the other yields Interval or ratio measures; the biserial correlation coefficient. Suppose that variable Y is measured dichotomously even though one might think that more extensive or refined techniques could produce a nearly normal distribution of scores on Y. We say that the measurement of Y has produced a dichotomy with an underlying normal distribution. Thus Y is measured in precisely the same way as either of the variables involved in the calculation of a tetrachoric correlation coefficient. However, in this instance the data produced by measurement of X can be thought of as scores on an interval or ratio scale that are approximately normally distributed. For example, the X scores might be scores on the Scholastic Aptitude Test, and the Y scores might be 0's and 1's on a test item measuring cognitive flexibility. With more elaborate tests of cognitive flexibility, it might be possible to produce a wide range of cognitive flexibility scores (Y scores) that are nearly normally distributed. These two arrays of normally distributed scores on X and Y could be correlated and rev found. What do the X scores and the dichotomous scores on Y tell about the value of  $r_{xy}$ ? The answer to this question lies in the biserial correlation coefficient  $r_{bas}$ . (In  $r_{bis}$  we meet another product of the genius of Karl Pearson.)

The biserial correlation coefficient is an estimate of the product-moment correlation between X and the normally distributed scores on Y that are assumed to underlie the dichotomous (0 or 1) scores. This situation is similar to the conditions out of which  $r_{tot}$  arose, except that for  $r_{tot}$  both variables were dichotomous with underlying normal distributions.

The data gathered for the computation of  $r_{bis}$  consist of an X score, which can be any one of several different values, and a Y score, which is either 0 or 1, for each of n persons.

Suppose that a teacher wishes to relate the amount of time (X) students spend studying balancing chemical equations and their ability (Y) to balance such equations. Measurements of X are gathered by students' own reports of the homework time they gave to the study of the subject. Although Y could probably be measured by an extensive achievement test in such a way that a nearly normal distribution of scores would result, suppose that the teacher's time available for testing will allow the administration of only one item, a chemical equation to be balanced. Thus a normal distribution of ability is assumed to underlie the 0-incorrect and 1-correct scores on the test item. Data that might result from such a study appear in Table 9.5. They show that 11 of the 18 students balanced the equation correctly and that they scored higher than did the 7 students who balanced the equations incorrectly.

The rationale that underlies the derivation of real is taken from regression theory. The regression coefficient for predicting X from the normally

distributed Y is

$$b_1 = \frac{r_{xy}s_x}{s_x},$$

where  $s_v$  is the standard deviation of the hypothetically normally distributed Y, not of the dichotomous measures taken on Y. The slope of the least-squares regression line for predicting X from Y can be approximated by the slope of the line that passes through the mean X score for those scoring 0 on Y (which is denoted  $X_0$ ) and the mean X score for those scoring 1 on Y (which is denoted by  $X_0$ ). This latter line is drawn on the scatter diagram in Table 9.5. This line is a least-squares line in a sense, because the mean of a group is the point around which the sum of squared deviations is at a minimum.

The slope of this line is equal to  $(X_1 - X_2)$  divided by the distance between the mean scores on the normally distributed Y for those scoring 0 and 1 on the dichotomy. This latter distance is difficult to find in any elementary manner (see Walker and Lev, 1953, for a more advanced discussion). The desired distance turns out to involve the height u of the unit normal curve above the point on the abscissa (i.e., the Y axis here) above which  $100(n_1/n)$  percent of the area lies.  $(n_1$  is the number of persons scoring 1 on the dichotomy.) Combining these facts appropriately yields the following computational formula for  $r_{D_B}$ :

$$r_{bis} = \frac{\bar{X}_{.i} - \bar{X}_{.0}}{s} \cdot \frac{n_1 n_0}{u n_2 \sqrt{n^2 - n}}, \qquad (9.9)$$

where  $X_{.1}$  and  $X_{.0}$  are the mean X scores for those scoring 1 and 0 on Y, respectively;

s, is the standard deviation of the X scores;

 $n_1$  and  $n_0$  are the numbers of 1's and 0's on Y, respectively  $(n_1 + n_0 = n)$ ; and

u is the ordinate (i.e., the height) of the unit normal distribution at the point above which lies  $100(n_1/n)$  percent of the area under the curve (see Table B in Appendix A).

The calculation of  $r_{tit}$  from Eq. (9.9) will be illustrated with the data in Table 9.5.

Since  $n_1/n = 11/18 = .61$ , the ordinate u of the unit normal curve for the point above which lies 61% of the area must be found. In Table B in Appendix A this height is found to be .3836. The summary statistics from Table 9.5 are substituted into Eq. (9.9);

$$r_{bis} = \frac{12.36 - 10.00}{2.55} \cdot \frac{11 \cdot 7}{(.3836)18\sqrt{18^2 - 18}} = .60.$$

variable is changed into a dichotomous variable by combining two adjacent categories).\*

#### Case H

172

Both variables yield ordinal measures. Spearman Rank-Correlation Coefficient. Raw data may be converted to ranks, or ranks may be gathered as the original data. "Rank in graduating class" is an example of the conversion of ordered scores to ranks: grade-point averages are computed for each of 500 students, say a rank of 1 is assigned the highest GPA, 2 to the next highest, ..., 500 to the lowest. "Judges' rankings of excellence of a recitation" is an example of gathering ranks as original data: 10 students recite a passage, and a judge assigns the rank 1 to the best recitation, 2 to the second best recitation, ..., and 10 to the worst recitation. Data are often gathered in this form when more refined measurements are not convenient, needed, or possible. Regardless of how the ranks 1, 2, ..., n-1, n are generated, two sets of ranks for the same n persons can be correlated in the same way.

Suppose that 12 students are ranked by a single judge on the basis of both overt hostility toward their teacher (X) and overt hostility toward other students (Y). A rank of 1 is given to the student showing the greatest hostility toward the teacher; 12 indicates that student who showed the least hostility toward the teacher. The same ranking procedure is used for hostility toward other students.

Hypothetical data appear in Table 9.6. Student A was ranked second on X and sixth on Y. An obvious method of describing the relationship between the two sets of ranks, X and Y, is to compute the product-moment correlation coefficient between the n paired ranks, i.e., simply substitute the data of Table 9.6 into the formula for r... This rather obvious technique was employed first by Francis Galton. Charles Spearman, the British psychologist, made more use of it. He has been rewarded by having the coefficient named in his honor, even though Karl Pearson apparently felt that the honor was due Galton (see Walker, 1929, p. 103). The productmoment correlation coefficient computed on two sets of the n consecutive, untied ranks 1, ..., n is known as the Spearman rank-correlation coefficient. We shall symbolize this coefficient by r.

Before you begin to calculate  $r_s$  using any of the usual formulas for  $r_{ss}$ . take note. The constraints that exist on X, Y,  $s_s^2$ , and  $s_s^2$  because X and Ytake on only the values  $1, 2, \dots, n$  lead to enormous simplifications in the

<sup>\*</sup> To avoid spuriously high  $r_{\rm tot}$ 's, however, one should be careful about such coalescing when done after the data (i.e., the X's and Y's) are collected. The particular combining of categories that works well in one sample might not do so in another, if the combining is done to maximize , in the first sample. Problems of crossvalidation and "shrinkage"

TABLE 9.6

THE TWO SETS OF RANKS ASSIGNED BY A JUDGE TO TWELVE STUDENTS ON HOSTILITY TOWARD THEIR TEACHER (X) AND HOSTILITY TOWARD OTHER STUDENTS (Y)

HOSTILITY TOWARD OTHER STU Hostility toward				Computations		
- Student	Teacher (X)	Students (Y	) X <sup>2</sup>	Y²	(X-Y)	
			4	36	16	
Α	2	•	64	25	9	
B	8	10	144	100	4	
C	12	10	144	49	16	
D	3		í	9	4	
E	1	3	36	16	4	
F	6	4		81	4	
G	7	9	49	64	4	
H	10	8	100	1	9	
ï	4	1	16	101	á	
<b>,</b>	9	11	81	121	7	
,	11	12	121	144		
·		2	25	_4		
L.	. <u>.</u> .	S v - 78	$\sum X^2 = \overline{650}$	$\sum Y^2 = 650$	$\sum (X-Y)^2 = \overline{84}$	

formula for  $r_{xx}$ . Notice in Table 9.6 that  $\sum X = \sum Y$  and  $\sum X^2 = \sum Y^2$ . You should realize immediately that these equalities must always hold, as long as no tied ranks are allowed. It can be shown that

$$\sum X = 1 + 2 + \ldots + n = \frac{n(n+1)}{2}$$

In Table 9.6, we have

$$\frac{n(n+1)}{2} = \frac{12(13)}{2} = 78.$$

The famous German mathematician and astronomer Karl Gauss is reputed to have demonstrated this fact as a child for his astonished tutor in the following manner:

If

$$\sum_{1}^{n} X = 1 + 2 + \ldots + (n-1) + n,$$

then of course

$$\sum_{1}^{n} X = n + (n-1) + \ldots + 2 + 1$$

also (i.e., the same series in reverse order). Add the two equations together vertically and obtain

$$2\sum_{i=1}^{n}X=(n+1)+(n+1)+\ldots+(n+1)=n(n+1).$$

because there are n of the (n + 1)'s. Divide both sides of the equation by 2; you obtain

$$\sum_{1}^{n}X=\frac{n(n+1)}{2},$$

the sum of the n consecutive, untied ranks  $1, 2, \ldots, n$ . Divide both sides of the above equation by n and you have

$$\bar{X}_{\cdot} = \frac{n+1}{2} \,, \tag{9.12}$$

the mean of the n consecutive, untied ranks  $1, 2, \ldots, n$ .

Using similar, but slightly more complicated techniques, it can be shown (see Siegel, 1956, or Edwards, 1964) that if X takes on the n consecutive and untied ranks 1, 2, ..., n, the variance of the X's is

$$s_x^2 = \frac{n^2 - 1}{12} \,. \tag{9.13}$$

Taking Eqs. (9.12) and (9.13) into consideration and performing other minor simplifications, we obtain the following formula for r, (see Pearson, 1907; Siegel, 1956, pp. 203-4);

$$r_{i} = 1 - \frac{6\sum_{i=1}^{n}(X_{i} - Y_{i})^{2}}{n(n^{2} - 1)}.$$
 (9.14)

 $X_i - Y_i$  is the difference between the ith person's rank on X and his rank on Y. The 12 values of  $(X_1 - Y_1)^2$  for the data in Table 9.6 appear in the last column of the table. Their sum is 84.

The value of r, for the two sets of ranks in Table 9.6 is found as follows:

$$r_{\bullet} = 1 - \frac{6(84)}{12(144 - 1)} = 1 - \frac{42}{143} = .71.$$

This value of r, indicates a fairly strong direct relationship between the students' overt hostility toward the teacher and toward other students as

When calculations are being performed on a desk calculator, the following formula, which is equivalent to Eq. (9.14), will be more convenient:

$$r_s = \frac{3}{n-1} \left[ \frac{4 \sum_{i=1}^{n} X_i Y_i}{n(n+1)} - (n+1) \right].$$
 (9.15)

 $X_iY_i$  is the product on person I's ranks on X and Y. Equation (9.15), which gives exactly the same value as Eq. (9.14), does not require the calculation of  $X_i = Y_i$ , which may need to be written down when calculations are

made on a desk calculator. Each journey of a datum from machine to paper

and back to machine is an opportunity to err. Often, though, the deviations can be squared and summed mentally. without writing down anything except the sum of the squared deviations [needed in Eq. (9.14)]. Stanley (1964, pp. 375-78) offers a table from which

 $r_i$  can be secured readily when one knows  $\sum_{i=1}^{n} (X_i - Y_i)^2$  and  $n_i$  for any value of n from 2 through 10. For further discussion of the use of  $r_s$  in practical situations, see Stanley (1964, pp. 98-100 and 379-81).

# The Problem of Tied Ranks

Ties in measurements often occur. When they do, there is a special rule for assigning ranks. For example, if the 12th and 13th highest ranking students in a graduating class of 245 each has a grade-point average of 4.76. it is customary to assign both students the average of the two ranks: (12 + 13)/2 = 12.5. Or a judge may find it impossible to discriminate between the quality of the handwriting of the top three students, so he gives all three students the average of the top three ranks, 2 = (1 + 2 + 3)/3. (As a generous gesture, some judges would give all three a rank of 1, but that fails to preserve the sum of the 3 untied ranks 1, 2, 3,)

When tied ranks occur, neither Eq. (9.14) nor Eq. (9.15) is equivalent to the  $r_m$  between the ranks. Even though the mean of the numbers  $1, 2, \ldots, n$ does not change when ties in the ranks occur [it is still (n + 1)/2], the variance of the ranks is less than (n2 - 1)/12. Consequently, the variance simplifications in the formula for ray that led to Eqs. (9.14) and (9.15) cannot be made

There are three possible ways to proceed in the calculation of r, when tied ranks occur: (a) use the computational formula for ray on the data-this will always give r, whether any ranks are tied or not; (b) use a formula (see Kendall, 1955) that incorporates corrections of s2 and s2 for the ties in the ranks; (c) compute an approximation to r, via Eq. (9.14) or (9.15), if there are few ties. With the current availability of fully automatic desk calculators and electronic computers we suggest method (a)-that you compute the rev between the ranks, having assigned ranks to tied measurements by the averaging method described above.

# Interpretation and Use of r,

No special interpretation can be given to r, over and above the statement that it equals the product-moment correlation coefficient calculated on ranks. The value of r, can never be less than -1 nor greater than +1. It equals +1 only if each person has exactly the same ranks on both X and Y.

The Spearman rank-correlation coefficient is especially useful when the original data are ranks, as when judges rank persons or things. It is sometimes regarded as a quick means of estimating  $r_{xy}$ . The original data are ranked,  $r_i$ , is computed, and a trigonometric function of  $r_i$  is supposed to convert  $r_i$  back into the value of  $r_{xy}$ . This procedure does not accomplish this except approximately so for very large n. Desk calculators and electronic computers have taken enough of the labor out of calculation so that there is no longer any reason not to calculate  $r_{xy}$  on unranked data.

The Spearman rank-correlation coefficient r, is very closely tied to the concept of a product-moment correlation coefficient. Another coefficient of correlation exists for data that are ranks on both X and Y; it involves a rather different concept of what "relationship" is and how it should be measured.

# Kendall's Tau, τ

All of the coefficients of correlation met in this book so far employ the product-moment rationale of Pearson in one form or another. Some coefficients are merely the Pearson product-moment formula applied to dichotomous or ordinal data, e.g.,  $\phi_t$ ,  $r_t$ , and  $r_{tb}$ ; others are attempts to approximate the Pearson  $r_t$ , e.g.,  $r_{tt}$ , and  $r_{tb}$ . The English statistician Maurice Kendall made one of the few attempts at conceptualizing the measurement of relationships between variables in a manner other than by use of the product-moment principle. His efforts have resulted in one of the few basically new approaches to statistical description in modern times. For a readable account of his work see Kendall (1955).

As was the case with the Speatman coefficients, the observations on both X and Y are the n consecutive and untied ranks  $1,2,\ldots,n$ . Kendall based his coefficient of correlation on the numbers of pairs of ranks that are ordered in the same direction on both X and Y. Thus, his measure, called the rankings on X and Y. The speak of X is the result of disagreement in the rankings on X and Y.

Suppose that the ranks in Table 9.7 were assigned to eight persons on X and Y. Suppose we select any pair of persons, e.g., A and B. A is ranked above B on X, but B is ranked above A on Y. This represents a departure from a direct relationship between X and Y; this inversion of the relative between X and B as we move from X to Y argues for an inverse relationship between X and B as as far as A and B are concerned. There are relationship between X and Y must be studied. For any pair of persons, and the contribution of each pair to the relationship between X and Y must be studied. For any pair of persons and B contribute on A and B is the same. For example, persons A and B contribute on A is the same. For example, on X (1 vs. 4) and on Y (3 vs. 5). For any pair of persons, an inversion is counted if their order on X and Y is different. Persons A and B contribute

TABLE 9.7 ILLUSTRATION OF THE CALCULATION OF KENDALL'S TAU, 7

Person	X	Y	Agreements	Inversions	
	1	3	5	2	_
Ĉ	. ,	1	6	0	n = 8
,	7	2	5	0	P - Q = 21
D	4	- 5	3	1	
н	7	7	1	2	$\tau = \frac{1}{n(n-1)/2}$
Ł	2		Ō	2	14
F		4	1	0	$=\frac{14}{28}=.50$
D	7	*	ń	0	28
G	8	6	$P = \overline{21}$	$o = \frac{2}{7}$	

an inversion; note that F and D also contribute one inversion; F has a higher rank than D on X but a lower rank than D on Y.

With no tied ranks, all n(n-1)/2 pairs of persons will contribute either an agreement or an inversion. When the rankings on both X and Y are identical, there will be n(n-1)/2 agreements and no inversions. What will be the total numbers of agreements and inversions when Y is the reverse of  $X^2$ ? Zero and n(n-1)/2, respectively.

These facts can be pieced together in various ways to define a coefficient of the relationship between X and Y. Kendall chose the following definition for his coefficient  $\tau$ :

or his coefficient 
$$\tau$$
:
$$\tau = \frac{\text{(total number of "agreements")} - \text{(total number of "inversions")}}{n(n-1)/2}.$$
(9.16)

The process of counting the agreements and inversions in the ranks on X and Y for all n(n-1)/2 pairs of persons can be laborious. Fortunately, it can be simplified.

The first simplification occurs by algebraic manipulation of Eq. (9.16). The first simplification occurs by algebraic manipulation of Eq. (9.16) et us call the total number of inversions O. In this notation, Eq. (9.16) becomes

$$\tau = \frac{P - Q}{n(n-1)/2}.$$

It is customary to denote P-Q by the symbol  ${\cal S}$  so that au may be written

$$\tau = \frac{S}{n(n-1)/2} \,. \tag{9.17}$$

Since the sum of P and Q must be n(n-1)/2, we can write Eq. (9.17) as

$$\tau = \frac{P - Q}{n(n-1)/2} = \frac{[n(n-1)/2 - Q] - Q}{n(n-1)/2} = 1 - \frac{4Q}{n(n-1)}.$$
 (9.18)

# Kendall's & With Tied Ranks

When ties occur in the ranks on either X or Y, P and Q are still determined as indicated in Table 9.7. The only alteration in the formula for  $\tau$  occurs in the denominator. The correction of the denominator of  $\tau$  involves the quantities  $K_x$  and  $K_y$  (both of which are functions of the numbers of persons tied at the various ranks on Y and Y)

The following formula is applied when ties occur in the ranks on X and Y.

$$\tau = \frac{P - Q}{\sqrt{[n(n-1)/2] - K_* \sqrt{[n(n-1)/2] - K_*}}},$$
 (9.20)

where  $K_x = (\frac{1}{2}) \sum f_i(f_i - 1)$  (where  $f_i$  is the number of tied observations in each group of ties on X), and

 $K_{\nu} = (\frac{1}{2}) \sum f_i(f_i - 1)$  (where  $f_i$  is the number of tied observations in each group of ties on Y).

For an illustration of the application of Eq. (9.20), see Siegel (1956, pp. 218-19) or a more recent textbook on nonparametric statistics.

# Case 1

X is ordinal and Y is interval or ratio. No coefficient has been developed and studied for this particular case. If you find yourself with variables measured in these ways, it would be advisable to convert the Y scores to ranks and proceed with the calculation of either Spearman's or Kendall's rank-correlation coefficients.

# Return to Case C

Rank-biserial correlation. At this point it is appropriate to discuss the coefficient first mentioned under Case C, above. One notable coefficient for correlating a dichotomous variable X and an ordinal variable Y has been investigated by Cureton (1956) and Glass (1966b). This coefficient is closely related to Kendall's  $\tau$  and incorporates in its definition the concepts of agreements and inversions. We shall denote this coefficient by  $r_{rb}$ , the rank-biserial  $r_r$ .

Let X be a dichotomous variable and Y a variable comprising the n untied ranks  $1, 2, \dots, n$ . Curreno sought a coefficient descriptive of the relationship between X and Y such that (a) it would have attainable limits  $\pm 1$  under all circumstances, (b) it would be +1 when the n, highest ranks are all 1 on the dichotomy, and (c) it would be strictly nonparametric, i.e., defined wholly in terms of inversions and agreements without such concepts

as mean, variance, regression, etc. Suppose that the following data are gathered on X and  $\tilde{Y}$  for n = 10 persons:

Person	X	Y
Λ	0	1
В	1	10
C	0	2
D	1	9
E	0	5
F G	0	8
	1	4
H	1	7
I	0	3
J	0	6

To compute res the data are arranged in the following manner:

X = 1	on $Y$ for $X = 0$	Agreements	Inversions
10		6	
9		6	
	8	0	
7	0		2
,		5	
	6		
	5		
4		3	1
	3	,	
	2		
	1		
		$P=\overline{20}$	$Q = \tilde{4}$

There is an agreement at any given rank under column I for every smaller rank under column 0. There is an *incersion* at any given rank under column 0 for every smaller rank under column 1. Thus there are three agreements corresponding to the rank "4" under column I since there are three smaller tanks, 3, 2, and 1, under column 0. P is the sum of all agreements in the data, and Q is the sum of all inversions. Cureton defined r., as follows:

$$r_{n} = \frac{P - Q}{P_{max}}. (9.21)$$

When no ties exist,  $P_{max} = n_0 n_1$ , where  $n_0$  is the number of persons at 0 on

the dichotomy and  $n_1$  is the number of persons at 1. Hence

$$r_{rb} = \frac{P - Q}{r_{rb}}. (9.22)$$

For the above data.

$$r_{rb} = \frac{20-4}{(4)(6)} = \frac{16}{74} = \frac{2}{3} = .67.$$

Glass (1966b) has shown that  $r_{rb}$  is algebraically equivalent to a coefficient analogous to  $r_{bt}$  for ordinal variables. The practical importance of this equivalence is that it provides an easy means of computing  $r_{rb}$  without counting agreements and inversions. The following formula can be shown to yield the same value as that given by Eq. (9.22):

$$r_{rb} = \frac{2}{r} (\bar{Y}_1 - \bar{Y}_0),$$
 (9.23)

where  $\overline{Y}_1$  is the average rank of those scoring 1 on X and  $\overline{Y}_0$  is the average rank of those scoring 0 on X.

The data on which the calculation of  $r_{rb}$  by Eq. (9.22) was illustrated will be used to illustrate use of Eq. (9.23).

Ranks of $X = 1$		Calc	ulations
10	8	$n_1 = 4$ $n_0 = 6$	
9	6		
7	5	$\overline{Y}_1 = \frac{3.6}{4} = 7.500$	$\overline{Y}_0 = \frac{2.5}{6} = 4.167$
4	3		
	2	$r_{rb} = \frac{2}{10}(7.500 - 4)$	$.167) = \frac{3.333}{5} = .67$
	1		

When no ties occur in Y, Eq. (9.23) can always be used in place of Eq. (9.22). For a discussion of the case when there are ties in Y see Cureton (1968).

Whitfield (1947) derived a coefficient for correlating one dichotomous and one ordinal variable. His rationale was to consider the dichotomous variable to be a ranking variable ted at two ranks. He then applied Kendall's  $\tau$  formula for tied ranks. The resulting coefficient has the same numerator as  $r_0$ , but a different denominator. The rank-biserial coefficient is to be preferred as a measure of correlation because Whitfield's coefficient may not attain +1 when some perfect relationships between X and Y exist. For example, when n=5,  $n_1=2$ , and  $n_0=3$  and the fourth- and fifth-ranking persons on Y have the two scores of 1 on X,  $r_0=1$ , but Whitfield's coefficient equals only .77.

# 9.4 PART CORRELATION AND PARTIAL CORRELATION

Concepts from simple linear regression and correlation are combined in part correlation and partial correlation. We shall begin this section with the development of part correlation, since partial correlation is its generalization—statistically, at least.

A researcher wishes to determine the correlation between a measure of intelligence X and learning performance during an instructional unit in social studies. He chooses to measure X with the Kuhlmann-Anderson intelligence test; however, he faces some important decisions about how to measure learning performance. He can construct a respectable achievement test of the content being learned. But to give the test and score it for "number of correct answers" is not what this researcher means by the words "learning performance." A large correlation of X with the achievement-test score might result even though no learning at all took place during instruction, because variability on the achievement test could be due to intelligence and testwiseness and not be at all due to differential learning during instruction. Administering the achievement test both before and after instruction and subtracting each student's initial score from his post-instruction score produces a measure that is far closer to the researcher's notion of a measure of "learning performance." One slight difficulty remains. Such "posttest minus pretest" measures of learning would have a predictable negative correlation with intelligence due, in part, to the manner in which measurement errors are combined by subtracting one fallible measure from another-In fact, it is almost certain that these "difference scores" will have a negative correlation with the pretest scores upon which they are based. This is considered a defect of such "difference scores" when we have reason to believe that amount of learning should not necessarily correlate negatively with pretest status. An alternative method is to measure learning or change by fitting a straight regression line to the pretest and posttest achievementtest data and taking the deviation from the regression line (errors of estimate) measured along the posttest axis. This deviation, called a residual change score, is illustrated in Fig. 9.6 where Y and Z denote the posttest and pretest, respectively. (The Z here is not the standard score z mentioned earlier.)

The residual gain score is denoted by  $\epsilon_s$ , because it is precisely the same as the error made in predicting Y from Z by the least-squares regression line. We know from previous experience with the regression model that the correlation of  $\epsilon_s$ , with Z is always zero. As a measure of learning, then,  $\epsilon_s$ , has the property that the measure of how much has been learned is






FIG. 9.6 Illustration of the definition of the residual gain score.

unrelated (r = 0) with initial performance. The researcher may consider this property desirable.\*

The correlation of X with  $e_{yx}$  is called a part correlation. In a sense, it is the correlation of X with Y after the portion of Y that can be predicted linearly from Z has been removed from Y. However, such an ambiguous verbal definition will not substitute for the unequivocal definition embodied in the symbol  $r_{xx}$ :

Of course, the part-correlation coefficient  $r_{xx_{y,z}}$  could be found by actually calculating the values of  $e_x$ , from the regression line of Y on Z and correlating these values with X. However, we shall now see how this computational labor can be bypassed.

By definition, rge is given by

$$r_{xx_{y,z}} = \frac{s_{xx_{y,z}}}{s_{x,x_{y,z}}}$$
, (9.24)

i.e., by the ratio of the covariance of X with  $e_{vz}$  to the product of the two standard deviations.

From Sec. 8.3, we know  $s_{e_{y,a}} = s_y \sqrt{1 - r_{ye}^2}$ . It remains to evaluate the numerator of Eq. (9.24)

$$s_{x_{t_{n-1}}} = s_{x_t y - b_n - b_1 x_1} = s_{xy} - 0 - b_1 s_{xy}.$$
 (9.25)

The properties of residual gain scores have been extensively studied; see Bereiter (1963), Lord (1963), and Tucker, Damarin, and Messick (1966).

(The zero occurs because the covariance of X with the constant  $b_0$  is 0.)

The regression slope  $b_1$  in Eq. (9.25) is for predicting Y from Z; thus

The regulation along of a facts in Eq. (9.25) produces
$$r_{xx_x} = \frac{s_{xx} - r_{x_x}(s_x)s_{xx}}{s_{xx_x} - s_{xx_x}(s_x)s_{xx}}.$$
(9.26)

Dividing the numerator and denominator of Eq. (9.26) by szs, yields

$$r_{ze_{y,z}} = \frac{(s_{zy}|s_z s_y) - r_{yz}(s_{zz}|s_z s_z)}{\sqrt{1 - r_{zz}^2}} = \frac{r_{zy} - r_{zz}r_{yz}}{\sqrt{1 - r_{yz}^2}}.$$
(9.27)

Thus we see that  $r_{se_{pe}}$  can be calculated directly from  $r_{xy}$ ,  $r_{xx}$ , and  $r_{yy}$ . For example, let X be the measure of intelligence from the Kuhlmann-Anderson test and Z and Y the measures of present and positest achievement, respectively. Suppose that  $r_{xy} = .70$ ,  $r_{xz} = .50$ , and  $r_{yz} = .80$ . The value of  $r_{xx_p}$  is

$$r_{xe_{x,x}} = \frac{.70 - (.50)(.80)}{\sqrt{1 - (.80)^2}} = \frac{.70 - .40}{.60} = .50,$$

Thus, although the correlation  $r_{xy}$  between X and Y is .70, when one eliminates the linear relationship of Y with Z, the residual relationship of X with  $e_y$ , is only .50.

By no means is part correlation limited to applications in which Z and Y are "pretests" and "posttests." One could, for example, apply the part-correlation technique with X a measure of reading speed, Y a measure of reading comprehension, and Z a measure of intelligence. In this instance,  $x_{ss_s}$ , measures the relationship between reading speed X and that part of reading comprehension  $e_s$ , that is unrelated to intelligence.

In one sense, partial correlation is a simple extension of part correlation. To find the partial correlation of X and Y with Z "held constant" or "partialed out," we merely calculate the two arrays of errors of estimate for predicting X from Z and Y from Z and correlate them. Symbolically, the partial correlation of X and Y with Z partialed out its given by

$$r_{xyz} = r_{e_1,x_2},$$
  
where  $e_{xz} = X - (b_0 + b_1 Z)$  and  $e_{yz} = Y - (b_0^* + b_1^* Z).$ 

In the numerical-subscript notation sometimes used this becomes

$$r_{11.3} = \frac{r_{11} - r_{12}r_{13}}{\sqrt{1 - r_{11}^2}}$$

Notice that there are two distinct regression lines involved in calculating the two sets of errors of estimate (or "residuals"): the line for predicting X from Z—which has constants  $b_0$  and  $b_1$ —and the line for predicting Y from Z—the constants for which have been labeled  $b_0^*$  and  $b_1^*$  to distinguish them from those for the X and Z regression line.

As in the case of part correlation, the actual calculation of the errors of estimate for all n persons is unnecessary in the calculation of  $r_{xy}$ . The partial correlation coefficient can be calculated directly from  $r_{xy}$ ,  $r_{xz}$ , and  $r_{yz}$ .

It can be shown, in a manner analogous to the derivation of Eq. (9.27) for the part-correlation coefficient, that  $r_{xyz}$  is given by the following formula:

$$r_{xy\cdot z} = \frac{r_{xy} - r_{xz}r_{yz}}{\sqrt{1 - r_{xz}^2}\sqrt{1 - r_{yz}^2}} .*$$
 (9.28)

Next we shall see what interpretation may be given to a calculated value of  $r_{avx}$ . One might hypothesize that there exists a positive correlation between reading performance X and visual perceptual ability Y (as evidenced by eye coordination, scanning speed, etc.). Suppose a sample of 30 children. ranging in age (Z) from 6 years to 15 years, yields a correlation of X with Y. r., of .64. The conclusion that some children read better than others because of greater perceptual abilities is tempting, but the cautious researcher will avoid drawing it. It is obvious that as children grow older they develop greater eye coordination and other perceptual abilities as a part of natural maturation. Moreover, the same children receive instruction in school which helps make them better readers year after year, up to a point. Could it not be that measures of both X and Y increase (improve) with age, in the one case as a consequence of physical maturation and in the other case as a function of mental maturation and increased exposure to instruction? This certainly could be the case. If the correlation of X and Y were zero at any one level of chronological age (instead of over the range from 6 years to 15 years), the  $r_{xy}$  of .64 for the sample of 30 children has far different implications. Indeed, even the cautious researcher would be tempted to conclude that the observed  $r_{xy}$  of .64 was due to a common relationship of both reading performance X and visual perceptual ability Y to chronological age Z, and was not due to any direct relationship between X and Y. How can we find what the value of  $r_{xy}$  would be for any single value of the chronological age variable Z? Under suitable assumptions, this desired correlation equals the partial correlation of X and Y with Z held constant:  $r_{xyz}$ .

r<sub>ev</sub>: is called a first-order partial-correlation coefficient, because the linear influence
of ne variable, Z, is partialed out.
 r<sub>ev</sub> is a zero-order correlation coefficient, because nothing
is partialed out.

Provided that Z has a linear relationship with both X and Y and that the strength of the linear relationship between X and Y is the same for persons at any level of Z, then  $r_{xx}$ , equals the value of  $r_{xy}$  we would obtain by correlating X and Y for a group of persons having identical scores on Z. For example, in the above illustration suppose that  $r_{xy} = .64$ ,  $r_{xx} = .80$ , and  $r_{yx} = .80$ . The value of  $r_{xx}$ , from Eq. (9.28) is

$$r_{xy\ z} = \frac{.64 - (.80)(.80)}{\sqrt{1 - .80^2}\sqrt{1 - .80^2}} = \frac{.64 - .64}{(.60)(.60)} = 0.$$

Thus we would estimate the value of  $r_{xy}$  for children of the same chronological age to be zero. If enough children of the same chronological age were available, we could calculate  $r_{xy}$  for them alone to check the above result. However, in our example there was not a sufficient number of children of the same age; there were 30 who ranged from 6 to 15 years of age. The partial-correlation coefficient serves the purpose of estimating  $r_{xy}$  for a single level of chronological age when there is an insufficient number of persons at any single chronological age to do the estimating by direct calculation.

More than one variable can be "partialed out," using the techniques of multiple regression to be discussed in the next section. For formulas and explanations, see Exekiel and Fox (1963). With partial correlation, especially, it is well to keep in mind that correlation does not necessarily mean causation. In the behavioral sciences causation is usually complex and indirect. One must be quite nimble intellectually to avoid assuming that he has isolated causal influences when other explanations of one's results are at least as plausible. For discussions of pitfulls see Lerner (1965).

# 9.5 MULTIPLE CORRELATION AND PREDICTION

The last correlational technique to be presented in this chapter is known as multiple correlation. As was true with the ordinary Pearson product-moment correlational technique of Chapter 7, multiple correlation has a second side known as multiple prediction. Stated broadly, the purpose of multiple prediction is the estimation of a variable Y, the dependent variable, from a linear combination of m independent variables  $X_1, X_2, \dots, X_m$ .

You will recall from Chapter 8 that when one variable X was used to estimate a second variable Y, the estimation criterion of "least squares" was equivalent to choosing values for  $b_0$  and  $b_1$  so that

$$\sum_{i=1}^n (Y_i - b_0 - b_i X_i)^2$$

was as small as possible. The equation  $b_0 + b_1 X_1$  provides the least-squares estimate  $Y_1$  of the ith person's score on the variable  $Y_2$ . This type of estimates

mation is sometimes termed univariate estimation or prediction because there is only one "predictor variable." A multivariate prediction of the Y variable given scores on m independent variables is

$$\hat{Y} = b_0 + b_1 X_1 + b_2 X_2 + ... + b_m X_m,$$
 (9.29)

Equation (9.29) provides a multiple-prediction or multiple-regression equation. Equation (9.29) is sometimes referred to as a linear regression equation since the b's appear only with exponents of 1 and never appear as squared terms, cubed terms, etc. Of course, Eq. (9.29) alone is of no value; the procedure by which "good" values for the b's are chosen must be specified. Again the least-squares criterion is invoked, and those values of  $b_{n_1}, b_{n_2}, \dots, b_{n_n}$  are chosen that minimize the quantity

$$\sum_{i=1}^{n} (Y_i - b_0 - b_1 X_{i1} - b_2 X_{i2} - \dots - b_m X_{im})^2$$
 (9.30)

for a given set of values of Y and the X's.

The values  $b_0, \ldots, b_m$  that minimize the above quantity provide us with  $b_0 + b_1 X_1 + \ldots + b_m X_m$  as a good estimate of Y:

$$\hat{Y}_{i} = b_{0} + b_{1}X_{i1} + \ldots + b_{m}X_{im}$$

The Pearson product-moment correlation between Y and  $\hat{Y}$  is a measure of how well the "best" linear weighting of the independent variables  $X_1, \ldots, X_m$  predicts or correlates with the single dependent variable Y. This special case of Pearson's r is called the multiple correlation coefficient and is denoted by  $R_{n,1,2,\ldots,m}$ . A second sense in which the values  $b_0,\ldots,b_m$  that minimize Eq. (9.30) are "best" is as follows: the maximum possible positive correlation between Y and any linear combination of  $X_1,\ldots,X_m$  is attained when the Y's are combined into

$$b_0 + b_1 X_1 + \ldots + b_m X_m$$

the b's being those values that minimize Eq. (9.30). Thus,

$$b_0 + b_1 X_1 + \ldots + b_m X_m$$

not only provides the least-squares estimate of Y but it also correlates higher with Y than could any other linear combination of the X variables. (As a consequence of how the weights for the X variables are derived, the multiple correlation coefficient will always be positive or zero.)

The theory and techniques of multiple prediction and correlation are involved and complex. A comprehensive treatment of these subjects would occupy many pages. Fortunately, there is no lack of excellent treatments of these topics in the pedagogical literature of applied statistics. You will find the theoretical and applied aspects of multiple regression developed fully in such texts as DuBois (1957), Rozeboom (1966), Draper and Smith (1966), and Acton (1959). For a briefer treatment see Williams (1968).

Provided that Z has a linear relationship with both X and Y and that the strength of the linear relationship between X and Y is the same for persons at any level of Z, then  $r_{xy}$ , equals the value of  $r_{xz}$  we would obtain by correlating X and Y for a group of persons having identical scores on Z. For example, in the above illustration suppose that  $r_{xy} = .64$ ,  $r_{xz} = .80$ , and  $r_{yz} = .80$ . The value of  $r_{xy}$ , from Eq. (9.28) is

$$r_{xy\,x} = \frac{.64 - (.80)(.80)}{\sqrt{1 - .80^2}\sqrt{1 - .80^2}} = \frac{.64 - .64}{(.60)(.60)} = 0.$$

Thus we would estimate the value of  $r_{xy}$  for children of the same chronological age to be zero. If enough children of the same chronological age were available, we could calculate  $r_{xy}$  for them alone to check the above result. However, in our example there was not a sufficient number of children of the same age; there were 30 who ranged from 6 to 15 years of age. The partial-correlation coefficient serves the purpose of estimating  $r_{xy}$  for a single level of chronological age when there is an insufficient number of persons at any single chronological age to do the estimating by direct calculation.

More than one variable can be "partialed out," using the techniques of multiple regression to be discussed in the next section. For formulas and explanations, see Ezekieland Fox (1963). With partial correlation, especially, it is well to keep in mind that correlation does not necessarily mean causation. In the behavioral sciences causation is usually complex and inferet. One must be quite nimble intellectually to avoid assuming that he has isolated causal influences when other explanations of one's results are at least as plausible. For discussions of pitfalls see Lerner (1965).

# 9.5 MULTIPLE CORRELATION AND PREDICTION

The last correlational technique to be presented in this chapter is known a multiple correlation. As was true with the ordinary Pearson product-moment correlational technique of Chapter 7, multiple correlation has a second side known as multiple prediction. Stated broadly, the purpose of multiple prediction is the estimation of a variable Y, the dependent variable, from a linear combination of m independent variables  $X_1, X_2, \dots, X_m$ .

You will recall from Chapter 8 that when one variable X was used to estimate a second variable Y, the estimation criterion of "least squares" was quivalent to choosing values for  $b_n$  and b, so that

$$\sum_{i=1}^{n} (Y_i - b_0 - b_1 X_i)^2$$

was as small as possible. The equation  $b_0 + b_1 X_i$  provides the least-squares estimate  $Y_i$  of the ith person's score on the variable  $Y_i$ . This type of estimates  $Y_i$  in the  $Y_i$  is the  $Y_i$  in the Y

mation is sometimes termed *univariate* estimation or prediction because there is only one "predictor variable." A *multivariate* prediction of the Y variable given scores on m independent variables is

$$\hat{Y} = b_0 + b_1 X_1 + b_2 X_2 + \ldots + b_m X_m. \tag{9.29}$$

Equation (9.29) provides a multiple-prediction or multiple-regression equation. Equation (9.29) is sometimes referred to as a linear regression equation since the b's appear only with exponents of I and never appear as squared terms, cubed terms, etc. Of course, Eq. (9.29) alone is of no value; the procedure by which "good" values for the b's are chosen must be specified. Again the least-squares criterion is invoked, and those values of b. b. . . . b. are chosen that minimize the quantity

$$\sum_{i=1}^{n} (Y_i - b_0 - b_1 X_{i1} - b_2 X_{i2} - \ldots - b_m X_{im})^2$$
 (9.30)

for a given set of values of Y and the X's.

The values  $b_0, \ldots, b_m$  that minimize the above quantity provide us with  $b_0 + b_1 X_1 + \ldots + b_m X_m$  as a good estimate of Y:

$$\hat{Y}_{i} = b_{0} + b_{1}X_{i1} + \ldots + b_{m}X_{im}.$$

The Pearson product-moment correlation between Y and  $\hat{Y}$  is a measure of how well the "best" linear weighting of the independent variables  $X_1, \ldots, X_m$  predicts or correlates with the single dependent variable Y. This special case of Pearson's r is called the multiple correlation coefficient and is denoted by  $R_{g,1,k,\ldots,m}$ . A second sense in which the values  $b_0,\ldots,b_m$  that minimize Eq. (9.30) are "best" is as follows: the maximum possible positive correlation between Y and any linear combination of  $X_1,\ldots,X_m$  is attained when the Y's are combined into

$$b_0 + b_1 X_1 + \ldots + b_m X_m$$

the b's being those values that minimize Eq. (9.30). Thus,

$$b_0 + b_1 X_1 + \dots + b_m X_m$$

not only provides the least-squares estimate of Y but it also correlates higher with Y than could any other linear combination of the X variables. (As a consequence of how the weights for the X variables are derived, the multiple correlation coefficient will always be positive or zero.)

The theory and techniques of multiple prediction and correlation are involved and complex. A comprehensive treatment of these subjects would occupy many pages. Fortunately, there is no lack of excellent treatments of these topics in the pedagogical literature of applied statistics. You will find the theoretical and applied aspects of multiple regression developed fully in such texts as DuBois (1957), Rozeboom (1966), Draper and Smith (1966), and Acton (1959). For a briefer treatment see Williams (1968).

In the remainder of this section we shall illustrate multiple prediction and correlation for the case in which two variables  $X_1$  and  $X_2$  are used to predict the criterion variable Y. Suppose that Y is grade-point average for college freshmen at the end of two semesters. One wishes to predict Y from  $X_1$ , high-school grade-point average, and  $X_2$ , verbal ability as measured by the Scholastic Aptitude Test. Many details of the following discussion will be simplified if it is assumed that Y,  $X_1$ , and  $X_2$  are transformed to standard scores with mean zero and variance 1. Thus Y,  $X_1$ , and  $X_2$  become  $z_2$ ,  $z_4$ , and  $z_2$ . We seek the values of  $b_0$ ,  $b_1$ , and  $b_2$  that minimize the quantity

$$\sum_{1}^{n} (z_{v} - b_{0} - b_{1}z_{1} - b_{2}z_{2})^{2}$$

for a group of n students for whom freshman GPA, Y, high-school GPA,  $X_2$ , and SAT Verbal scores,  $X_2$ , are available.

Without substantiating the claim here, we state that the least-squares estimates  $b_0$ ,  $b_1$ , and  $b_2$  are as follows:

$$b_0 = 0$$
,  $b_1 = \frac{r_{v1} - r_{v2}r_{12}}{1 - r_{12}^2}$ ,  $b_2 = \frac{r_{v2} - r_{v1}r_{12}}{1 - r_{12}^2}$ . (9.31)

The fact that  $b_0$  is zero is a consequence of having transformed Y,  $X_1$ , and  $X_2$  to standard scores;  $b_0$  would not be 0 if these variables had not been standardized. The quantities  $r_{01}$ ,  $r_{22}$ , and  $r_{12}$  are simply the correlation coefficients between Y and  $X_1$ , Y and  $X_2$ , and  $X_1$  and  $X_2$  calculated on the data from the n students.

The best (in the least-squares sense) estimate of the *i*th person's standard score on Y given his standard scores on  $X_1$  and  $X_2$  is

$$\hat{z}_* = b_1 z_1 + b_2 z_2, \tag{9.32}$$

where  $b_1$  and  $b_2$  are as defined in Eq. (9.31). Transforming Eq. (9.32) back into the scale of the original variables gives the multiple-regression equation for estimating Y from  $X_1$  and  $X_2$ :

$$\hat{Y}_{t} = \left(b_{1} \frac{s_{2}}{s_{1}}\right) X_{t1} + \left(b_{2} \frac{s_{2}}{s_{2}}\right) X_{t2} + \left(\bar{Y}_{t} - b_{1} \frac{s_{2}}{s_{1}} \bar{X}_{,1} - b_{2} \frac{s_{2}}{s_{2}} \bar{X}_{,2}\right). \quad (9.33)$$

The correlation coefficient between the n actual freshman grade-point averages ( $\Upsilon$ ) and the n predicted freshman grade-point averages ( $\Upsilon$ ) obtained from Eq. (9.3) is the multiple correlation coefficient  $R_{n,1}$  between freshman grade-point average and the best linear composite of high-school grade-point average and SAT Verbal scores. One could compute  $R_{n,1}$  directly from the n pairs of values of  $Y_n$  and  $Y_n$ . However, once  $Y_{n+1}, Y_{n+1}$  and  $Y_n$ , are known, the multiple correlation coefficient is given conveniently by the following equation:

$$R_{r,1,2} = \sqrt{b_1 r_{s1} + b_2 r_{s2}}. (9.34)$$

The values of  $b_1$  and  $b_2$  in the above equation are, of course, obtained from the formulas in Eq. (9.31). Note that the positive square root is taken so that R will never be negative. It is not as simple to see that the term under the radical will never be negative, but such is the case.

Data from a study conducted by Dizney and Gromen (1967), shown in Table 9.8, will be used to illustrate multiple prediction and correlation for

TABLE 9.8 SUMMARY DATA FROM A MULTIPLE-PREDICTION STUDY (DIZNEY AND GROMEN, 1967) \*

	Intercorrelations				Standard
	$X_1$	$X_{\mathbf{a}}$	Y	Means	deviations
X <sub>1</sub> (MLAReading)	1.00	.58	.33	25.55	10 20
X <sub>2</sub> (MLAWriting)		1.00	.45	63.22	11.91
Y (German grade)			1.00	2,61	0.50

the case of two independent variables. Dizney and Gromen studied the relationship of reading proficiency  $X_1$  and writing proficiency  $X_2$  (both as measured by the Modern Language Association Foreign Language Proficiency Tests) to course grades in the second quarter of a college German class. A total of n=111 students participated.

The values of  $b_1$  and  $b_2$  for predicting standard scores on Y from standard scores on X, and  $X_0$  are found from Eq. (9.31) as follows:

$$b_1 = \frac{r_{y1} - r_{y2}r_{12}}{1 - r_{12}^2} = \frac{.33 - .45(.58)}{1 - (.58)^2} = .104$$

$$b_2 = \frac{r_{y2} - r_{y1}r_{12}}{1 - r_{12}^2} = \frac{.45 - .33(.58)}{1 - (.58)^2} = .390.$$

Hence, the best estimate of standard scores on Y given  $z_1$  and  $z_2$  is

$$\dot{z}_{..} = .104z_1 + .390z_2$$

From Eq. (9.33) and the data in Table 9.8 we can construct the multipleprediction equation for raw scores.

$$b_1 \frac{s_2}{s_1} = .104 \left( \frac{0.50}{10.20} \right) = .005 \qquad b_2 \frac{s_2}{s_2} = .390 \left( \frac{0.50}{11.91} \right) = .016,$$

$$\dot{Y} = .005X_1 + .016X_2 + [2.61 - (.005)25.55 - (.016)63.22]$$

$$= .005X_1 + .016X_2 + 1.47.$$

<sup>\*</sup> Dirney and Gromen report their statistics to too few significant figures for the desirable level of accuracy in the computations here. The r's, for example, should be to the nearest four deciral places, rather than only two. They probably used more figures in their computations than they reported to the reader, however.

The multiple correlation coefficient  $R_{1,1,2}$ , which is the product-moment correlation between Y and the best-weighted composite of  $X_1$  and  $X_2$ , could either be calculated directly or found more conveniently by Eq. (9.34):

$$R_{y_1,2} = \sqrt{b_1 r_{y_1} + b_2 r_{y_2}} = \sqrt{(.104).33 + (.390).45} = .46.$$

Notice that the best combination of  $X_1$  and  $X_2$  is scarcely any better as predictor of Y than is  $X_2$  alone. Combining  $X_1$  with  $X_2$  in an optimal way increases the correlation of  $X_2$  with Y from A5 to A6. This "expendability" of  $X_1$  arises from the fact that  $X_1$  and  $X_2$  correlate about equally with Y (with  $X_2$  correlating slightly higher), and  $X_1$  and  $X_2$  are substantially correlated themselves  $(r_1 = .58)$ . The net effect of combining two predictors  $X_1$  and  $X_2$  increases when  $X_2$  and  $X_2$  are both substantially correlated with Y but have a low correlation with each other. Note below how the value of  $R_{1,1,2}$  depends upon the value of  $r_{12}$ :

		Case 1			C	ase 2	
	X <sub>1</sub>	X <sub>2</sub>	Y		$X_1$	$X_2$	Y
X1 .	1.0	0	.50	<i>X</i> <sub>1</sub>	1.0	.5	.5
$X_2$	l	1.0	.50	X <sub>2</sub>		1.0	.5
Y			1.0	Y			1.0
1	$b_1 = .50$ $b_2 = .50$ $R_{y1,2} = \sqrt{.50^2 + .50^2} = .71$			R <sub>v 1,2</sub> =	$b_1 = \sqrt{.33(.5)}$		.50) = .5°

The multiple R is substantially larger in the case in which  $r_{12}=0$ . As you explore the relationships between the correlations of  $X_1$  and  $X_2$  with  $Y_2$ , the correlation of  $X_1$  with  $X_2$ , and the value of  $R_{1,k,n}$  you may have need of an inequality that relates  $r_n$  and  $r_{n1}$  to  $r_{12}$ . If  $r_{n1}$  and  $r_{n2}$  are given, then  $r_{12}$  must satisfy the following inequality:

$$r_{y1}r_{y2} - \sqrt{(1 - r_{y1}^2)(1 - r_{y2}^2)} < r_{12} < r_{y1}r_{y2} + \sqrt{(1 - r_{y2}^2)(1 - r_{y2}^2)}.$$
 (9.35)

This result comes readily by algebraic manipulation of the inequality  $-1 < r_{12}, < 1$ , where  $r_{12}$ , is the  $r_{22}$  of Eq. (9.28). More generally the correlation between variables 1 and 2 cannot be outside the limits

$$r_{13}r_{23} \pm \sqrt{(1-r_{13}^2)(1-r_{23}^2)}$$

For  $r_{11} = r_{22} = 0$ , the limits that  $r_{11}$  may attain are the conventional  $\pm 1$ . For  $r_{12} = r_{12} = 1$  or  $r_{12} = -1$ ,  $r_{12}$  must be 1. For  $r_{13}$  and  $r_{12} = 1$ ,  $r_{13}$  must be 1. For  $r_{13}$  and  $r_{12} = 1$ ,  $r_{13}$  must be 1. Try putting these equations into words to be sure you understand what they mean. See Stanley and Wang (1969) and Glass and Collins (1969) for further details.

Interpreting multiple-regression results causally is fraught with hazards. Probably sociologists have struggled with this problem more than have psychologists or educators. For example, see Coleman et al., (1966). Helpful methodological articles have been provided by Werts (1968) and Pugh (1968).

#### PROBLEMS AND EXERCISES

1. The following are characteristics of people and methods of measuring them:

Characteristic	Measurement
A. Sex	Dichotomous; males-1, females-0
B. Age	Measured in months to nearest month
C. Height	Measured to nearest inch
D. Political preference	Dichotomous; Democrat-1, Republican-0
E. Anxiety	Measured by the judgment of clinical psychologists in ranks from 1 through n for a group of n persons.

F. Intelligence Measured by converting IQ scores to the ranks 1 through n for a group of n persons.

In each of the following instances, identify one or more correlation coefficients

- appropriate for describing the relationship between the two variables in question;
- a. Sex and height.
- b. Anxiety and intelligence.
- c. Sex and anxiety.
- d. Age and height.
  e. Sex and political preference.
- f. Political preference and intelligence.
- 2. The following data are illustrative of data gathered by Kennedy, Van de Riet, and White (1963) and Terman and Mertiil (1960). (Also see Jensen, 1968.) A sample of n = 100 ten-year-old children was drawn from the population of ten-year-olds. Fifty of the children were Negro, and 50 were white. The children were given both digit-span and vocabulary subtests. The following data were obtained:

	Digit span					Voca	Dulary	
		Fail (O)	Pass (1)			Fail (0)	Pass (	1)
	White (1)	25	25	50	White (1)	22	28	50
Race	Negro (O)	29	21	50	Negro (O)	39	11_	50
		54	46	100		61	39	100

- a. Calculate the phi coefficient between race X and performance on the digitspan test Y.
- b. Calculate the phi coefficient between race X and performance on the vocabulary test Y.

- c. Calculate the maximum possible phi that could occur in each of the two situations for the p's there. Hint: p<sub>rect</sub> = .5 each time, but p<sub>digit-span</sub> = .46 and p<sub>sacobalate</sub> = .39.
- Compare the two values of φ calculated above and attempt to interpret
  the difference, both statistically and substantively.
- 3. In a sample of 100 adults, the mean score on an object-assembly test is 104.00 and the unbiased estimate of the sariance of the scores is 256.00. The 50 women in the group have a mean score of 100.00. Scoring females 0 and males 1, calculate the point-biserial coefficient of correlation between sex and object-assembly ability.
- 4. Data were gathered on a test of writing skill (Y) and dichotomously on verbal reasoning (X). The variable X was scored 0 for "below average" and 1 for "above average"; Y was measured by taking raw scores on a 70-item test:

Student	X	r
A	0	52
В	1	52
C	0	44
D	0	55
E	1	58
C D E F	0	52
G	0	61
H	0	38
1	1	53
ſ	0	29
K	0	40
L	0	40
M	0	45
N	1	59
O P	1	57
P	1	50

Calculate r<sub>br</sub>, as an approximation to the correlation between the "underlying" verbal-reasoning scores and writing skill.

5. Equation (9.11) shows that the ratio of rote to rpb is given by

$$\frac{r_{bis}}{r} = \frac{\sqrt{n_1 n_0}}{r}.$$

You are told that  $n_1 = 25$ , n = 50 and  $r_{pb} = -.60$ . Find the value of  $r_{bip}$ 

6. A clinical psychologist and a speech therapist jointly ranked n = 20 children on two variables: X—emotional adjustment (1 = best adjustment, 20 = worst adjustment); Y—severity of stutering (1 = least severe, 20 = most severe). There are C<sub>1</sub><sup>20</sup> = 190 distinct pairs of children in this group. In 80% of these pairs, the child in the pair who had a higher rank on X also had a higher rank.

on Y; in the remaining 20% of the pairs, the child with the higher rank on X had a lower rank on Y than the other child of the pair. What is the value of Kendall's  $\tau$  for these data?

7. For a sample of 89 children, Kabot (see Delacato, 1966, Chap. 14) gathered dichotomous data on reading performance X and laterality Y (consistency with which one "side"—eye, hand, foot—of the body is used). Data were in the form of judgments of poor (0) and good (1) reading performance and low (0) and high (1) consistency in the use of one side of the body:

		Reading p Poor (O)		•
( -1 ( 1	High (1)	18	31	49
Laterality	Low (0)	28	12	40
		46	43	- 00

Assume that normally distributed variables underlie both dichotomies and that their joint distribution is bivariate-normal. Calculate the value of  $r_{nt}$  for the above data as an approximation to the product-moment correlation of reading performance and laterality.

8. The raw scores of 12 high-school students on tests of abstract and verbal reasoning are reported below:

Student	Abstract reasoning	Verbal reasoning		
A	40	37		
В	49	42		
С	44	25		
D	42	40		
E	24	19		
F	48	39		
G	36	27		
H	25	14		
1	45	43		
J	28	16		
K	31	20		
L.	39	35		

- a. Convert the raw scores to ranks (1-12) for each variable and calculate  $r_s$ . b. Using the same ranks generated in (a), calculate  $\tau$ .
- Themes written by ten pupils were judged to be either "creative" (X = 1)
  or "not creative" (X = 0). A ranking of the same students on intelligence
  Y was available (10 = highest, 1 = lowest). Calculate the rank-biserial

correlation coefficient from these data:

Student	X	Y
Α	1	2
В	1	6
С	ī	1
D	0	7
E	0	3
F	0	10
G	1	9
H	1	5
1	0	4
j	0	8

 In a group of female physical-education students Leyman (1967) found the following intercorrelations of measures of grade-point average X, the Scott Motor Ability Test Y, and intellectual aptitude Z:

	Х	Y	Z
X	1.00	.30	.22
<u> </u>		1.00	04
z			1.00

- Find the correlation of the Scott Motor Ability Test Y and GPA in physical education X, with intelligence Z partialed out.
- b. Determine the value of the multiple correlation coefficient R<sub>2,1,4</sub>, i.e., the correlation of physical education GPA with the optimal linear combination of intelligence and motor ability.
- c. Try to interpret these results.
- Goolsby (1967) found the following correlations among the School and College Ability Test (SCAT), the Florida Twelfth Grade Achievement Test (FTGAT), and freshman Grade-Point Average (GPA):

	SCAT	FTGAT	GPA	
SCAT FTGAT GPA	1.00	.70 1.00	.40	
017			1.00	

Intercorrelations

Find the value of the multiple correlation coefficient  $R_{2,1,2}$  between GPA Y, and the best linear combination of SCAT  $X_1$  and FTGAT  $X_2$ . In the process, determine the weights  $b_1$  and  $b_2$  for obtaining the least-squares estimate of the  $\hbar\hbar$  person's standard scores on Y from  $\hbar$  is standard scores on  $X_1$  and  $X_2$ .

 The correlation of Y with X<sub>1</sub> is .91; the correlation of Y with X<sub>2</sub> is .87. Can the correlation of X<sub>1</sub> with X<sub>2</sub> be as low as .10?

# 10

## PROBABILITY

#### 101 INTRODUCTION

The scientist is always attempting to go beyond his data. Even when he is most objective and wants least to generalize, he makes tacit assumptions that his set of data has some stability; if he gathered more data tomorrow it would reflect approximately, though not exactly, the same trend. When least objective, he makes wide, general inferences from what he sees today to what he will see in different places, under different conditions, tomorrow.

Every inference contains some degree of uncertainty. The likelihood of an inference's being true often does not rush upon us, as when we hear: "The Yankees will beat the Sox tomorrow because I saw them beat the Sox today." Other inferences are much more cogent: "I've noted that the sun has risen every day for the past twenty years, hence it will also rise tomorrow risen every day for the past eventy years, hence it will also rise tomorrow morning." Inferences differ in their likelihood of being valid all the way from "extremely unlikely" to "almost certain." By its very nature, no inference is certain to be valid, although some approach certainty.

196 PROBABILITY CHAP, 10

Much of the work of the statistician is derivation of methods that assign probabilities to inferences. This is indeed an important occupation. Inferential reasoning is the method of science; what a clumsy method it would be if scientists had no objective and systematic means of assigning probabilities to the inferences they make. The language of everyday life with which people refer to inferences as "extremely unlikely" or "almost certain" does not serve science adequately. These subjective estimates vary from person to person depending on the words chosen to express the likelihood and on what the words themselves mean. It is far better for communication among scientists that they can independently arrive at the same statement of the probability of the validity of an inference, and that the statement can be made in unambiguous terms carrying the same meaning for all scientists.

We have idealized the actions of scientists only slightly in the above paragraphs to make the point stronger. Scientists and statisticians are not unanimous on the questions how to assign probabilities to statements and to which statements should probabilities be assigned. Nonetheless, although they may construct adjacent but nonintersecting systems, their preference for one aystem springs from a value judgment about what the role of the scientist is and ought to be. Their systems are open for all to see, and anyone who is capable of understanding them may. Although statisticians do not always agree on the proper method, their methods of assigning probabilities to inferences are far better than the use of everyday language and unquantified

Much of the remaining material in this text deals with assigning a probability value to an inference. We shall examine the methods the statistician has developed that allow him to state, for example, "The relationship between IQ and grade-point average is not random in children II years of age, and I make this statement with probability .99 that it is true." Obviously, you must set yourself to the task of learning the mathematical conventions that constitute the beginnings of probability theory. You cannot expect to learn all about probability in the space that is available, for probability is a large and complex body of knowledge. Hopefully, you can learn enough to make elementary statistical methodology meaningful.

#### 10.2 PROBABILITY AS A MATHEMATICAL SYSTEM

We shall begin by being entirely arbitrary. Probability will be looked at as a system of definitions and operations pertaining to a sample space. The idea of a sample space is basic. We shall never make a probability statement

107

that is not related to a sample space of some sort. Indeed, statements of probability are simply statements about sample spaces and their characteristics.\*

A sample space is a set of points. These points can represent anything: persons, numbers, balls, etc. An event is an observable happening like the appearance of "meads" when a coin is flipped or the observation that a person whose name you have selected at random from a telephone book is watching television. There may be several points in the sample space, each of which is an example of an event. For instance, the sample space has 9 points. An event might be "A ball is white." This event has 6 sample-space points. How many points in the sample space does the event "A ball is black" have? The event "A ball is this urn is red" has no sample points. "A ball in this urn is either white or black" is also an example of an event. Notice that many different events can be defined on the same sample spame to

A statement of probability is made about the occurrence of an event that is associated with a sample space. For the next few pages, we shall let a capital letter,  $A, B, C, \ldots$ , stand for an event; the "probability of the event A" will be denoted by P(A).

Definition: The probability of the event A, P(A), is the ratio of the number of sample points that are examples of A to the total number of sample points, provided all sample points are equally likely.

Let A be the event "A ball is white," where the sample space is the set of 9 balls (6 white, 3 black) in an urn. How many sample points are examples of the event A? Obviously, the answer is 6. What is the total number of sample points 79. Hence, the probability of the event A ("A ball is white") is

$$P(A) = \frac{\text{number of examples of } A}{\text{total number of sample points}} = \frac{6}{9} = \frac{2}{3}.$$

If B is the event "A ball is black" in our example, find P(B). Add together P(A) and P(B). In our example, what is P(C) if C is the event "A ball is red"? D is the event "A ball is either white or black." What is P(D)? E is the event "A ball is both white and black." Find P(E).

Suppose you have a second urn which has 4 white balls in it and an unspecified number of black balls. What is the probability of the event that a ball is white? You cannot know; at least, within the system we have developed so far, the question cannot be answered. A probability statement

The notion of a sample space is actually a relatively recent development in probability theory, dating back only to the 1920's and the work of von Mises (1931).

can be made if you know fully the characteristics of the sample space, but in this example you do not.

There exists an alternative route by which we may establish the definition of the probability of an event. Consider a sample space composed of a countable number of elementary exerts. An elementary exert is a sample point. Denote each sample point by a lower case letter "a": a<sub>1</sub>, a<sub>2</sub>....a<sub>n</sub>.

Every event defined on the sample space is composed of a set of elementary events.

Definition: A probability function is a rule of correspondence that associates with each event A in the sample space a number P(A) such that (1) P(A) > 0, for any event A; (2) the sum of the probabilities for all distinct events is 1; (3) if A and B are mutually exclusive events, i.e., have no sample points in common, then P(A or B) = P(A) + P(B).

If we assume that the probability of an elementary event  $a_t$  is 1/n, where n is the total number of sample points, then the probability of the event A that is composed of  $n_t$  sample points is

$$P(A) = \frac{1}{n} + \frac{1}{n} + \ldots + \frac{1}{n} = \frac{n_1}{n}$$
.

 $n_1/n$  is the ratio of the number of sample points that are examples of A to the total number of sample points.

Both routes bring us to the same definition for P(A). While the latter definition might have the preference of the mathematician, the former definition of P(A) will probably seem clearer to you.

### 10.3 COMBINING PROBABILITIES

Suppose we have an urn that contains 4 red, 3 white, and 3 black balls. Three events might be of interest to us: (1) A, a ball is red; (2) B, a ball is white; (3) C, a ball is black. These three events are mutually exclusive: each sample point is an example of one and only one event.

The question arises, "What is the probability that a ball is red or white?" We shall denote this event by the symbol  $A \cup B$  and its probability by  $P(A \cup B)$ .

#### First Addition Rule of Probabilities

When the events A and B are mutually exclusive,  $P(A \cup B)$ , the probability of either A or B, is P(A) + P(B).

In our example.

$$P(A \cup B) = P(A) + P(B) = \frac{4}{10} + \frac{3}{10} = \frac{7}{10}$$

Find  $P(A \cup C) = P(A) + P(C)$ . What is the value of  $P(B \cup C)$ ?

In some sample spaces two events may not be mutually exclusive. i.e.. a single sample point may be an example of both events A and B. Let's a single sample point may be an example of both events A and B. Let's consider the possible outcomes ("heads" or "tails") of flipping a coin three times in a row. We consider these outcomes abstractly; as yet we don't want to talk about the physical act of flipping a coin. The eight possible outcomes are the sample space:

1. <i>HHH</i>	5. TTT
2. HHT	6. TTH
3. HTH	7. THT 8. THH
4. HTT	0

Assuming that all of the eight outcomes are equally likely, i.e., each has probability 1/8, what is the probability of "heads" on the first "flip"? What is the probability of "heads" on flips 1 and 2?

We shall now define two events A and B on the above sample space:

The sample points that are examples of event A are the first two outcomes (numbers 1 and 2 above). Two sample points are examples of event B. Which two are they?

We shall let the symbol  $A \cap B$  denote the new event, "A and B." In we small let the symbol  $A\cap B$  is the event "heads" on flips 1 and 2 and "heads" on our example  $A\cap B$  is the event "heads" on flips 2 and 3. If we assume that all the eight sample points are equally likely, then the probability of the event  $A \cap B$  is as follows:

ly, then the probability of the event 
$$A \cap B$$
 is number of sample points that are examples of  $A \cap B$  total number of sample points

The total number of sample points is 8. Only one sample point is an example of the event "heads" on flips 1 and 2 and "heads" on flips 2 and 3. Which one is it? So the probability of the event  $A \cap B$  is 1/8.

Look back at the First Addition Rule of Probabilities and you'll see Look wask at the 1 as Adomotic Rate of Floradinius and you'll see the assumption that the two events A and B are mutually exclusive. In the and assumption that the Anna B were not mutually exclusive. The outcome example just discussed A and B were not mutually exclusive. The outcome example f was an example of both events A and B. Keep the same definitions for A and B and let's see what the probability of A or B,  $A \cup B$ , is when A and B are not mutually exclusive.

#### Second Addition Rule of Probabilities

The probability of either event A or event B or both,  $P(A \cup B)$ , is  $P(A) + P(B) - P(A \cap B)$ .

This expression might look strange; a Venn diagram, an illustration of the relationship between events defined on sample spaces, should help clear up the mystery of the term  $^{-}P(A \cap B)$ ." See Fig. 10.1 where the events A and B, two overlapping groups of sample points, are depicted in the sample space S. We'll assume that the probability of event A is the area of circle A, and that the probability of event B is the area of circle B.

The probability of A or B or both is that area covered by the intersecting circles A and B. The shaded portion in Fig. 10.1 is that set of sample points in both events A and B, i.e., those points that are examples of  $A \cap B$ .

How do we find the area covered by A and B? We first find the area of A that is not shared by B, add to it the area of B not shared by A, and then add the area of A and B.

$$P(A \cup B) = [P(A) - P(A \cap B)] + [P(B) - P(A \cap B)] + P(A \cap B).$$

The first two terms on the right side of the above equation give the area of A minus the area in common with B. Terms three and four give the area of B minus its area in common with A. We complete the area we wish to find by adding in the area common to A and B,  $P(A \cap B)$ . The above equation simplifies to:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

Hence, we have established the Second Addition Rule of Probabilities. Notice that if we had simply added P(A) and P(B) to find  $P(A \cup B)$ , the portion in common to A and B,  $P(A \cap B)$ , would contribute twice to the sum. It must contribute only once, so consequently it must be subtracted once.

Even though we have presented two separate rules for the addition of probabilities, the first rule is just a special case of the second rule—that is, the case when  $P(A \cap B) = 0$ .

If A and B are mutually exclusive events in S, then they don't overlap. See Fig. 10.2; there is no area in common to A and B, so  $P(A \cap B) = 0$ .



FIG. 10.1 Venn diagram of the intersecting events A and B in the sample space S.

701




FIG. 10.2 Venn diagram of the mutually exclusive events A and B in the sample space S.

In general,  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ . If A and B are mutually exclusive, then  $P(A \cap B) = 0$ . Therefore, if A and B are mutually exclusive then

$$P(A \cup B) = P(A) + P(B) - 0 = P(A) + P(B)$$

So far we have treated probability rather abstractly. In much the same manner as the mathematical systems of geometry and algebra, probability theory can be developed from a small set of axioms and definitions. But also in the same manner as geometry and algebra, probability theory can serve as a model for what is going on in a certain class of events in the world around us. Geometry has applications to surveying and architecture. The numerous applications of algebra are obvious. Probability theory also applies to a certain group of events.

The principle that relates probability statements to physical events is due to James Bernoulli (1654-1705). An example of the application of a formal probability statement to an actual set of actions should suffice to show the relationship between theory and application.

Suppose we have an urn that contains 4 white and 6 black balls. The balls are identical in size, shape, and weight and thoroughly mixed so that if we were to reach in and pull one out, it is equally likely that any one of the ten balls would be selected; each ball has one chance in 10 of being chosen. We reach into the urn, bull out a ball, and record its color. The ball is returned to the urn, the balls in the urn are stirred thoroughly, and the act is repeated under the same conditions. We perform this act a very large number of times, say 10,000. After the 10,000th drawing of a ball, suppose we count the number of times a white ball was drawn. Your intuition would tell you that the ratio of the number of times a white ball was drawn. Your intuition would be very close to 4/10. The ratio will very probably not be exactly 4/10 but it will be close.

If we regard the 10 balls as a sample space, and if we say that A is the event "a ball is white," then P(A) is 4|10, exactly. The question arises, Will the formal probability of an event as calculated from theory correspond closely to the relative frequency of the occurrence of the event? The answer to this question is the key to the relationship of probability theory and its application, and the answer is "yes."

We shall attempt a formal statement of this relationship. Suppose an event A either does or does not occur on every trial of an act. The prob-

202 PROBABILITY **CHAP. 10** 

> ability that A will occur is the same for all trials of the act and equals P(A). For example, the act may be flipping a symmetrical coin, A may be the event "heads," and it is assumed that the probability of heads, 1/2, is the same from one flip to the next. It is also assumed that every trial is independent of (in no way affected by) every other trial. Now after n trials of the act. the proportion of times A has occurred is p. It can be proved (the proof is so difficult there is no point presenting it here) that p gets closer and closer to P(A) as n becomes larger and larger. We can make the proportion of times A occurs as close as we want to P(A), the probability calculated from the sample space, by performing the act enough times. So P(A) tells what will happen in the long run if we actually performed the actions under the conditions laid down above.

> The preceding paragraph is a verbal statement of the law of large numbers. The law of large numbers is important for the application of probability and since statistics is one such application, it is important for statistical inference. In spite of its importance, we shall have little more to say about the law of large numbers.

#### Multiplicative Rule of Probabilities

There exists a multiplicative rule for probabilities that will be of considerable importance to us in later work. Suppose we are flipping a coin five times in a row. Assume that the probability of "heads" is 1/2 on each flip and that the flips are independent. The multiplicative rule for probabilities states that the probability of getting five straight "heads" is 1/2 · 1/2 · 1/2 ·  $1/2 \cdot 1/2 = 1/32$ . A general statement of the rule follows:

Multiplicative Rule of Probabilities: The probability that A, which has probability P(A) of occurring on any one trial, will occur n times in n independent trials is

$$P(A) \cdot P(A) \cdot \ldots \cdot P(A) = P(A)^n$$

The following examples will illustrate the probability rules developed so far. Suppose one rolls a die (singular of dice) on which the probability of any one of the 6 faces coming up is 1/6. What is the sample space of outcomes for one toss? The sample space is the set of 6 outcomes 1, 2, 3, 4, 5, and 6. What is the probability of the die's coming up 2? What is the probability that an even number will appear on a single toss of the die?

 $P(\text{even number}) = P(2) + P(4) + P(6) = \frac{1}{4} + \frac{1}{4} + \frac{1}{4} = \frac{3}{4} = \frac{1}{4}.$ 

What is the probability of an even or odd number appearing on a single toss? Suppose we consider two rolls of the die. The sample space of possible

	1, 1	2,1	3, 1	4, 1	5, 1	6.1
	1, 2	2,2	3, 2	4, 2	5, 2	6, 2
	1, 3	2,3	3, 3	4, 3	5, 3	6.3
FIG. 10.0 G . 1	1,4	2,4	3,4	4,4	5, 4	6,4
FIG. 10.3 Sample space of out-	1,5	2,5	3, 5	4,5	5, 5	6,5
comes of tossing a die twice.	1,6	2,6	3, 6	4,6	5, 6	6,6

outcomes has 36 points, as shown in Fig. 10.3. Let event A be "a 1 on toss 1" and event B "a 2 on toss 2." Find  $P(A \cap B)$  by dividing the number of sample points that are examples of  $A \cap B$  (both A and B) by 36. Verify that  $P(A \cap B)$  is equal to  $P(A) \cdot P(B)$ . Find  $P(A \cup B)$ , the probability of event A or event B, remembering that  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ . Here, this is 6/36 + 6/36 - 1/36 = 11/36; the point 1, 2 is common to A and B.

We shall accept as a definition the statement that two events are independent if and only if  $P(A \cap B) = P(A) \cdot P(B)$ . Independence is an important concept in statistics and probability, and we shall have more to say about it later.

# 10.4 PERMUTATIONS AND COMBINATIONS

Two additional concepts that crop up repeatedly when one begins examining the outcomes of experiments are permutations and combinations.

A permutation of a set of objects (the letters A, B, C, and D, for example) is an arrangement of them in which their order is considered. A different ordering of the objects is a different permutation. How many different permutations (orderings) are there of the letters A, B, C, and D? To find out we can set about the task of writing them down and counting them, as shown in Table 10.1.

If we have n distinct objects we can make n(n-1)(n-2), ...2 · 1 different permutations of them. Instead of writing n(n-1)(n-2) ...2 · 1 we can denote this product by n!, read "n factorial," n! is the product of the numbers from 1 through n and equals the number of permutations of n distinct objects. (We agree to let 0! equal 1.)

It can be shown fairly easily, and you should satisfy yourself that it is true, that

$$n(n-1)(n-2)\dots(n-n_1+1)=\frac{n!}{(n-n_1)!}$$
;

merely write n! as

$$n(n-1)(n-2)\dots[(n-n_1)+1](n-n_1)[(n-n_1)-1]\dots 1$$

Write out n! in the numerator and  $(n - n_1)!$  in the denominator and cancel the terms that are common to both numerator and denominator and express  $(n-n_1)[(n-n_1)-1]\dots$  as  $(n-n_1)!$  Then perform the division. Consequently.

$$\binom{n}{n_1} = \frac{n!}{n_1! (n - n_1)!}.$$
where of parameters are not to the second state of the second state

To summarize, the number of permutations of n objects is n!. The number of permutations of n objects taken n, at a time is

$$n(n-1)\ldots(n-n_1+1).$$

The number of combinations of n things taken n<sub>1</sub> at a time is given by

$$\binom{n}{n_1} = \frac{n!}{n_1! (n-n_1)!}.$$

How many different combinations are there of 3 things taken from 5 things? In this problem n = 5 and  $n_1 = 3$ .

$$\binom{5}{3} = \frac{5!}{3!(5-3)!} = \frac{5!}{3!2!} = \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{(3 \cdot 2 \cdot 1)(2 \cdot 1)} = \frac{5 \cdot 4}{2 \cdot 1} = \frac{20}{2} = 10.$$

Note that there are  $5 \cdot 4 \cdot 3 = 60$  permutations of 5 things taken 3 at a time because each of the 10 combinations of 5 things taken 3 at a time has 3 · 2 · 1

Ten men are eligible to serve on a committee. The committee must be composed of only seven men. How many different committees could be formed from the 10 available men? Let n = 10 and  $n_1 = 7$ ; then evaluate

An ice cream manufacturer has six different flavors of ice cream. He can sell them individually, he can mix any two different flavors, or he can mix any three different flavors to make new flavors. How many different flavors (and combinations of flavors) can he make? You must evaluate three separate quantities of the form  $\binom{n}{n}$  to solve this problem.

#### BINOMIAL DISTRIBUTION

Practically every topic discussed so far in this chapter can be applied to the solution of a problem in statistics. The following problem concerns the repeated independent performance of an act that can result in either "success" or "failure" with a constant probability. We shall develop a binomial distribution that will describe one aspect of the results of such a series of acts, namely, the probability of a given number of successes.

Suppose a fair coin, i.e., symmetrical and of homogeneous composition, is to be flipped five times in succession. Each flip is independent of every other flip in the sense that if the first flip results in "heads," then a "heads' on the second flip is no more nor less likely than if the first flip had given a "tails." Assume further that the probability of "heads" is the same from the first to the fifth flip. The five flips are five "trials", independent trials that can result in one of two outcomes with a constant probability are called Bernoulli or binomial trials. (James Bernoulli was a 17th-century mathematician whose Ars Conjectandi was one of the first treatises on probability theory.) The five coin flips are five binomial trials. How does one find the probability that three "heads" will result in five trials?

One possible outcome of the five flips that gives three "heads" and two "tails" is H, H, H, T, T. From the multiplicative rule of probabilities, we know that the probability of H, H, H, T, T is  $|V_2| + |V_2| + |V_2| + |V_2| + |V_2| + |V_2| + |V_2|$  since the probability of "heads" and the probability of "tails" are both  $|V_2|$ . However, the sample space of outcomes of the five flips has  $|V_2| + |V_2| +$ 

$$\binom{5}{3} = \frac{5!}{3! \, 2!} = \frac{4 \cdot 5}{2 \cdot 1} = 10.$$

There are 10 sample points that are examples of the event and each has probability 1/32; consequently, we use the addition rule of probabilities so that the probability of obtaining three "heads" in five flips of a fair coin is 10(1/32) = 10/32.

In summary, 10/32 was found by multiplying out  $\binom{5}{3} \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2$ .

In general, the probability of obtaining  $n_1$  "successes" in n binomial

trials is equal to

$$\binom{n}{n_1} p^{n_1} q^{n-n_1},$$

where p is the probability of a "success" on any one trial,

q is the probability of a "failure" and equals 
$$1-p$$
, and  $n_1=0,1,2,\ldots,n$ .

Note that "binomial trials" refers only to trials that have only two possible outcomes: yes or no, success or failure, heads or tails, red or not red, etc.

A die is tossed four times, and we wish to find the probability of three "sixes." In this example, n=4,  $n_i=3$ , and we take the probability of a "six" on any trial, p, to be 1/6. Hence, q=1-p=5/6, the probability of not a 6. The probability of 3 "sixes" thus equals

$$\binom{4}{3} \left(\frac{1}{6}\right)^3 \left(\frac{5}{6}\right)^1 = \frac{4!}{3! \cdot 1!} \left(\frac{1}{6}\right)^3 \left(\frac{5}{6}\right) = \frac{4 \cdot 5}{6^4} = \frac{20}{1296}.$$

You can see that the probability of three "sixes" in four tosses of a die is small, less than .02.

The expression

probability 
$$(n_1) = \binom{n}{n_1} p^{n_1} q^{n-n_1}$$

is the formula for the binomial distribution. This distribution has many important uses in theoretical and applied statistics.

An interesting relationship exists between the term  $\binom{n}{n_1}$  in the binomial distribution and a table known as Pascal's triangle [after the French mathematician and philosopher Blaise Pascal (1623-1662)]. Each row in Pascal's triangle (see Fig. 10.4) is formed by adding together certain numbers in the row above it. Each number is the sum of the two numbers diagonally above it. Thus, the 6 in row 4 is the sum of the two 3's above it. You could casily construct the eighth, ninth, tenth, and subsequent rows yourself.

It can be proved that  $\binom{n}{n_i}$  equals the  $n_i+1$  st number in the nth row of Pascal's triangle. For example,  $\binom{3}{i}$  equals the second,  $(n_i+1)$ st, number in row 3:

$$\binom{3}{1} = \frac{3!}{1! \ 2!} = 3.$$

 $\binom{5}{0} = 1$  is the first,  $(n_1 + 1)$ st, number in row 5; it is the number of combinations of zero things, taken from 5. Note that the first and the last

FIG. 10.4 Pascal's triangle.

number in each row is 1; because  $\binom{n}{0} = \binom{n}{n} = 1$ .

Calculating, for example, the probability of 14 successes in 30 trials when p=3/8 and q=5/8 would be an arduous task. Fortunately, excellent tables for the binomial distribution exist. We recommend you use them whenever your calculations are more complicated than the elementary once in this chapter. A brief table for the binomial distribution appears in Pearson and Hartley (1966). Much more extensive tables have been published. See Staff, Harvard University Computational Laboratory (1956) and "Tables of the Cumulative Binomial Probabilities" (1953). (The "cumulative binomial" is simply the sum of successive terms in the binomial distribution and gives the probability of at least  $n_s$  successes in n trials.)

Figure 10.5 illustrates the entire binomial distribution for eight trials when p = q = 1/2. Note that when p = q the distribution is symmetrical. For p > q it will be skewed to the left because high values will predominate, thou will it be skewed for p < q?

In Fig. 10.6 the binomial distribution for n=5 and p=.30, q=.70 appears. You can see from Fig. 10.6 that the probability of five successes when n=5 and p=.30 is very small. It is only  $(.30)^5=.00243$ , i.e., (.34) times in  $(.30)^5=.302$ . What values of  $(.30)^5=.302$  what values of  $(.30)^5=.302$  what values of  $(.30)^5=.302$  when  $(.30)^5=.302$ 

The binomial distribution affords a good example of the general type of reasoning used in testing statistical hypotheses. A statistician may feel



FIG. 10.5 Binomial distribution for n = 8, p = 1/2.

210 PROBABILITY CHAP. 10



FIG. 10.6 Binomial distribution for n = 5, p = .30. that the process by which the data he has observed were generated was a

series of binomial trials. Suppose he has flipped a coin ten times and observed nine "heads." He is suspicious that the coin is not fair, i.e., the probability of getting "heads" is not 1/2. His observations appear to be evidence that the coin is unfair. He formalizes his evidence in the following manner:

- 1. Entertain the notion temporarily that the probability of "heads" for this coin truly is 1/2. This is a hypothesis.
- 2. If the probability of "heads" is 1/2, what is the probability that I would have observed the event that I did (nine "heads") or one more extreme (10 "heads")?
- 3. This probability is

$$\binom{10}{9}\binom{1}{2}^{\bullet}\binom{1}{2}^{\bullet}+\binom{10}{10}\binom{1}{2}^{10}\binom{1}{2}^{0}=.0107.$$

(Recall that any number with an exponent of zero equals 1.)

- 4. If this coin is fair, I have observed an event that is extremely unlikely.
- 5. If this coin is biased in favor of "heads," the event I observed would be more probable. For example, if the probability of "heads" for this particular coin were actually 8/10, then the probability of getting nine or 10 "heads" in 10 flips would be

$$\binom{10}{9} \left(\frac{8}{10}\right)^{\!\!\!\!0} \! \left(\frac{2}{10}\right)^{\!\!\!0} + \binom{10}{10} \! \left(\frac{8}{10}\right)^{\!\!\!10} \! \left(\frac{2}{10}\right)^{\!\!\!0} = .3758.$$

6. It appears that on the basis of my observations it is very unlikely that this coin is fair. Therefore, the hypothesis in (1) is rejected; the coin appears to be biased in favor of "heads."

Notice that it is certainly not impossible to obtain nine "heads" in 10 flips of a coin; in fact, we should expect it to happen about once (1.07) in every 100 sets of 10 flips. The possibility exists that the coin that yielded nine "heads" actually is a fair coin, but the probability that this is true is small.

The experimenter's logic often follows the same pattern as the reasoning above. He hypothesizes that certain features of a model for his experiment the hypothesis: the hypothesis a model have certain values (e.g., the probability of "heads" is 1/2). He needn't believe this hypothesis; he only entertains it temporarily to see where it leads him. The experimenter then takes his observations (e.g., flips the coin 10 or 20 times), after which he calculates the probability of obtaining the actual result he did obtain, or a more extreme one, if his original hypothesis was true. If this probability is very small (say, .05 or .01), he questions the truth of the hypothesis. If his original hypothesis was false and some alternative hypothesis was true (e.g., the coin is biased), his results might be much more likely. This process, called "hypothesis testing," is one of rejecting or failing to reject some explanation (hypothesis) for the obtained results on the basis of the probability of obtaining the particular results if the hypothesis were true. If the probability of the obtained results (or more extreme ones) is large (for example, .30 or more) when one calculates this probability under a certain hypothesis, then the hypothesis is not rejected at that moment. If the probability of the obtained results (or more extreme ones) is small (.05 or less) when this probability is calculated under another hypothesis. then this hypothesis is rejected. It has not been proven that the hypothesis is false: there is some small probability (.05 or less) that it is true. However, the evidence for its falsity is great.

The study of probability can be interesting and entertaining. Historically, probability concepts arose in connection with games of chance. Those who make use of probability theory are generally awed by the intricacy and excitement of the system and the way in which it produces results quite in disagreement with intuition, unless one's intuition has been developed by experience with calculated probabilities. A few examples of surprising results will convince you of the untrustworthiness of your intuition at this stage.

What is the probability that at least two people in a group of 23 have the same birthdate? (Assume that the people are drawn randomly from population of persons in which all 365 birthdates (not counting February 29) are equally likely.) Is the probability 000 or .0001 or even smaller? Actually, the probability that at least two people out of 23 have the same birthdate is .501? You should expect multiple birthdates to occur in slightly more than half the groups of size 23. It is practically certain that in a group of 150 persons at least two people will have the same birthdate! (See Feller, 1957, pp. 31-32.)

Suppose John and Jim have a fair coin (probability of "heads" is 1/2)

educational and psychological research. Careful study of the proper randomsampling procedure and how it can be abused will be worthwhile.

The condition of equiprobability in the definition of random sampling states that at each stage in the sampling process all of the remaining elements have the same probability of being chosen. Suppose two cards are to be selected at random from a deck of 52. If the first choice is to be random. then all 52 cards must have an equal chance of being chosen, namely 1/52. If the second choice is to be random, all of the 51 remaining cards must have the same probability of being chosen, namely 1/51. How do you achieve this goal of equiprobability in practice? There are numerous ways in which equiprobability is approximated; some are much better than others. Probability statements about various "hands" in poker and bridge assume that the cards are shuffled in a purely random manner. The greater incidence of "royal flushes" and "perfect hands" than would be predicted by probability attests to the fact that a casual shuffling of the cards is a poor approximation to randomness. Drawing names from a hat is an equally bad attempt to approximate randomness. The best method of achieving random sampling approximate randomness. The oest method of achieving random sampling available to the practitioner of statistics is use of a table of random numbers. Tables of random numbers are tables of digits, 0 through 9, ordered by

a mechanical process that approximates equal probability of the occurrence of any one digit as nearly as technology can. If you select your sample with the aid of a well-known table of random digits and if you use an approximate procedure, you need have no worry that your sample will not be random. A table of random digits appears as Table A in Appendix A.

The following is an example of random sampling using a table of random digits. From the 900 male members of the PTA, Principal Jones wishes to estimate the percent of fathers with high-school diplomas from a random sample of 200. The percent with diplomas in the random sample of 200 will not differ much from the corresponding percent in the population of 900. Jones instructs his secretary to assign a three-digit number from 001 to 900 by starting at the beginning of the alphabetical file of PTA members. He then goes to his table of random numbers to make his selection. With the book open to the first page, he lays the point of his pencil on the page of random digits with his eyes closed. The two digits closest to his pencil point are 3 and 6. He moves to the intersection of row 3 and column 6 of the table to begin selections. Moving along row 3 starting with column 6 Jones sees the digits 78 2 1 2 1 6 9 9 3 3 5 9 0 2 9 1 3 ... By grouping digits into groups of three, he obtains the numbers 782, 121, 699, 335, 902, 913, . . . His secretary will now pull the cards of those persons who have identification numbers 782, 121, 699, 335, etc. But what about 902 and 913? There are only 900 persons, so no one has identification numbers 902 and 913. These numbers are simply disregarded. What happens when 782 comes up again, if it does, or the number of any other person who has already been selected? SEC. 10.7 PANDOM MARIABLE 215

studies at the master's and doctor's level and many others that are published in journals stumble on this basic problem. Examples of proper and improper analyses will be given throughout this text; you are best advised, however, to seek expert advice concerning your sampling technique when you are designing an experiment.

A common fault of naïve researchers in education is to choose "classrooms" randomly to participate in an experiment and then to analyze the data as if "students" had been chosen randomly. If classroom A is chosen for method A of teaching and classroom B for method B, then surely the selection of "students" has not been an independent one. If John and George are both in classroom A, then John and George must both receive method A. There's no chance for John to receive A and George to receive B. Any analysis that treats the 30 students in both A and B as though they constituted 60 separate observations is quite likely to be wrong because the two classrooms are just two "clusters" of students. The average of the 30 observations in A may be treated as independent of the average of the 30 observations in B. if other conditions are met. Hence, one has two observations that constitute a random sample. This is a difficult point to comprehend. The general problem assumes a slightly different form for all of its special instances. The proper method of sampling and analysis sometimes manages to elude the most sophisticated researchers. We hope to throw more light on the problem later in this text.

We conclude this section on random sampling with an alternative statement of the definition of simple random sampling. Try to satisfy yourself that this statement is equivalent to the one given at the beginning of this section

Simple random sampling: If sampling from n observations is random, then regardless of what the first  $n_1$  choices were, the probability of any particular observation's being chosen on the  $(n_1 + 1)$ st selection is  $1(n_1 - n_1)$ .

#### 10.7 RANDOM VARIABLE

A random variable is defined in terms of two concepts, one of which has already been introduced: sample space and function. We have already discussed the meaning of a sample space. It is a collection (finite or infinite) of objects or events. A function is any set of ordered pairs of elements, no two of which have the same first element. As you can see, the definition of a function is quite general;  $\{(a, 1), (b, 2), (c, 3)\}$  is a function, so is  $\{John, Alice), \{Joe, Mary\}, (Ted, Sharon), (Jim, Joyce)\}. A function is formed when$ 

we specify a rule that associates a unique element with every element of some set. We form a function when we associate every person with his age: (Mr. Jones, 37), (Mark Smith, 5),... A random variable is a function such that all of the first elements are points in a sample space. The previous example is a random variable. The sample space is all persons; each is associated with his age.

The following are all examples of random variables (the random variables of most interest to statisticians and to us in this text are those in which the elements of a sample space are associated with numbers):

1. Sample space: outcomes of the flip of a coin ("heads," "tails").

	variable X 2nd elemen		
"heads"	1		
"tails"	0		

2. Sample space: the 6 different outcomes of rolling a die.

Random variable Y		
1st element	2nd element	
the face with 1 dot	1	
the face with 2 dots	2	
•		
•		
the face with 6 dots	6	

3. Sample space: a bushel of oranges.

R 1st element	andom variable Z  2nd element (the weight of the orange in ounces)
Orange #1 Orange #2 Orange #3	3 5
Orange #3	2
-	•

The idea of a random variable is so simple and seems so obvious that a fancy definition appears to be "window dressing." Confusion can arise, however, if the definition is not made explicit.

We shall speak later of "the value of the random variable Z." If Z is the random variable as defined in (3) above, then Z assumes a value, the weight in ounces, for each element of the sample space, each orange. For the first orange, Z assumes the value 3.

You have already met a random variable in this chapter, but no attempt was made to point it out. The binomial distribution is that of a random variable. The sample space is the collection of points that are the distinct outcomes of n binomial trials. The random variable X takes on the values  $0, 1, 2, \ldots, n$  according to the number of "successes" that occur in n trials. If a coin is flipped four times and a "heads" is called a "success," then X has the value 3 for the event H, T, H, H, since 3 "successes" occurred.

A sample space can become complex conceptually. For example, a child is to read a given page of a reader. He is to do this a large number of times; each separate reading of the page is a sample point in the sample space of all readings. A random variable X can be defined on this sample space by associating with each reading the time from beginning to end. X takes on a value for each reading: reading 1—3 min 5 sec; reading 2—2 min 48 sec; ... X is a random variable that takes on values expressed in "minutes" and "seconds."

In subsequent chapters we shall see how probabilities are associated with values of a random variable. For example, suppose a fair coin is being flipped. The random variable X is the same as example (1) above. The probability that X will take on the value 1 is 1/2, the probability that the event "heads" associated with 1 will occur. The probability that the random variable X will assume the value 0 is 1/2.

# 10.8 TYPES OF RANDOM VARIABLES

The statistician finds the distinction between discrete and continuous random variables a useful one. We have had no need to use this distinction yet; it becomes a useful one when certain operations on random variables are defined and performed.

The distinction between discrete and continuous refers to the nature of the numbers that are associated as second elements of the random-variable function with sample-space points. A discrete random variable is one that can assume only certain values on the real-number line. We can think of all real numbers as points on a line that extends from  $-\infty$  to  $+\infty$ , as in Fig. 10.7. Between two points on the real-number line, a discrete random variable

can assume some values but not others. An example of a discrete random variable is X in (1) in Sec. 10.7 above. It can assume only the values 0 and 1; it is impossible for X to equal 1/2. A continuous random variable is one

CHAP. 10 218 PROBABILITY

> that can assume any value on the real-number line between two points. An example of a continuous random variable is age. The value of the random variable "age" for a person can equal 5 years, 6 months, 4 days, 11 hours, 14 minutes, 6.132 ... seconds. A person who has just become 10 years old has possessed an age equal to every possible number on the real number line from 0 to 10 years.

Which of the following random variables are discrete and which are continuous?

- 1. Number of "heads" in six flips of a coin.
- 2. Time required to solve a concept formation task.
  - 3. Height of mercury in a barometer.
- 4. Highest temperature of the air during the daylight hours.
- 5. The number of teeth in an infant's mouth.
- 6. The amount of money in the pocket of a corporation president.

Variables 1, 5, and 6 are discrete; the others are continuous. Can a corporation president have 11.5 cents (not counting trading stamps) in his pocket? Can a subject require 51.23 seconds to solve a concept formation task? Can be require 46.721 seconds or 38.50 seconds?

A distinction must be made between the values that a random variable can assume theoretically and the values that one's measuring instruments yield. A variable such as length can assume theoretically any possible real number between 0" and 5", say. For example, analytic geometry tells us that the hypotenuse of a right triangle both sides of which are 1" in length has length  $\sqrt{2}$ . The number  $\sqrt{2}$  is an unending number the first few digits of which are 1.414.... No physical act of measurement will yield a value exactly equal to  $\sqrt{2}$ . The most sensitive measuring instrument must finally give up in its attempt to add more figures after 1.414 . . . . Perhaps a finely calibrated ruler could give lengths to the nearest 1/128th of an inch; if so, length as measured by this ruler could not assume a value between 6/128" and 7/128\*, for example. Although "length" is theoretically a continuous random variable, any measurement of it yields discrete values. Nonetheless, it will be helpful to retain the distinction between discrete and comingous while remembering that the physical act of measurement yields discrete numbers

#### 10.9 PROBABILITY AS AN AREA

The probabilities of observing values of continuous variables, e.g., height, are conveniently represented by mathematical curves known as probability distributions. Suppose we have a continuous random variable X that can




FIG. 10.8 The probability density function of the variable  $X_t$ , time required to solve a puzzle.

take on values from 0 to 10. For example, X could be the time required for subjects to solve a certain puzzle. They may solve it almost immediately, or they may take as long as 10 minutes but no longer. Presumably the length of time required to solve the problem is known for a huge number of different subjects. A graph is drawn in which the "time to solution" is graphed against "proportion of subjects requiring that time" (see Fig. 10.8).

The proportion of subjects requiring between 2 and 4 minutes to solve the puzzle can be regarded as the probability that a subject selected at random from the population will require between 2 and 4 minutes to solve the puzzle. The area under the curve in Fig. 10.8 is 1, so the area under the curve in Fig. 10.8 is 1, so the area under the curve between any two points  $X_1$  and  $X_2$  is the probability that a randomly selected subject will require between  $X_1$  and  $X_2$  minutes to solve the puzzle. The probability that a randomly selected subject will take more than 5.3 minutes is equal to the shaded area in Fig. 10.8. What area corresponds to the probability that a randomly selected subject will take less than 0.5 minutes? (Theoretically, the probability of a subject's taking exactly 4 minutes, say, is zero because there is no area between 4 and itself.) If the area under the curve in Fig. 10.8 between 6 and 10 were .07, then in a group of 100 randomly chosen subjects we would expect about 7 of them to take between 6 and 10 minutes to solve the puzzle.

The statistician frequently plots the values a continuous random variable can assume in such a way that the area between any two values of the variable equals the probability that the variable will assume a value between those two values. The resulting graph is called a probability density function. The graph can often be expressed as a mathematical function in such a way that the ordinate P(X) can be found by substituting any value of the random variable X. For example, assume X is a random variable that can take on any value between 0 and 2 with equal probability. If we let P(X) = 1/2 for all X, then the resulting graph (Fig. 10.9) will be the probability density function of X.

The area under the curve, e.g., of the rectangle in Fig. 10.9, is exactly 1  $(0.5 \cdot 2.0)$ . The shaded area is the probability that X takes on a value between 0 and 1. What does this probability equal?




FIG. 10.9 Probability density function of the variable X that assumes all possible values between 0 and 2 with equal probability.

CHAP. 10

# 10.10 EXPECTATIONS AND MOMENTS

220

Moments are characteristics of distributions defined in terms of expectations. We shall consider the definition of the expectation of a random variable X first.

Definition: If X is a discrete random variable that takes on the values  $X_1, X_2, \ldots, X_n$  with probabilities  $p_1, p_2, \ldots, p_n$ , then the expectation of X denoted by E(X) is defined as

$$E(X) = p_1 X_1 + p_2 X_2 + \ldots + p_n X_n = \sum_{j=1}^{n} p_j X_j,$$
where  $p_1 + p_2 + \ldots + p_n = 1$ .

Note how similar this is to a formula for computing the mean from grouped data:

$$\overline{X}_{\cdot} = \left(\frac{f_1}{N}\right) X_1 + \left(\frac{f_2}{N}\right) X_2 + \ldots + \left(\frac{f_n}{N}\right) X_n.$$

However, expectations are only for infinite populations, not for samples.

Another symbol denoting the expectation of X is  $\mu$ , the Greek letter "mu."  $\mu = E(X)$ , the mean of the infinite population of X's.\*

\* The mathematically sophisticated reader may note that

$$\mu = E(X) = \lim_{n \to \infty} \left( \frac{\sum_{i=1}^{N} X_i}{N} \right),$$

i.e., the mean of the infinite number of  $X_i$ 's. In this notation,

$$p_i = \lim_{N \to \infty} \left( \frac{f_i}{N} \right)$$
,

the proportion of all the scores that are X/s. n, the number of different X's, may be finite, but N must be infinite in order to define the p/s, or the p/s may be theoretical values, such as  $p_n = .5$  for one random flip of an unbiased coin.

The names "expectation" and "expected value" are synonymous. Some examples of expectations are as follows:

 Suppose X is the random variable that has 6 possible values, 1,2,...,6. The events of the sample space could be the 6 sides of a die. Assume that a probability of 1/6 is associated with each value of X. What is the value of E(X)?

$$E(X) = \frac{1}{6} \frac{1}{6} + \frac{1}{6} \frac{2}{6} + \frac{1}{6} \frac{3}{6} + \frac{1}{6} \frac{4}{6} + \frac{1}{6} \frac{5}{6} + \frac{1}{6} \frac{6}{6} = \frac{1}{6} (1 + 2 + \dots + 6)$$

$$= \frac{1}{6} \frac{1}{6} = \frac{3}{6} \frac{1}{6} = \frac{3}{6} \frac{1}{6} + \frac{1}{6} \frac{1}{6} + \frac{1}{6} \frac{1}{6} + \frac{1}{6} \frac{1}{6} = \frac{1}{6} \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} \frac{1}{6} + \frac{1}{6} \frac{1}{6} + \frac{1}{6} \frac{1}{6} + \frac{1}{6} +$$

halfway between the smallest value, 1, and the largest value, 6:

$$\frac{1+6}{2} = 3.5.$$

In this example,  $E(X) = \mu = 21/6 = 3.5$ . In repeatedly rolling the die, one can "expect" to average 3.5 points.

2. A particular slot-machine has payoffs of \$0.00, \$0.50, \$1.00, and \$2.00. The probabilities associated with each of these occurrences are .80, .15, .04, and .01, respectively. Define a random variable X that takes on the four values 0, 50, 100, and 200 cents with probabilities .80, .15, .04, and .01. What is the value of E(X)?

$$\mu = E(X) = .80(0) + .15(50) + .04(100) + .01(200)$$

$$= 0 + 7.5 + 4.0 + 2.0 = 13.5.$$

If it costs \$0.25 for each trial on this slot-machine, would you like to play?

 Let X be the random variable that corresponds to the number of "heads" in 4 flips of a fair coin. X can take on the values 0, 1, 2, 3, and 4. Find E(X).

First we must calculate the probabilities that X will take on each of the values from 0 to 4, i.e., the probabilities that there will be 0 "heads," . . . , 4 "heads" in 4 flips of a fair coin.

Probability 
$$(X = 0) = {4 \choose 0} \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^4 = \frac{4!}{0! \ 4!} \cdot 1 \cdot \frac{1}{16} = \frac{1}{16}$$

Probability 
$$(X = 1) = {4 \choose 1} \left(\frac{1}{2}\right)^1 \left(\frac{1}{2}\right)^3 = \frac{4!}{1! \ 3!} \cdot \frac{1}{16} = \frac{4}{16} = \frac{1}{4}.$$

If you have forgotten how to compute the remaining three probabilities, refer back to the relevant sections of this chapter. Then show that  $E(X) = \mu = 2$ .

Actually, one need not compute the mean of a binomial distribution this long way, because that mean is always merely np, where n is the number of trials and p is the probability per trial. Here, np = 4(1/2) = 2, which agrees with the result computed above via the  $\sum_{i} p_i X_i$  formula.

222

How does this result differ from that for example 1, above? If we had wanted to know the average value that would be obtained in repeatedly wanted to know the average value that would be obtained in repeatedly flipping an unbiased coin, where T=0, H=1, and  $p_T=p_H=1/2$ , we would have computed 1/2(0)+1/2(1)=1/2, not the 2 found above for a different problem.

Analogously, if in example 1 we had wanted to know the number of, say, 5's to be expected in rolling an unbiased die 4 times, we would have been working with the binomial expansion (1/6 + 5/6)4; 1/6 is the probability of securing a 5 on any roll of the die. The number 5 can occur 0, 1, 2, 3, or 4 times in 4 rolls of the die. The answer would be

$$p_6(0) + p_1(1) + p_2(2) + p_3(3) + p_6(4) = \frac{1}{6}(4) = 2/3$$


not the 3.5 found in example 1 for another problem.

If we rolled a die 4 times, and then 4 more times, etc., until we had rolled it 6 sets of 4 times each, we would expect a total of (2/3) (6 sets of rolls) = 4 fives to appear. This is reasonable, for we would have rolled the die independently 4(6) = 24 times, each time with a probability of 1/6 of rolling the number five: 24(1/6) = 4.

Obviously, one must state his probability problem carefully and then solve it, rather than a similar-sounding but different problem. This can be quite tricky, as even some eminent mathematicians have regretfully learned.

If X is a continuous variable instead of a discrete one, then an algebraic function describes the form of its probability distribution. As we saw earlier. if X is continuous we cannot assign a probability to a single value of X. Instead, we make statements about the probability that X lies in an interval. For these reasons, the definition given above for E(X) cannot be applied to a continuous random variable. Unfortunately for those who have no knowledge of calculus, there seems to be no sensible way to define the expectation of a continuous variable without recourse to the calculus concept of integration. We shall attempt to sketch heuristically the idea of E(X)when X is continuous so that those without a knowledge of calculus will not be at any disadvantage in our later discussions.

Suppose X is a continuous random variable and the probability distribution of X looks like the one in Fig. 10.10. There is an algebraic rule that gives the height of the curve in Fig. 10.10 for every value of X. The area under the curve is 1 unit. The probability that X will assume a value between, for example, 2 and 3 is equal to the area under the curve between those two points.



Definition: The expectation of the continuous random variable X is the sum of the products formed by multiplying each value that X can assume by the height of the probability function curve above that value of X.

Since X can take on infinitely many values, you might wonder how you could physically multiply each of the separate values of X by the height of the curve at X to find its expectation. This is the problem that recourse to the integral calculus solves. We ask that you take it on faith that it can be done, in a precise but somewhat indirect way, by "integration."

The expectation of a continuous random variable X is denoted by E(X) or  $\mu$ , as is the expectation of a discrete variable.

#### Moments

Moments are quantities that describe the distributions of variables.

Definition: The first moment of a random variable X is  $\mu$ , the expectation of X.

The first moment is also called the "population mean."  $\mu$ , the first moment of X, describes the general location of the distribution along a line. For some distributions (for most met in practice), E(X), or  $\mu$ , is a good indicator of the central point toward which the values of X tend. Suppose X and Y are both normally distributed random variables, but  $E(X) = \mu_p = 10$  and  $E(Y) = \mu_p = 5$ . Then we know that the distribution of X is generally to the right, on the number line, of the distribution of Y, as in Fig. 10.11.



FIG. 10.11 Distributions of X and Y where E(X) = 10 and E(Y) = 5.

**Definition:** The second moment of a random variable X is  $E(X^2)$ .  $E(X^2) = p_1X_1^2 + p_2X_2^2 + \dots + p_nX_n^2$ 

$$E(X^2) = p_1 X_1^2 + p_2 X_2^2 + \dots + p_n X_n^2$$

if X is discrete.

We have little use for the second moment directly. The idea of a "second moment" is used to define a very important concept, however. is the probability that a child entering kindergarten will require special teaching. i.e., will have either perceptual or emotional problems or both?

- 3. a. Find 41
  - h Find 15!/13!.
  - c. Find 61/13! (6 3)11
  - d. For what value of n is (n + 1)! exactly twenty times larger than n!?
  - 4. An experimenter wishes to have subjects learn a list of paired associations in all possible orders of the pairs. There are 10 paired associates and 150 subjects: each subject can learn the list only once. Are there enough subjects so that at least one subject can learn every possible ordering of pairs?
  - 5. a. Find  $\binom{6}{3}$ .
    - b. Find (4)
    - c. Find (15).
  - 6. The varsity basketball team has 13 members. How many possible "starting fives"—the five players who start the game—could the coach form from his team of 13 players?
  - 7. Verify that  $\binom{4}{0}$  +  $\binom{4}{1}$  +  $\binom{4}{2}$  +  $\binom{4}{3}$  +  $\binom{4}{4}$  is equal to 2<sup>4</sup>. (In general,
    - $\sum_{i=1}^{n} \binom{n}{i} = 2^{n}.$
  - 8. In how many ways may a ten-item test be split into two tests of five items each? (Hint: Find the number of ways to select five items from ten to comprise one half. Since selecting five items for one half is the same as not selecting them for the other half, divide the first quantity you calculated by 2.)
  - 9. A pupil takes a 10-item true-false test in which half of the statements are true. If he guesses randomly between "true" and "false" on each item, what is the probability that he will earn a score X of 4 or more correct? (Hint: Find the probabilities of either 0, 1, 2, or 3 items correct and subtract the sum of these probabilities from 1.00.)
  - Graph the expected distribution of total test scores X that would be obtained by 256 completely ignorant examinees who will guess at random the answers to each of four four-option multiple-choice items. (Hint; Regard the test items as four binomial trials in which the probability of success is 1/4. Multiply each term of the binomial distribution with n=4 and p=1/4 by 256 to obtain the number of examinees that would be expected to obtain each test score from 0 to 4.) Stanley (1964, p. 197) shows the distribution of chance scores for 90 five-option items. Compare the skewness of your results with that of his Curve 2.

- 11. It can be proved that if n binomial trials result in nasuccesses, then the probability of this happening is maximal if p, the probability of success on any one trial, equals  $n_0/n$ . (When p is unknown and the outcome of n trials is n, successes. the ratio nin is termed the "maximum likelihood estimate" of the unknown p because this value of p, namely n.ln, makes the outcome more probable than does any other value of p.) Suppose 4 binomial trials yield 3 successes. Verify that this outcome has higher probability if p = 3/4 than if p = 1/2 or 4/5.
  - 12. Ten convicts volunteered for an experiment on the relationship between smoking and lung cancer. The convicts were matched into five matched pairs so that both pair mates are of the same age. Within each pair of convicts a coin was flipped to determine which convict would smoke two packs of cigarettes a day and which one would not smoke for the duration of the experiment. At the end of the 30-year experimental period the five smokers in each pair had lung cancer; none of the nonsmokers had lung cancer. (This experiment is pure fiction.) Suppose that at the outset of the experiment, five convicts either had lung cancer or were destined to develop it in the next 30 years whether they smoked or not. What is the probability that if smoking is truly unrelated to lung cancer, the five initially cancerous convicts were totally by chance assigned to be the five experimental smokers?
    - 13. In the general population, Stanford-Binet IQ's are nearly normally distributed with a mean of 100 and a standard deviation of 16. By referring to Table B in Appendix A, determine the following probabilities to two decimal places:
      - a. That a randomly sampled person will have an IQ between 80 and 120.
      - b. That a randomly sampled person will have an IQ above 140.
      - c. That three independently randomly sampled persons will all have IQ's above 92.
    - 14. The variable X takes on the values 0, 1, 2, 3, and 4 with probabilities 0, 2/5, 1/5, 1/5, and 1/5, respectively. What is the value of E(X), the expected value of X?
    - 15. The variable X is binomially distributed with n = 4 and p = 1/2, i.e., X takes on the following values with the following probabilities:

X	Probability (X)
0	44
1	ie.
2	
3	17
4	11

- a. Determine the value of E(X). Compare this value with np = 4(1/2) = 2.
- b. Determine the variance of X, which is

$$\sigma_x^2 = \sum_{i=0}^4 [X_i - E(X)]^2$$
 probability  $(X_i)$ .

Compare this value with npq = 4(1/2)(1/2) = 1.

16. Trials of an event that can happen in one of three ways are called trinomial. If the event A can occur in three possible ways (a<sub>1</sub>, a<sub>2</sub>, a<sub>3</sub>) with probabilities p<sub>1</sub>, p<sub>2</sub>, and p<sub>3</sub> (p<sub>1</sub> + p<sub>3</sub> + p<sub>3</sub> = 1), respectively, then it can be shown that the probability that n trials of A will produce r occurrences of a<sub>1</sub>, s occurrences of a<sub>2</sub>, and r occurrences of a<sub>2</sub> is given by the general formula for the trinomial distribution:

$$P(r, s, t) = \frac{n!}{r! s! t!} p_1^r p_2^s p_3^t.$$

- a. Show that if  $p_3 = 0$  the formula for the trinomial distribution reduces to the formula for the binomial distribution. (Hint: If  $p_3$  is zero, t can never be anything except zero and s must equal n r.)
- b. The First Methodist Church is planning a Sunday evening pot-luck supper. Each of 20 families is asked to bring either a salad or a "hot dish" or a dessert. Assume that the families make their choice of a contribution to the supper independently and with the following probabilities: for salad,  $p_1 = .10$ ; for "hot dish."  $p_2 = .60$ ; for desert,  $p_3 = .30$ . Calculate the probability that—to the chagtin of the parents and to the delight of the children—each of the 20 families brings a dessert to the pot-luck supper.

## H

# THEORETICAL DISTRIBUTIONS FOR USE IN STATISTICAL INFERENCE

## II.I

Part of the "working tools" of inferential statistical methods are a group of theoretical distributions of some special variables. In this section we shall study four such distributions: the normal, which was the subject of Chapter 6: the chi-square distribution; the I-distribution; and the I-distribution.

#### 11.2 NORMAL DISTRIBUTION

In Chapter 6 we learned that a normal distribution is a particular type of mathematical curve. The graph of a normal distribution is symmetric about its mean,  $\mu_i$  it is unimodal; it has a kurtosis of 3.0, etc.

The normal distribution—in fact there are many, one for every different set of values of  $\mu$  and  $\sigma$ —is very important in statistical inference. Many inferential statistical techniques rest on the assumption that the frequency distribution of scores on a variable in a population is adequately described as a normal distribution with a certain mean and standard deviation. It will be seen in this section that the other theoretical distributions build upon the normal distributions.

This primacy of the normal distribution probably can be attributed to three facts, one empirical, the other two mathematical. First, the normal distribution happens to be a fairly accurate representation of the frequency distributions of scores on a large number of different variables. In Sec. 6.6 we saw that the normal distribution was a good representation of the frequency polygon of heights of 8585 adult men born in Great Britain during the 19th century and of the distribution of scores on the Stanford-Binet Intelligence Test.

Second, it is a mathematical fact that averaging individual scores that are not normally distributed results, in many instances, in an average that is nearly normally distributed. This fact—stated and proved as the "Central Limit Theorem" of mathematical statistics—finds important applications in modern inferential statistical techniques, although it does not play the crucial role it once did. (See Sec. 12.3.)

Third, the normal distribution is mathematically convenient. A most convenient feature of the normal distribution is that when samples are drawn from a normal distribution and  $\mu$  and  $\sigma$  are estimated in a certain way, the two estimates produced for repeated samples are independent.

#### 11.3 CHI-SQUARE DISTRIBUTIONS

Imagine a huge population of scores that are essentially normally distributed amagine a nuge population of 300103 mass are community distributed with mean 0 and standard deviation 1. Suppose that a single score  $X_1$  is selected at random from this population and the standard score z= $(X_1-0)/1$  is formed. Over repeated selections of an X score, z will have a normal distribution with mean 0 and standard deviation 1. Denote the square of z as follows:

 $\gamma_1^2 = z^2,$ 

i.e., the square of a standardized score selected from a normal distribution is symbolized by  $\chi_1^2$ , where the superscript 2 tells us that squaring has taken symbolized by X<sub>1</sub>, where the supersorpe 2 can be shat squaring has taken place and the subscript 1 tells us that only one score has been "combined" to produce  $\chi^2$ . We can conceive of repeating an unlimited number of times to produce  $\chi^2$ . We can expect of  $\chi^2_1$  was found. Each time a new X score is the process by which a value of  $\chi^2_1$  was found. the process by which a value of  $\lambda_1$  and squared. We can build a frequency randomly selected, standardized, and squared. nature into some  $x_1$  statement  $x_2$  and  $x_3$  are polygon of the values of  $x_1^2$  so obtained. If this frequency polygon is smoothed polygon of the values of  $\chi_1^2$  have been recorded and if the scale off after many thousand values of  $\chi_1^2$  have been recorded and if the scale on ancer many thousand values of A1 and the resulting curve is 1, of the ordinate is adjusted so that the area under the resulting curve is 1, of the orthpare is adjusted so that the graph of the chi-square distribution with one degree of freedom would be obtained. A graph of this distribution appears in Fig. 11.1.

The mathematical curve that is graphed in Fig. 11.1 and that describes The maintenantial curve many  $\chi_1^2$  has a complex formula we will not study. (If interested the distribution of  $\chi_1^2$  has a complex formula we will not study. in this formula, see Graybill, 1961, p. 31.) We denote the mathematical




FIG. 11.1 Graph of the chisquare distribution with one degree of freedom. (The shaded area comprises 30% of the area under the curve.)

curve in Fig. 11.1 by  $\chi_1^{1}$ . The curve takes its name from the Greek letter  $\chi$  used to denote it. (The mathematical curve for the chi-square distribution was derived by Karl Pearson in 1900.)

The area under the curve for  $\chi_1^2$  is set equal to one unit so that  $\chi_1^2$  is a probability distribution, e.g., the probability of obtaining a value of  $\chi_1^2$  between 0.5 and 2.5 equals the area under the curve between 0.5 and 2.5. In Fig. 11.1 we see that .30 or 30% of the area under the curve lies to the right of 1.07. Thus, we know that the probability of obtaining a value of  $z^2 = \chi_1^2$  that exceeds 1.07 is .30. In other words, 30% of the z scores randomly selected from a normal distribution will have squares that exceed 1.07. An equivalent statement of this fact is that the 70th percentile in the chi-square




FIG. 11.2 Graphs of χ<sup>\*</sup><sub>1</sub> and χ<sup>\*</sup><sub>2</sub>, the chi-square distributions with two degrees and three degrees of freedom.

distribution with one degree of freedom equals 1.07. We also write this as-

$$r_0 r_1^2 = 1.07$$
.

where  $\chi_1^4$  denotes the chi-square distribution with one degree of freedom and .70 indicates the 70th percentile of that distribution.

Now we shall develop the chi-square distribution with two degrees of freedom,  $\chi_{\tau}^2$ . Suppose we go back to the original normal distribution of X. Instead of drawing out just one X score, draw two X scores at random and independently. Standardize each of these scores by subtracting  $\mu$  from it and dividing the difference by  $\sigma$ . Call the first (by order of selection, not by size) standardized score  $\tau_1$  and the second  $\tau_2$ . Now square and sum the two  $\tau^2$ 's to form the quantity

$$\chi_2^2 = \left(\frac{X_1 - \mu}{2}\right)^2 + \left(\frac{X_2 - \mu}{2}\right)^2 = z_1^2 + z_2^2.$$
 (11.2)

This process of determining a  $\chi_1^2$  could be repeated thousands of times with new pairs of z scores. A frequency polygon of these  $\chi_1^2$  scores could be constructed, smoothed, and reduced so that the area under the curve was one unit. The resulting curve would look like the graph of the mathematical curve  $\chi_1^2$ , the chi-square distribution with two degrees of freedom. Fig. 11.2 shows a graph of  $\chi_1^2$ .

### The Chi-Square Distribution with n Degrees of Freedom, X:

A chi-square variable with n degrees of freedom,  $\chi_n^2$ , is formed by adding together the squares of n independent z scores from a normal distribution:

$$y_{-}^2 \approx z_1^2 + z_2^2 + \ldots + z_n^2$$
 (11.3)

If a large number of these  $\chi_n^*$  values are generated from separate sets of nz scores, their frequency polygon will have the same shape as the mathematical curve  $\chi_n^*$ . Figure 11.3 illustrates the graphs of  $\chi_n^*$  and  $\chi_n^*$ .

The area under each curve in Fig. 11.3 is one unit. One-half the area under  $\chi_{10}^2$  lies above the point 9.34. Hence, we know that the probability is .50 that the sum of the squares of ten z scores drawn at random from a normal distribution will exceed 9.34. Equivalently  $_{10}\chi_{10}^2 = 9.34$ , the median of the chi-square distribution with ten degrees of freedom.

There is a different chi-square distribution for each integer value of n (1, 2, 3, ...). The properties of the curve  $\chi_1^k$  depend upon the value of n. The following facts provide a partial description of the family of chi-square distributions:

 The mean of a chi-square distribution with n degrees of freedom is equal to n. For example, the average value of χ<sup>2</sup><sub>12</sub> one would




FIG. 11.3 Graphs of 2 and 210.

expect to obtain by squaring and summing 12 independent, standardized normal scores is 12.

- 2. The mode of  $\chi_n^2$  is at the point n-2 for n=2 or greater.
- 3. The standard deviation of  $\chi_n^2$  is  $\sqrt{2n}$ .
- The skewness of χ<sup>2</sup><sub>2</sub> is √8[n. Hence, every chi-square distribution is positively skewed, but the asymmetry becomes very slight for large n.
- As n becomes large, χ<sup>2</sup> approaches more nearly a normal distribution with mean n and standard deviation √2n.

An important theorem concerning combinations of chi-square variables can be stated as follows:

If  $\chi^2_{n_1}$  has a chi-square distribution with  $n_1$  degrees of freedom and if  $\chi^2_{n_1}$  has a chi-square distribution with  $n_2$  degrees of freedom and is independent of  $\chi^2_{n_1}$ , then  $\chi^2_{n_1} + \chi^2_{n_2}$  has a chi-square distribution with  $n_1 + n_2$  degrees of freedom.

The pth percentile in the chi-square distribution with n degrees of freedom is denoted by  $\chi_{L_{*}}^{2}$ . The percentiles of the chi-square distribution play a prominent role in inferential statistical techniques, particularly as applied to nominal data. Various percentiles in thi-square distributions for n=1 up to n=30 appear in Table C of Appendix A to this text. The following an example of how Table C is read: Suppose one wishes to find the 50th percentile in the chi-square distribution with four degrees of freedom, i.e.,  $\omega Z_{*}^{2}$ . First, the row labeled 4 in Table C is located. Second, the column headed "50th percentile" is found; it is near the center of the table. At the intersection of the appropriate row and column, the number 3.36 is found. This to the value of  $\omega Z_{*}^{2}$ , the median of the chi-square distribution with four degrees of freedom.

Imagine that a chi-square variate with five degrees of freedom  $\chi_1^2$  is formed from  $z_1^2 + \ldots + z_2^2$ . Now suppose a second, independent chi-square variate with 10 degrees of freedom  $\chi_1^2$  is formed by sampling values from a unit normal distribution, squaring, and summing the 10 squared terms. A variate called an *F-ratio* with 5 and 10 degrees of freedom is formed as follows:

$$F_{5,10} = \frac{\chi_{5}^{2}/5}{\chi_{10}^{2}/10} \,. \tag{11.4}$$


By repeatedly determining values of  $\chi_s^2$  and  $\chi_{10}^2$ , dividing each by its respective degrees of freedom, and forming the ratio F<sub>5,10</sub>, the distribution of  $F_{8,10}$  could be determined. It is known from mathematical statistics that the distribution of  $F_{5,10}$  is an F-distribution with 5 degrees of freedom for the numerator and 10 degrees of freedom for the denominator. The Fdistribution with 5 and 10 degrees of freedom is a positively skewed distribution with mean (10)/(10-2) = 1.25 and a median less than 1. Because of the squaring, only nonnegative values of  $F_{5,10}$  may occur; hence the  $F_{8,10}$ distribution has all of its area to the right of zero. The area under the  $F_{8.10}$ distribution between any two values of  $F_{5,10}$  is equal to the probability of obtaining an F-ratio between these two values. Later in this text selected percentile points in F-distributions will be useful. Certain percentile points in the F-distributions were tabulated long ago for easy reference. We know, for example, that the 90th percentile in the F-distribution with 5 and 10 degrees of freedom, 90F5,10, is equal to 2.52. That is, the probability is .90 of obtaining a value of F<sub>8,16</sub> between 0 and 2.52. <sub>98</sub>F<sub>8,10</sub> equals 3.35, and

There exists a different F-distribution to describe the distribution of each F-ratio with a unique combination of degrees of freedom for the chi-square variates in the numerator and denominator. In general, if two independent chi-square variates, one with  $n_1$  degrees of freedom  $\chi^2_{n_1}$ , and the other with  $n_2$  degrees of freedom  $\chi^2_{n_1}$ , are combined as

$$F_{n_1,n_2} = \frac{\chi_{n_1}^2/n_1}{\chi_{n_2}^2/n_2},$$
 (11.5)

then  $F_{n_1,n_2}$  has an F-distribution with  $n_1$  and  $n_2$  degrees of freedom. Such an F-distribution with  $n_1$  degrees of freedom for the numerator and  $n_2$  degrees of freedom for the denominator has the following properties:

- 1. It is positively skewed.
- 2. It is unimodal.



- 3. It has a median of I or less.
- 4. It has a mean equal to  $n_2/(n_2-2)$  for  $n_2>3$ .

The F-distributions with 4 and 4 degrees of freedom and 4 and 25 degrees of freedom appear in Fig. 11.4.

Percentiles above the 50th in the distribution  $F_{n_1,n_2}$  are related to those percentiles below 50 in the following manner:

$$1-pF_{n_1,n_2} = \frac{1}{pF_{n_2,n_1}}$$
 (11.6)

For example, the fifth percentile in the distribution  $F_{10.5}$  equals

$$\frac{1}{35F_{5,10}} = \frac{1}{3.33} = 0.30.$$

Table E of Appendix A is a tabulation of the upper percentile points in the F-distributions for numerous values of  $n_1$  and  $n_2$ . The columns of Table E correspond to values of  $n_1$ ; the rows of Table E correspond to values of  $n_1$ . Check the table to see if you can verify that the 99th percentile in  $F_{10,20}$  is 3.37, i.e.,  ${}_{10,20} = 3.37$ .

#### 11.5 t-DISTRIBUTIONS

Imagine that a unit normal distribution and a chi-square distribution with 10 degrees of freedom are at hand. One observation is randomly sampled from each distribution: z and  $z_{1}^{1}$ . Next, the following ratio is formed:

$$t_{10} = \frac{z}{\sqrt{z_{10}^2/10}}.$$
 (11.7)

The quantity f<sub>18</sub> in Eq. (11.7) is an observation from the r-distribution with 10 degrees of freedom. In other words, if the process of randomly drawing

one observation each from the unit normal distribution and from  $\chi_{10}^2$  and forming  $t_{10} = z/\sqrt{\chi_{10}^2/10}$  was repeated an infinite number of times, the values of  $t_{10}$  would form a t-distribution with ten degrees of freedom.

The t-distribution with 10 degrees of freedom is described by a symmetric, unimodal curve. The mean of the distribution is 0; the standard deviation is slightly greater than 1, approaching 1 as the degrees of freedom increase. The distribution is somewhat leptokurtic, i.e., it has kurtosis greater than 3; hence, it is more peaked and has more area in the extreme "tails" of the distribution than the normal distribution.

There is not just one t-distribution; as was true of the  $\chi^2$  and F-distributions, there exists a family of t-distributions. There is a different t-distribution for every distinct number of degrees of freedom for the chi-square variable in the denominator of Eq. (11.7). If the denominator of Eq. (11.7) involves a chi-square variable with 5 degrees of freedom, then

$$t_5 = \frac{z}{\sqrt{\chi_5^2/5}}$$

has a *t*-distribution with 5 degrees of freedom. In general, if z has a unit normal distribution and  $z_n^2$  is a chi-square variable with n degrees of freedom and is independent of z, then

$$t_n = \frac{2}{\sqrt{r^2/n}} \tag{11.8}$$

has a t-distribution with n degrees of freedom.

All of the t-distributions are described by symmetric, unimodal curves with a mean of 0. The variance of the t-distribution with n degrees of freedom is n(n-2). They are all slightly leptokurtic. As n becomes larger and larger, the distribution  $t_n$  begins to look more and more like a normal distribution. When n is infinitely large—a theoretical possibility that is empirically impossible—the t-distribution is the same as the normal distribution. The t-distributions with degrees of freedom 1, 5, and 25 appear along with the normal distribution in Fig. 11.5.

In subsequent discussions of statistical inference, selected percentile points in a r-distribution will have to be found. The prin percentile in the t-distribution with 10 degrees of freedom will be denoted by \*10. The most often used percentiles in the t-distributions appear in Table D in Appendix A. There we read, for example, that the 95th percentile in the t-distribution with 10 degrees of freedom, i.e., \*10.16\* is equal to 1.812.

Only the upper-percentile points in the t-distributions appear in Table D. Because of a simple relationship it is unnecessary to tabulate both upper- and lower-percentile points. The symmetry of all t-distributions implies that

$$_{1-p}t_n = -_pt_n, \tag{11.9}$$




FIG. 11.5 The t-distributions with 1, 5, and 25 degrees of freedom and the unit-normal distribution.

i.e., the negative of the pth percentile in the t-distribution with n degrees of freedom equals the (1-p)th percentile in the same distribution. For example,

$$_{.05}I_{10} = 1.812$$
, therefore  $_{.05}I_{10} = -1.812$ ,  $_{.05}I_{20} = 2.528$ , therefore  $_{.01}I_{20} = -2.528$ ,  $_{.00}I_{5} = 1.476$ , therefore  $_{.15}I_{5} = -1.476$ .

# II.6 RELATIONSHIPS AMONG THE NORMAL, 1-, CHI-SQUARE, AND F-DISTRIBUTIONS

The tr. x<sup>4</sup>, and F-distributions are all based on the normal distribution. In each instance, sampling from a normal distribution underlies the new distribution. For example, a chi-square variable is formed by summing squared, unit normal variables; in turn, chi-square variables are combined to form F-variables. In this section, the relationships among the various water than the combined of distributions will be made explicit.

We have already seen that a t-distribution with infinite degrees of freedom is the same as the normal distribution. Suppose we look at the square of the t-variable with n degrees of freedom:

$$I_n^2 = \frac{z^2}{\chi_n^2/n}$$
 (11.10)

In the numerator of Eq. (11.10) we find the square of a unit normal variable  $d\pi L dcd\ by\ 1$ ; in the denominator is an independent chi-square variable




FIG. 11.6 The family of F-distributions and their relationship to the normal, t-, and  $\chi^2$  distributions.

divided by its degrees of freedom. Stated in slightly different form,

$$t_n^2 = \frac{\chi_1^2/1}{\chi_n^2/n}.$$
 (11.11)

We recognize, however, that Eq. (11.11) is an F-variable with 1 and n degrees of freedom. Therefore, the square of a t-variable with n degrees of freedom is an F-variable with 1 and n degrees of freedom.

It is somewhat more difficult to prove another interesting fact, which we shall simply state: Any F-distribution with n degrees of freedom for the numerator and infinite degrees of freedom for the denominator is the same as the 2<sup>nd</sup> distribution divided by the constant n:i.e.,

$$F_{\pi,\infty} = \frac{\chi_n^2}{n}$$
.

All of these facts are depicted in Fig. 11.6. Figure 11.6 is a crossclassification of the family of F-distributions with respect to the degrees of

• The proof depends on the fact that 
$$\lim_{n_3\to\infty}\frac{\chi_{n_2}^2}{n_1}=1$$
.

freedom (1 through  $\infty$ ) of numerator and denominator. Each cell in Fig. 11.6 corresponds to an F-distribution. When a cell happens to coincide with a special case of either the normal, t-, or  $\chi^2$  distributions, the symbol for the F-distribution does not appear; this should not be taken to mean that no such F-distribution exists for that cell, however, for an F-distribution could be shown in every cell.

The pth percentile in the  $\chi_n^2$  distribution is the same as the pth percentile in the  $n(F_{n,m})$  distribution. However, if you square the pth percentile in the t-distribution with n degrees of freedom, you obtain the 2p-1 percentile in the distribution  $F_{1,n}$ . For example, the 95th percentile of the  $t_n$ -distribution is the 2(.95)-1=.90=90th percentile of the  $t_n^2$ -distribution is the 2(.95)-1=.90=90th percentile of the  $t_n^2$ - $F_{1,n}$ -distribution. This is true because 5% of the cases exceed the 95th percentile in  $t_n$  and 5% lie below the 5th percentile. When the t values are squared both the top 5% and the bottom 5% take on a positive sign; hence, 10% of the values in  $F_{1,n}$  exceed the square of 10% of the values in

#### PROBLEMS AND EXERCISES

 Complete the following table by finding the value of the designated percentile in the tables in Appendix A:

Distribution	Degrees of freedom	Percentile	Value of the percentile
. Normal		54th	
). <i>t</i>	20	97.5th	
. Normai		90th	
5. t	120	90th	
c. Chì-square	6	1st	
f. F	4 and 60	99th	
g. Chi-square	15	99.9th	

- Compare the 95th percentile in the F-distribution with 8 and infinite degrees of freedom with the 95th percentile of the chi-square distribution with 8 degrees of freedom divided by 8.
- What is the probability that a randomly selected observation from a chi-square distribution with 21 degrees of freedom will exceed 36.34? (Refer immediately to Table C in Appendix A.)
- 4. Recall that the pth percentile in the F-distribution with n<sub>1</sub> and n<sub>2</sub> degrees of freedom is equal to the reciprocal of the (100 p)th percentile in the F-distribution with n<sub>2</sub> and n<sub>4</sub> degrees of freedom. Find the 5th percentile in the F-distribution with 4 and 8 degrees of freedom.
  - 5. The mean of the F-distribution with  $n_1$  and  $n_2$  degrees of freedom equals  $n_2$   $(n_2-2)$ . When  $n_1=n_2$ , the median of the F-distribution is 1. In which

- direction is the F-distribution with 12 and 12 degrees of freedom skewed? (Report your reasoning.)
- 6. The mean of the chi-square distribution with n degrees of freedom is n. The one and only mode is at n-2 for n greater than 2. We know that the chi-square distribution is positively skewed. Is the median of the chi-square distribution with n degrees of freedom above, below, or equal to n?
- 7. For reasonably large n, the chi-square distribution with n degrees of freedom is nearly a normal distribution. The mean and standard deviation are n and  $\sqrt{2n}$ , respectively, regardless of the size of n. For n = 20, find the 95th percentile in the chi-square distribution from Table C in Appendix A and compare it with the 95th percentile in a normal distribution with mean n = 20 and standard deviation  $\sqrt{2n} = \sqrt{40}$ .
- 8. Prove that the variance of the *t*-distribution with *n* degrees of freedom is n/(n-2). (Hint:  $t_n^2$  is the same as the F-distribution with 1 and *n* degrees of freedom. Since  $E(t_n) = 0$ , and  $\sigma_n^2 = E(t E(t))^2$ , the variance of  $t_n$  is  $E(t_n^2)$ .)

sample from a population. A sample is a part, or subset, of a population. The sample is generally selected in a deliberate fashion from the population no order that the properties of the population can be studied. Theoretically, populations can be either infinitely large or finite in size. The truly infinite populations that come easily to mind are somewhat artificial or conceptual: the collection of all positive numbers, the collection of all possible lengths of a stick, the collection of tosses of two dice which could be made throughout eternity. Almost any interesting population of physical items—as opposed to conceptual possibilities—is finite in size: all persons in the Western Hemisphere, the refrigerators produced in Canada in the last decade, the school districts in the United States of America. A finite population may be extremely large, e.g., the proverbial "grains of sand on earth" or 150!, but if it is conceivable that the process of counting the elements of the population could be completed, then the population is finite. At times, but not often, in inferential statistics it is important to distinguish between finite and infinite populations. However, for the purposes of statistical inference it is generally not necessary to worry about the distinction between finite and infinite populations whenever the size of the population is more than 100 times greater than the sample taken from the population. If the ratio of population size to sample size is larger than 100, the techniques appropriate to making inferences to finite populations and those appropriate for infinite populations give essentially the same results. It is customary to use statistical techniques based on the assumption that infinite populations are being sampled whenever the population is reasonably large (containing several hundred or more elements) and the sample from the population does not constitute an appreciable proportion of the population. It is common to speak of a population as being "virtually infinite" when one means to say that it is huse but finite and that statistical techniques that assume infinite populations will be used on it. We shall not discuss inferential statistical techniques that have been developed for "finite populations," i.e., for small populations or when a sample being studied is more than 1/100, say, of the population. An excellent treatment of these "finite techniques" can be found in William G. Cochran's Sampling Techniques (1963).

Measurements taken on populations of things can be described in the ways we have discussed in the preceding chapters. We can compute means, medians, variances, and percentiles on the data gathered from a population; we might calculate the correlation between height and weight for the population of high-school sophomores in the United States of America. The values of various descriptive measures computed for populations are called parameters. For samples, these same descriptive measures are called statistics. The parameter describes a population in the way a statistic describes a sample. It is customary to denote statistics by Roman letters and parameters by Greek letters. The symbol  $\vec{X}$  stands for the sample mean, and the Greek

letter  $\mu$  stands for the population mean. The sample variance is denoted by  $s^2$ ; the population variance by  $\sigma^2$ .

A statistic computed on a sample can be regarded as estimating a parameter in the population. An estimator is some function of the scores in a sample that produces a value, called the estimate; an estimate gives us some information about a parameter. For example, the sample mean Xis an estimator of the mean or average score in the population. A random sample of 100 eight-year-olds might yield 104.65 for a sample mean on the California Test of Mental Maturity; this value, 104.65, would be an estimate of the mean test score in the population from which the pupils were sampled.

#### 12.2 RANDOM SAMPLING

247

Samples, and estimates calculated from them, give us some idea about the characteristics of the population sampled. There are many different ways in which samples can be chosen from a population. Simple random sampling, which was discussed at length in Sec. 10.6, is just one of many ways, though it happens to be the most widely used. Other sorts of sampling plans can become quite elaborate and lead to complex methods of estimation. We shall restrict our attention in this text to simple random sampling. It is so common to speak of only simple random samples in elementary statistics that we will occasionally use only the word "sample" when we mean a "simple random sample." More sophisticated sampling plans, such as stratified sampling, cluster sampling, and two-stage sampling, are dealt with in more advanced statistical textbooks (Cochran, 1963).

Before a sample will serve well as a basis for making estimates of population parameters, it must be representative of the population. However, this criterion of representativeness presents a problem. How would one know whether a sample is representative of a population unless the characteristics of the population are known? And if the characteristics of the population are known, why does one need a sample with which to estimate them? This quandary is a genuine one. It is resolved when one realizes that random sampling of a population will produce samples which in the long run are representative of the population. It will happen that some random samples from a population will not be very representative of it. A random sampling of 20 from the population of all high-school teachers in California might yield—though it is unlikely—20 female French teachers. Some random samples will not be very representative of the population; others will be. One never knows whether the sample one has chosen is representative. If it is a random sample, the most that can be said is that it is randomly representative of the population in all respects.

An important feature of a random sample is that one can determine what sorts of nonrepresentativeness are to be expected in the long run or in a large group of random samples. This is not possible or feasible with many other types of sampling plan. For example, if one were to choose the first 50 men to walk down the street whose names began with the letter "T," one would not have a random sample of the population of adult Americans. Furthermore, this sample of 50 is nonrepresentative of the population, and it is nonrepresentative in unknown ways and to unknown extents. However, if 50 persons were randomly sampled from the population of adult Americans it would be mathematically possible to answer such questions as "How likely is it that a randomly drawn sample of 50 adult Americans will have a mean height that is more than one inch above the mean height of all adult Americans?"


The process of inferential statistical reasoning involves finding an estimate of a parameter from a sample and then determining how representative such a sample can be expected to be for the purpose of estimating the parameter. It is not surprising, then, that inferential statistics is based on assumptions of random sampling from populations

## 12.3 THE CONCEPT OF A SAMPLING DISTRIBUTION

The statistician assesses the representativeness to be expected from random samples by studying sampling distributions. The concept of a sampling distribution is basic to one entire branch of inferential statistics or estimator calculated on a sample is said to possess a certain sampling distribution. You can imagine the process of choosing sample after sample of size n from a certain population and recording for each sample the value of some estimator, e.g., the sample mean \( \bar{X} \). If this process of drawing a sample from the population were repeated thousands of times, it would be possible to construct a frequency distribution of the thousands of sample means that were obtained. The frequency distribution so constructed would look like the sampling distribution of the mean of samples of size n for the population being sampled. If a frequency polygon were drawn for the data to a scale so that the area under the curve was one unit, the curve would be almost identical to the sampling distribution of the sample mean.

Suppose, for example, that a certain population has several thousand elements and that measurement of any one element will yield a score of 0, 1, 2, ..., 9 with egual probability. Hence, the random variable X takes on any one of the values 0, 1, 2, ..., 9 with probability .10. A probability distribution for the population has the form shown in Fig. 12.1.

One hundred random samples of size n=2 were drawn from the above population. For each sample, the mean was calculated. The first sample contained the digits (3, 2), which yield a mean of (3 + 2)/2 = 2.5. Ninetynine other sample means were calculated, and all 100 means were graphed.



FIG, 12.1 Probability distribution for a population.

The frequency polygon of the 100 sample means appears in Fig. 12.2.

Figure 12.2 gives us some idea of what the actual sampling distribution of X looks like for samples of size 2 from the population in Fig. 12.1. The graph in Fig. 12.2 is an empirical approximation to the sampling distribution of X in this situation.

Generally, the statistician does not have to rely on empirical procedures (such as were reported in Fig. 12.2) to determine the sampling distribution of a statistic. Advanced mathematical techniques can be used to answer such questions as "What is the distribution of the mean of samples of size n from a normal distribution?" or "What is the distribution of the product moment coefficient of correlation of X and Y in samples of size n from a population in which the correlation of X and Y is zero?" Because of the difficult mathematics underlying the derivations of most of the sampling distributions we shall use, we shall simply report the results without proof.

One of the principal theorems of inferential statistics concerns the sampling distribution of the sample mean X. This theorem is called the




FIG. 12.2 Frequency distribution of 100 means of 100 samples (n = 2) from the population in Fig. 12.1,

central limit theorem. Suppose that samples are being drawn from an infinitely large population. The mean of this population is denoted by  $\mu$  and the variance by  $\sigma^2$ . Random samples of size n will be drawn from the population. What will the sampling distribution of the sample mean  $\mathcal{X}$  look like? If n is "sufficiently large" (and it is not possible to be more specific about the size of n), the sample mean will be very nearly normally distributed. Furthermore, the mean of all of the sample means will equal  $\mu$ , the population mean; and the variance of the sample means will equal  $\sigma^2/n$ , where  $\sigma^2$  is the population variance.

The following example will be used to illustrate the central limit theorem. Suppose that the population in Fig. 12.3 has a mean  $\mu$  equal to 15 and a variance  $\sigma^2$  equal to 100. Random samples of size 100 will be drawn from the population in Fig. 12.3. The central limit theorem tells us that the distribution of the means  $\bar{X}$ , of these samples will be nearly normal with a mean of 15 and a variance of  $\sigma_2^2 n = 100/100 = 1$ . The sampling distribution of  $\bar{X}$ , in this instance is depicted in Fig. 12.4.

It may seem incredible to you at first that, regardless of the shape of the population being sampled, the means of "sufficiently large" samples will have a normal distribution. Such is the case, however. This is one of three reasons why the normal distribution is so important in statistics. Just how large n must be before the sampling distribution of X is nearly normal depends on the shape of the population. Samples of size 100 are probably large enough to yield nearly normal sampling distributions of X for most populations one might meet in practice.





FIG. 12.3 Distribution of X for a population.



A proof of the fact that the sampling distribution of X has a mean of  $\mu$  and variance of  $\sigma^2/n$ , where  $\mu$  and  $\sigma^2$  are the mean and variance of the population sampled and n is the sample size, is not too difficult to develop. X is the variable being measured on the population; its mean is  $\mu$  and its variance is  $\sigma^2$ . A random sample of size n has a first element  $X_1$ , a second element  $X_2, \ldots$ , and an nth element  $X_n$ . The ordering of the subscript is not indicative of the size of the score in the sample;  $X_1$  is merely the first score chosen in each sample. Therefore, the collection of all possible  $X_1$ 's, i.e., all first scores chosen in all possible random samples from the population, forms a population with mean  $\mu$  and variance  $\sigma^2$ . In other words,  $X_1$ ,  $X_2, \ldots, X_n$  are each random variables with a mean of  $\mu$  and a variance of  $\sigma^2$ .

The sample mean equals  $(X_1 + X_2 + \ldots + X_n)/n$ . The mean of the sampling distribution of  $\bar{X}$  equals the expected value of  $\bar{X}$ .

$$E(\bar{X}) = E[(X_1 + X_2 + \dots + X_n)/n]$$

$$= \frac{1}{n} E(X_1 + X_2 + \dots + X_n)$$

$$= \frac{1}{n} [E(X_1) + E(X_2) + \dots + E(X_n)].$$
 (12.1)

Now  $X_1$ , the arbitrarily chosen first observation in a sample, has the same distribution over samples as does  $X_2$  or any other  $X_{\mu}$ . Its mean is  $\mu$ and its variance is  $\sigma_x^2$ . Hence, the last term in Eq. (12.1) equals

$$\frac{1}{n}(\mu+\mu+\ldots+\mu)=\frac{1}{n}(n\mu)=\mu.$$

Stated in words, the expected value of X, which is the population mean of the sampling distribution of the sample means, is equal to  $\mu$ , the mean of the population being sampled.

How much will X vary randomly from sample to sample? In other words, if a population (sampling distribution) were formed that consisted of the means of random samples of size n from a population with mean  $\mu$ and variance  $\sigma^2$ , what would the variance of this population of sample means be? You might expect that the variance of the sample means will be smaller than the variance of the population sampled. This will certainly be true if each sample contains two or more observations (i.e., n > 2). Notice that if we chose samples with an n of 1, the sample mean would be the same as the single observation drawn. If we then took repeated samples with n=1 and built a sampling distribution of these "sample means," we would simply be building a sampling distribution of the original population. In this case, the variance of the original population,  $\sigma^z$ , and the variance of the sampling distribution of the "means" of samples of size 1 would be the Suppose we try to find the variance of the means of samples of size n=2 from a population. Let the population variance be  $\sigma^2$ . For each sample,  $X=(X_1+X_2)/2$  is calculated.  $X_1$  and  $X_2$  are arbitrary designations for the first and second observations randomly drawn and are not related to the size of the scores. Consequently, over all random samples,  $X_1$  has variance  $\sigma^2$  and so does  $X_2$ . Because the samples are randomly drawn, there is no relationship between the sizes of the first and second observations in any sample. If we were to construct a scatter diagram for graphing the points  $(X_1, X_2)$  from sample to sample, after hundreds of samples the scatter diagram would show no correlation between  $X_1$  and  $X_2$ . This is depicted in Fig. 12.5.

Clearly, X<sub>1</sub> and X<sub>2</sub> are uncorrelated. Thus, the correlation coefficient and covariance between the first and second observations in a sample over infinitely many random samples from a population is zero.

Now the variance over random samples of  $\overline{X}_1 = (X_1 + X_2)/2$  is denoted as follows:

$$\sigma_{r}^{2} = \sigma_{(x_{1}+x_{2})/2}^{2}$$
 (12.2)

The  $\sigma^2$ 's are simply the expected, long-run average values of  $s^2$ ;  $E[s_2^3] = \sigma_x^2$ . Equation (12.2) shows that the variance of  $\tilde{X}$  is the same as the variance of 1/2 times the sum of  $X_1$  and  $X_2$ . We saw in Sec. 5.8 that the effect on the variance of a variable of multiplying the variable by a constant was to multiply the variance by the square of the constant. Therefore,

$$\sigma_{(x_1+x_2)(1/2)}^2 = (\frac{1}{2})^2 \sigma_{(x_1+x_2)}^2$$
 (12.3)

In Sec. 7.9 we learned that if two variables are uncorrelated then the variance of the sum of the two variables is the sum of their variances. Above we argued that X<sub>1</sub> and X<sub>2</sub> are uncorrelated. Hence,

$$(\frac{1}{2})^2 \sigma_{x_1+x_2}^2 = \frac{1}{4} (\sigma_{x_1}^2 + \sigma_{x_2}^2 + 2\rho_{x_1x_2}\sigma_{x_1}\sigma_{x_2}) = \frac{1}{4} (\sigma_{x_1}^2 + \sigma_{x_2}^2).$$
 (12.4)



FIG. 12.5 Scatter diagram of the relationship between the first  $(X_1)$  and second  $(X_2)$  observations in random samples from a population.

First observation in sample,  $X_1$ 

The variance of  $X_1$  over repeated random samples is  $\sigma_x^2$ , and so is the variance of  $X_2$ . Therefore, we can write Eq. (12.4) as follows:

$$(\frac{1}{4})(\sigma_{x_1}^2 + \sigma_{x_2}^2) = (\frac{1}{4})(2\sigma_x^2) = \sigma_x^2/2.$$
 (12.5)

Equation (12.5) expresses the conclusion of the argument; the variance of the mean of samples of size 2 from a population with variance  $\sigma^2$  is equal to  $\sigma^2/2$ . In this instance, n=2 and  $\sigma_L^2=\sigma^2/2$ . This is no coincidence. It is true in general that for random samples of size n,  $\sigma_L^2=\sigma^2/n$ . Let's explore this general statement further.

If random samples of size n are taken from a population with variance  $\sigma^z$ , then the variance of the mean,  $\bar{X} = (X_1 + X_2 + ... + X_n)/n$ , over samples is given by

$$\sigma_L^2 = \sigma_{(x_1+x_2,...+x_n)/n}^2$$
 (12.6)

The right-hand side of Eq. (12.6) shows  $\sigma_2^2$  to be the variance of (1/n) times the sum of the n uncorrelated variables  $X_1, X_2, \ldots, X_n$ . Therefore,

$$\sigma_{z}^{2} = \left(\frac{1}{n}\right)^{2} \sigma_{(x_{1}+x_{2}+...+x_{n})}^{2}.$$

Each variable  $X_i(i=1,2,\ldots,n)$  has a variance of  $\sigma^*$  and is uncorrelated with the other n-1 variables. Therefore, the variance of the sum of the n uncorrelated variables is the sum of the variances of the variables, because each of the n(n-1)/2 covariances is 0. Thus

$$\left(\frac{1}{n}\right)^3 \sigma_{(x_1+x_2+\ldots+x_n)}^2 = \left(\frac{1}{n}\right)^3 (\sigma_{x_1}^2 + \sigma_{x_2}^2 + \ldots + \sigma_{x_n}^2).$$
 (12.7)

Because each variable has the same variance  $\sigma^2$ , Eq. (12.7) can be written as

$$\left(\frac{1}{n}\right)^{2}(\sigma^{2}+\sigma^{2}+\ldots+\sigma^{2})=\left(\frac{1}{n}\right)^{2}(n\sigma^{2})=\frac{\sigma^{2}}{n}.$$
 (12.8)

A fundamental relationship is expressed in Eq. (12.8). The variance of the means of random samples of size n from a population with variance  $\sigma^2$  is equal to  $\sigma^2$ [n.

The expression o'ln has traditionally been called the rariance error of the mean. The positive square root of Eq. (12.8) is another important expression known as the standard error of the mean,

$$\sigma_{t.} = \sigma/\sqrt{n}$$
. (12.9)

The standard error of the mean, Eq. (12.9), is the standard deviation of the sampling distribution of the means of an infinite number of samples, each of size 7, from a population with variance \(\epsilon^2\). Notice that in Figs. 12.3 and 12.4 the population from which samples were drawn had a standard deviation of 10, and the standard deviation of the sampling distribution of

means of random samples of size 100 from that population was 1. This is consistent with Fq. (12.9).

$$\sigma_{z} = \frac{10}{\sqrt{100}} = 1.$$

The estimation of a population correlation coefficient provides another illustration of the concept of a sampling distribution. Probably the variables "verbal intelligence" X and "reaction time" Y are virtually uncorrelated in the population of all twelve-year-olds in the U.S.A. Imagine for the moment that someone administered the Wechsler Intelligence Scale for Children to measure X and a reaction time test to measure Y to all children age 12 in the country. With these measures, a scatter diagram of the X and Y scores could be constructed and, also, the correlation between X and Y could be calculated. Suppose, further, that the normal bivariate surface (see Sec. 6.6) proved to be an adequate description of the scatter diagram for the X and Y scores. Also, suppose that the value of the product-moment correlation coefficient, which we will denote by the Greek letter a turned out to be zero, i.e.,  $\rho_{-}=0$ . (Since the correlation coefficient describes the population in which we are interested, it is a parameter instead of a sample statistic. We have followed the widely accepted custom of denoting all parameters by Greek letters; statistics are denoted by Roman letters. Hence. ρ is the correlation coefficient in a population, and r is a correlation coefficient in a sample from the population.) A sample of 80 children could be drawn randomly from the population and their X and Y scores observed. For this sample, the value of r,, the sample correlation of "verbal intelligence" and "reaction time," could be computed. The sample of 80 could then be returned to the population. A second sample of 80 could then be drawn at random, ray computed, and the sample returned to the population. This process could be repeated indefinitely; a large number of values of rem for samples of size 80 from a bivariate normal population in which p... equals 0 could thus be accumulated. What would the frequency distribution of the large collection of sample correlation coefficients look like? The statistician can answer this question without going through the actual process of drawing thousands of samples and computing r each time. He can show mathematically that the distribution of these values of r for random samples of size 80 from a bivariate normal population in which  $\rho = 0$  is nearly a normal distribution with mean 0 and variance 1/(n-1) = 1/79. The theoretical sampling distribution of r for random samples of size 80 from a population with a = 0 appears in Fig. 12.6.

The standard deviation of the sampling distribution of r is called the standard error of the correlation coefficient. In this particular case only, it happens to be equal to 0.11. The standard error of r is denoted by  $\sigma_r$ .

Approximately 68% of the samples will yield values of r between -.11 and +.11. About 95% of the samples will yield values of r between -.22




FIG. 12.6 Theoretical sampling distribution of r for random samples of size 80 from a bivariate normal population in which  $\rho$  is zero.

and +.22. Most, but not all, samples will yield a value of r between -.33 and +.33. If you were investigating the relationship between X and Y and obtained a value of r equal to .08 on a random sample of 80 persons from a population for which the value of  $\rho$  was unknown to you, would you be prepared to declare categorically that the unknown value of  $\rho$  was not zero? If you were rational about it, you probably would not. If you will measure the proportion of the area under the curve in Fig. 12.6 to the right of a value of r of .08, you will find that about 23% of the samples of size 80 from a population for which  $\rho = 0$  will produce values of r larger than .08. Note also that 23% of the samples of 80 will give values of r less than -.08—in other words, 46% of the samples of 80 will give values of r less than -.08—in other words, 46% of the believe that an r of .08 on a sample of 80 makes a population value of  $\rho$  equal to zero unlikely. This line of reasoning closely parallels one theory of inferential statistics that will be studied in greater detail later in this chapter.

#### 12.4 PROPERTIES OF ESTIMATORS

In spite of the general importance of statistical estimation, attempts at formalizing the properties of estimators were not pursued seriously until the early twentieth century. The most successful single attempt was the work of Sir Ronald Fisher. The concepts of consistency and relative efficiency that will be met in this section are due to Fisher.

It was pointed out earlier that an estimate is a value of a statistic for a sample which gives information about a population parameter. For example, the sample mean  $\mathcal X$  is an estimator of the population mean  $\mu$ . There is a close analogy between the way in which a sample mean is calculated and the of  $\mathcal X$  as estimating  $\mu$ . However, there are other ways of treating sample data to arrive at a value that estimates  $\mu$ . Why not use the sample median or

the sample mode as an estimate of  $\mu$ ? It is certainly possible to do this; we shall see, however, that by the criteria used in assessing the properties of an estimator, X turns out to be a better estimator of  $\mu$  than either the sample mode.

In this section we shall also be concerned with the properties of estimators of a population median, variance  $(\sigma^2)$ , standard deviation  $(\sigma)$ , and correlation coefficient  $(\rho)$ . What are the different ways in which these parameters can be estimated? Is one estimator to be preferred over all others for estimating a certain parameter, and why? We shall look closely at three properties of estimators.

#### Unbiasedness

An estimator is said to be unbiased for estimating a parameter if the mean of the sampling distribution of the estimator equals the value of the parameter being estimated.

Regardless of the nature of the population being sampled, the sample mean  $\tilde{X}$ , is an unbiased estimator of the population mean  $\mu$ . Notice in Figs. 12.3 and 12.4 that the value of the population mean  $\mu$  is 15 and that the mean of the sampling distribution of  $\tilde{X}$ , is also 15. This example illustrates the unbiasedness of  $\tilde{X}$  as an estimator of  $\mu$ . If samples are drawn randomly from a normal distribution (or any other symmetric distribution), then the sample median is also an unbiased estimator of the population mean  $\mu$ . In other words, the average of the medians of an infinite number of random samples from a normal distribution equals  $\mu$ , the mean of the normal distribution (which is, of course, also its median and its mode).

There are many examples of biased estimators. Suppose we wish to enormal distribution in the population. Imagine that for a particular population  $\rho=.75$ . The mean of the sampling distribution of the sample correlation coefficient r will be less than .75 for any finite sample size. Thus, r is in general a biased estimator of  $\rho$ . If you have already looked back at Fig. 12.6, you might doubt this statement. In Fig. 12.6, we saw that the mean of the sampling distribution of r was 0 for samples of size 80 from a population in which  $\rho=0$ . Consequently, r was then an unbiased estimator of  $\rho$ . It is o happens that r estimates  $\rho$  unbiasedly only when  $\rho=0$ . If  $\rho$  is any other value from -1.00 to +1.00, there will be a bias in r as an estimator of  $\rho$ . (Okin and Pratt, 1958, have derived the unbiased estimator of  $\rho$ . Its calculation on a sample is rather complex. Olkin and Pratt provided tables for finding an unbiased estimate of the population correlation coefficient.)

Though we are formally dealing with the property of unbiasedness of estimators here for the first time in this text, this property influenced the methods used to describe variation in Chapter 5. In Sec. 5.5 we chose to

measure the variation in a sample by the quantity  $s_x^2 = \sum (X_i - X_i)^2 | (n-1)$ . It might have been more natural to measure variability by simply taking the average of the squared deviations around the sample mean, but instead it was decided to place (n-1) and not n in the denominator of  $s_i^2$ . Now we are in a position to elaborate on the motivation of this choice. The quantity  $s_i^2$  is an unbiased estimator of the population variance  $\sigma_i^2$ , whereas  $\sum (X_i - X_i)^2 | n$  is negatively biased as an estimator of  $\sigma_i^2$ . That is,

$$E\left[\sum \frac{(X_i - \bar{X}_i)^2}{n}\right] < \sigma^2,$$

approaching equality only as  $n \to \infty$ .

Suppose that we took random samples from any population with variance  $\sigma^2$  and calculated  $s_s^2$  each time. The average of a huge number of these sample variances would be exactly equal to  $\sigma^2$ . Hence,  $s_s^2$  is an unbiased estimator of  $\sigma^2$ . If  $\sum (X_s - \bar{X}_s)^2/n$  had been calculated on each sample instead, the average of these quantities would have been smaller than  $\sigma^2$ , namely  $[(n-1)/n]\sigma^2$ . Of course, if n was quite large—[00 or more, for example—the difference between  $s_s^2$  and  $\sum (X_s - \bar{X}_s)^2/n$  would be quite small, because the value of (n-1)/n would approach 1. Nonetheless, the latter would contain some small bias as an estimator of  $\sigma^2$ . It is sometimes said that  $\sum (X_s - \bar{X}_s)^2/n$  is "asymptotically unbiased" as an estimate of  $\sigma^2$  because the bias becomes smaller as n increases.

Suppose that one has a normal distribution with mean  $\mu=0$  and variance  $\sigma^2=100$ . If an infinite number of random samples of size n=6 were drawn from the population and both  $s_1^2$  and  $\sum (X_1-X_1)^2/6$  were calculated for each sample, the two sampling distributions in Fig. 12.7 would be obtained.

Notice that the mean of the sampling distribution of  $s_n^2$  is 100, the value of  $\sigma^2$ . This illustrates the unbiasedness of  $s_n^2$  in this instance. The mean of the sampling distribution of  $\sum (X_i - X_i)^2/6$  is equal to 83.33. In this instance, the bias introduced into the estimation of  $\sigma^2$  by using n in place of n-1 in the denominator of the sample variance is sizable—i.e., (n-1)/n=5/6 here.

You are probably wondering how it was determined that the denominator of the sample variance should be n-1 and not n-2, or n-3, or  $n-\frac{1}{2}$ . There are several ways of proving mathematically that  $\frac{1}{2}$  is an unbiased estimator, estimator of  $\sigma^2$ . The algebraic proof is cumbersome, although not as difficult to comprehend as you might imagine. We shall not present the proof here. It can be found in Edwards (1964, pp. 29–36).

The quantity s<sup>2</sup> is an unbiased estimator of σ<sup>3</sup>. Does this imply that sample standard deviation, is an unbiased estimator of σ, the population standard deviation? As a matter of fact, it does not. A nonlinear transformation of an unbiased estimator is not itself an unbiased estimator.

253

FIG. 12.7 Sampling distributions of  $s_s^2$  and  $\sum (X_t - X_t)^2/6$  for random samples of size 6 from a normal distribution with variance  $\sigma^2 = 100$ .

so the sample standard deviation is a biased estimator of the population standard deviation. The amount of bias depends on the shape of the population being sampled. If the population is normal, the mean of the sampling distribution of s is slightly less than or. Specifically,

$$E(s) = \mu_s = \left(\frac{4n - 4}{4n - 3}\right)\sigma. \tag{12.10}$$

If n is fairly large, the bias in s is quite small. Nonetheless, s remains a biased (but consistent) estimator of  $\sigma$ . As  $n \to \infty$ , the expression in parentheses in Eq. (12.10) approaches 1 and the bias disappears.

TABLE 12.1 BIASEDNESS OR UNBIASEDNESS OF VARIOUS ESTIMATORS OF PARAMETERS OF VARIOUS POPULATIONS

Parameter	Nature of population	Estimator	Status of the estimator
μ	Any population	X	Unbiased
μ	Symmetric	Median	Unbiased
įt.	Symmetric and unimodal	Mode	Unbiased
μ	Skewed	Median	Biased
μ	Skewed	Mode	Biased
σ <sup>2</sup>	Any population	S.	Unbiased
σ	Normal	5,	Biased negatively*
Pzs	Bivariate-normal	rev	Biased negatively

For interesting discussion of unbiased estimation of the standard deviation, see Cureton (1968b), Jarrett (1968), Cureton (1968c), and Bolch (1968) in that order.

Table 12.1 presents some parameters, their estimators, and statements that the estimator is biased or unbiased. As you study Table 12.1, remember that several different sample statistics may be used to estimate the same parameter and that whether an estimator is biased or unbiased depends in

part upon the shape of the distribution of measures in the population from which samples are drawn.

#### Consistency

254

A second property of estimators is their consistency. A consistent estimator, even though it may be biased, tends to get closer and closer to the value of the parameter it estimates as the sample size becomes larger and larger. Some estimators that are biased are consistent. For example, the sample standard deviation is a biased but consistent estimator of o. By taking a large sample, s will be close to  $\sigma$  in value; the larger the sample becomes, the closer s gets to  $\sigma$ . This can be seen algebraically if you let n approach infinity in Eq. (12.10). In an inexact sense, the condition of consistency implies that a consistent estimator of a parameter is calculated on a sample in such a way that if the same calculation was performed on the entire population it would yield the value of the parameter. The sample mean is a consistent estimator of  $\mu$  since, if the sample were made as large as the population,  $\bar{X}$  would equal  $\mu$ . (The sample mean is also an unbiased estimator of  $\mu$ .) This requirement of consistency makes good sense. It is difficult to find any estimators that are taken seriously which are not consistent. The few examples of inconsistent estimators that might be presented are quite contrived.

#### Relative Efficiency

The third property of estimators we shall consider is their efficiency. "Efficiency" refers to the precision with which an estimator estimates a parameter; it refers to the variability of the estimate from sample to sample. In a few previous examples we have measured this variability (or efficiency) by taking the variance or standard deviation of the sampling distribution of the statistic. The variance error of the sample mean,  $a_r^2$ , is a measure of the efficiency of  $\chi^2$  as an estimator of  $\mu$ . The variance error of the sample of  $\rho$ .

The variance error of an estimator is one of its most important properties. The cariance error of any statistic is the variance of the sampling distribution of the statistic.

Suppose we wish to estimate the value of the population mean of a particular normal distribution. One way of estimating  $\mu$  is to find the mean X, of a sample of size n. However, the sample median Md is also an unbiased estimator of  $\mu$ . Both are consistent estimator of  $\mu$ . Which is to be preferred? This question could be answered by considering the relative efficiencies of the two estimators. Which estimator of  $\mu$ , the sample mean

or the sample median, varies less from sample to sample? Which has a smaller variance error?

If the variance  $\sigma^2$  of the normal population being sampled is 50 and sample size n is 10, then the variance of  $\tilde{X}$ , over repeated random samples is  $\sigma^2 h = 50/10 = 5$ . What about the variance error of the sample median,  $\sigma^2_{Md}$ ? If thousands and thousands of random samples, each of size n, are drawn from a normal population with mean  $\mu$  and variance  $\sigma^2$ , and the median Md is calculated for each sample, the frequency distribution of these sample medians will be normal with mean  $\mu$  and variance  $(1.57)\sigma^2 h$ . Hence, the variance error of the sample median is  $(1.57)\sigma^2 h$ . Figure 12.8 depicts the sampling distributions of  $\tilde{X}$  and Md for samples of size 10 from a normal distribution with variance 50.




FIG. 12.8 Sampling distributions of the sample mean X, and the sample median Md for random samples of size 10 from a normal population with mean  $\mu=20$  and variance  $\sigma^2=50$ .

The variance error of the sample median in Fig. 12.8 is equal to  $(1.57)\sigma^2/n = (1.57)(5) = 7.85$ . This figure reveals that the sample median will vary more than the sample mean over repeated samples. Note that while only 16% of the sample means will be larger than 25, about 26% of the sample medians will be larger than 25. X, is a more efficient estimator of  $\mu$  than Md. Seeking greater descriptive precision, the statistician defines the efficiency of X, relative to Md as the ratio of their variance errors. In this instance,

relative efficiency = 
$$\frac{\sigma^2/n}{(1.57)\sigma^2/n} = \frac{1}{1.57} = .637 = 63.7\%$$
,

meaning that for normally distributed measures the median is less than two-thirds as efficient as the arithmetic mean, regardless of the magnitude of n.

One interpretation of the coefficient of relative efficiency is that if the median of a sample of 100 observations is used to estimate  $\mu$ , the same degree of precision of estimation could be attained by drawing a sample of 64 observations and computing X.

Statisticians have generally combined the criteria of unbiasedness and efficiency when making their choice of a "best" estimator of a parameter. For example, the sample mean, median, and mode might all be worthy contenders for "best estimator" of  $\mu$  in a normal population. The first

question a statistician would probably ask is "Which ones are unbiased estimators?" All three qualify on this criterion. The next question would probably be "Which one is the most efficient?", i.e., "Which one has the smallest variance error?" The sample mode is least efficient, and we saw that the sample median is less efficient than the sample mean. The sample mean wins. In fact, the sample mean wins relative to any competition. The primary reason that  $\bar{X}$ , is used almost exclusively to estimate the population mean  $\mu$  of any population is that it has a smaller variance error than any other unbiased estimator of  $\mu$ . In this sentence you can see that both the properties of unbiasedness and efficiency are important.

#### 12.5 INTERVAL ESTIMATION

The topics we have discussed thus far in this chapter belong to one branch of the theory of estimation of parameters. That branch is called point estimation because we considered as the estimate of a parameter a single value or number—a point on a line, if you will. For example, a sample of 100 teachers might be drawn from the population of all female elementary-school teachers in the nation for the purpose of estimating the average number of years in service for the population. For the sample,  $\mathcal{X}$ , might equal 3.47 years. The number 3.47 is a single point on the number line and thus provides a point estimate of  $\mu$ , the population mean.

The other type of estimate that is typically used builds on the concept of the point estimate and is called an interval estimate. Interval estimation is a highly useful inferential statistical technique. We shall encounter it repeatedly in the remainder of this text.

An interval estimate of a parameter is a segment on the number line, and the value of the parameter is thought to lie somewhere on that interval. For example, the result of drawing a sample from a population in order to estimate  $\mu$  might be the interval (25.9), 13.65), which probably contains between its bounds—lower bound 25.91 and upper bound 38.65—the value of  $\mu$ . Instead of calculating a single point as an estimate of a parameter, we have now found a whole set of adjacent points, an interval, and one of arises to make interval estimation a difficult notion to comprehend is the way in which one determines exactly how probable it is that the parameter lies of constructing an interval estimate of a parameter in such a way that it limits.

For our first example of interval estimation, let us go back to  $\mathcal{R}$ , as an estimator of  $\mu$  and what the central limit theorem tells us about the sampling




FIG. 12.9 The sampling distribution of X, for random samples of size n from a population.

distribution of X. We have seen that if random samples of size n are drawn from a population with mean  $\mu$  and variance  $\sigma^2$ , the sampling distribution of X will have mean  $\mu$  and variance  $\sigma^2/n$ , and will be nearly normal if n is sufficiently large. The sampling distribution of X is shown in Fig. 12.9.

In Fig. 12.9 the standard deviation of the sampling distribution of X. the standard error of the mean, is  $\sigma l \sqrt{n}$ . Since the distribution is normal. 68% of the observations lie within one standard deviation of  $\mu$ , i.e., 68% of the sample means that would be obtained in repeated random sampling would lie on the interval  $\mu - (\sigma/\sqrt{n})$  to  $\mu + (\sigma/\sqrt{n})$ . Approximately 95% of the means lie on the interval  $\mu - (2\sigma/\sqrt{n})$  to  $\mu + (2\sigma/\sqrt{n})$ , since about 95% of the area under the normal curve lies within two standard deviations of the mean. By referring to a table of the unit normal distribution, we can determine exactly how many standard deviations we must go out from  $\mu$  in each direction to establish an interval that includes 80%, 90%, 99%, or any other percent of the area under the curve. For example, we can see from Table B in Appendix A that 5% of the area under the normal curve lies above a z score of 1.64, and 5% lies below a z score of -1.64. In other words, 90% of the area under the normal curve lies within 1.64 standard deviations on either side of the mean  $\mu$ , i.e., within the range  $\mu \pm 1.64\sigma$ . For Fig. 12.9, this means that 90% of the sample means obtained in repeated random sampling would lie on the interval  $\mu - (1.64\sigma/\sqrt{n})$  and  $\mu + (1.64\sigma/\sqrt{n})$ . Stated in probabilistic terms, the probability is .90 that a randomly drawn sample will have a mean  $\vec{X}$  that is larger than  $\mu - (1.64\sigma/\sqrt{n})$  and smaller than  $\mu + (1.64\sigma/\sqrt{n})$ .

samples which might be drawn from the population, adding  $1.64\sigma/\sqrt{n}$  to X and subtracting  $1.64\sigma/\sqrt{n}$  from X will establish an interval that does not contain  $\mu$  between its lower and upper limits. Such an interval is depicted in Fig. 12.11.

Suppose we now consider the entire collection of means of random samples of size n from the population in question. This is an infinitely large collection and any single element of it is denoted by  $\tilde{X}$ . We can think of constructing an interval around each of these sample means by adding and subtracting 1.64 $\sigma$ / $\sqrt{n}$  from each mean. Now we have an infinite collection of intervals. Ninety percent of these intervals contain  $\mu$  between their lower and upper limits; ten percent do not. What is the probability that a randomly selected interval from this infinite collection of intervals will contain  $\mu$  between its lower and upper limits? Of course this probability is 90. The probability must be .10, then, that an interval randomly selected in this manner will not contain  $\mu$  between its bounds.

The mathematical statement of what we have been discussing is concise, unambiguous, and very revealing. Since  $\vec{X}$  is distributed normally with mean  $\mu$  and variance  $\sigma^2/n$ , the variable  $(\vec{X}, -\mu)/(\sigma/\sqrt{n})$  has a normal distribution with mean 0 and variance 1. We know, then, that

probability 
$$\left(-1.64 < \frac{\bar{X}. - \mu}{\sigma/\sqrt{n}} < 1.64\right) = .90.$$
 (12.11)

Stated in words, the probability that the distance of  $\overline{X}$ , from  $\mu$  in units measured by  $\sigma/\sqrt{n}$  is greater than -1.64 and less than 1.64 is .90. Multiplying the inequality inside the parentheses by  $\sigma/\sqrt{n}$ , subtracting  $\overline{X}$ , throughout the inequality, and then multiplying by -1—which changes the direction of the inequality—changes Eq. (12.11) into the following:

he inequality—changes Eq. (and y)
$$\text{probability}\left\{\left[\overline{X}_{\cdot} - \left(1.64 \frac{\sigma}{\sqrt{n}}\right)\right] < \mu < \left[\overline{X}_{\cdot} + \left(1.64 \frac{\sigma}{\sqrt{n}}\right)\right]\right\} = .90.$$




FIG. 12.11 Illustration of how an interval established around X, may not capture  $\mu$  within its limits.

260

STATISTICAL INFERENCE; ESTIMATION

Thus, the probability is .90 that  $\mu$  is greater than  $\bar{X} = (1.64\sigma/\sqrt{n})$  and less than  $\bar{X} + (1.64\sigma/\sqrt{n})$ .

When an interval estimate of a parameter is constructed so that it has a certain known probability of including the value of the parameter between its limits, the interval is called a confidence interval. The probability that the confidence interval "captures" the parameter between its limits should be used in identifying the interval. In the example developed above, a ".90 confidence interval," or "90% confidence interval" was constructed. The confidence coefficient is the probability that a randomly selected interval from the collection of all possible confidence intervals as defined in the above way (i.e.,  $X \pm z\sigma/\sqrt{n}$ ) will capture the parameter. In the above example, the confidence coefficient we chose to use was equal to .90. Another way of saying this is as follows: "A confidence interval with confidence coefficient .90 was constructed." Finally, we speak of constructing a confidence interval around a sample statistic and on a parameter, because for a given population the parameter assumes just one value. For example,  $X \pm (1.64 \sigma / \sqrt{n})$  is the .90 confidence interval around X, and on  $\mu$ .

Example of the construction and use of a confidence interval. We will now present an example of how interval estimation would be applied. The example will be developed in considerable detail so that more can be learned about the rationale of applying the theory we have developed.

A researcher has set out to determine, as well as he can with his limited resources, the average IQ, as measured by the Wechsler Intelligence Scale for Children (WISC), of the 35,000 fifth-grade pupils in his state. The WISC is an individual intelligence test that yields measures of performance and verbal intelligence and a total IQ that is a combination of both; it must be administered by trained examiners. The available funds for this research will cover 900 test administrations, but no more.

The researcher has good reason to believe that the fifth-grade pupils in his state are as heterogeneous as the pupils used to norm the WISC, but he has some reason to believe that their average score might deviate from that of the norm group. Hence, he is willing to believe that the variance of WISC total IQ's in his state is 225—the same as the variance in the norm

ullet You may have sensed that our argument as to why  $\sigma^z$  is known and  $\mu$  is not known is somewhat artificial. In almost all instances, both are either known or unknown. We assumed here that  $\sigma^a$  is known and  $\mu$  is unknown in order to keep the problem of interval estimation simpler than it would be if both  $\sigma^4$  and  $\mu$  had to be estimated from the same sample. This latter case, in which both  $\sigma^{4}$  and  $\mu$  are unknown, is by far the more realistic situation. The distribution theory that led to the solution of the problem of interval estimation of  $\mu$  when  $\sigma^{\dagger}$  is unknown might well be taken as the dawn of modern inferential statistical methods; this solution was not presented until early in this century. It is due to W. S. Gossett who wrote under the pseudonym of "Student" (1908). We will return to "Student" and his epoch-making research in a later chapter.

Our researcher is in the position of taking a random sample of 900 WISC Total IQ scores from a population of 35,000 scores in which the variance is 225. He will calculate  $\bar{X}$  as an estimate of the unknown  $\mu$ . He wishes to establish a confidence interval around  $\bar{X}$ , and he would like the confidence coefficient for this interval to be .99. With samples as large as 900, one can be confident in this situation that in repeated random samples  $\bar{X}$  is very nearly normally distributed around  $\mu$  with a standard error of  $\sigma/\sqrt{n} = \sqrt{225/900} = 0.5$ . From Table B in Appendix A, it can be determined that 99% of the area under the unit normal curve lies within 2.58 standard deviation units of the mean. The sampling distribution of  $\bar{X}$  for samples of 900 from a population with  $\sigma^2 = 225$  appears in Fig. 12.12, where distance along the baseline is in terms of  $\sigma_x = 0.5$ . Thus, 99% of the area under this curve lies within  $\mu \pm 1.29$  because 2.58(0.5) = 1.29.

Confusion often arises in the interpretation of a confidence interval. For example, some persons believe mistakenly that if the 95% confidence interval on  $\mu$  around a sample mean of 46.25 extends from 36.25 to 56.25, then 95% of the sample means would be expected to fall between 36.25 and 56.25. This is an incorrect interpretation of a confidence interval. If 46.25 happens to lie far (perhaps  $3\sigma_3$ ) above (or below)  $\mu$ —and there is no way of knowing this from the sample—then less than half of the subsequent sample means would fall between 36.25 and 56.25.

It is correct to state that "the probability is .95 that  $\bar{X}_{.} \pm (1.96\sigma/\sqrt{n})$  will span  $\mu_{.}$ " or to write the same in symbols:

prob 
$$\left\{\left[\bar{X}_{\cdot} - \left(1.96 \frac{\sigma}{\sqrt{n}}\right)\right] < \mu < \left[\bar{X}_{\cdot} + \left(1.96 \frac{\sigma}{\sqrt{n}}\right)\right]\right\} = .95.$$
 (12.12)

It is understood that the probability statement refers to the sample space of all intervals that could be formed by computing one interval for each sample.




FIG. 12.12 Sampling distribution of X for random samples of size 900 from a population with unknown mean  $\mu$  and known variance  $\sigma^2 = 225$ .

The mathematical statement in Eq. (12.12) makes perfect sense because it involves a variable  $\bar{X}_i$  about which probability statements can be made. However, after a single sample has been drawn and a single value like 46.25 However, after a incorrect to say that  $46.25 \pm 1.969 \sqrt{n}$  has probability .95 of spanning  $\mu$ . It is not legitimate to write

prob 
$$\left\{ \left[ 46.25 - \left( 1.96 \frac{\sigma}{\sqrt{n}} \right) \right] < \mu < \left[ 46.25 \pm \left( 1.96 \frac{\sigma}{\sqrt{n}} \right) \right] \right\} = .95.$$

The above statement of a probability makes no sense because there is no quantity that varies from sample to sample.

The chances are 99 in 100 that the mean of a random sample will lie less than  $2.8\sigma/\sqrt{n}$  units from  $\mu$ . In this case,  $2.8\sigma/\sqrt{n} = 2.58(0.5) = 1.29$ . The researcher knows this is true even though we do not know the value of  $\mu$ . Consequently, if he conceives of adding and subtracting 1.29 from all sample means he would obtain in a series of random samples, he would have a probability of .99 of capturing the true value of  $\mu$  between the limits of the confidence intervals he so constructed. Suppose now that he draws his singles sample and obtains a mean of 103.72. The .99 confidence interval on  $\mu$  around R is calculated as follows:

$$\vec{X}$$
.  $\pm 2.58 \frac{\sigma}{\sqrt{n}} = 103.72 \pm 1.29 = (103.72 - 1.29, 103.72 + 1.29)$   
= (102.43, 105.01).

It is sometimes said that "the probability that (102.43, 105.01) captures  $\mu$  is .99." This is a confusion and is not true. The value of  $\mu$  either does or does not lie between 102.43 and 105.01. Since the value of  $\mu$  is unknown, we do not know whether it does or does not lie in this interval; we know, however, that only these two possibilities are logical. The interval (102.43, 105.01) is not a sample space (see Sec. 10.2), and it does not make sense to speak of probabilities in connection with it. Probabilities and the confidence coefficient apply to the repeated process of constructing confidence intervals. R. von Mises, one of the two or three persons most responsible for modern, statistical notions of probability, has said (1939, pp. 11-14): "Our probability theory has no relation to questions such as: 'Is there a probability of Germany being again involved in a war with Liberia? ... we may say that in order to apply the theory of probability we must have a practically unlimited sequence of similar observations." If the researcher in the above example were to repeat his actions of drawing samples and constructing confidence intervals on  $\mu$  indefinitely, then 99% of the intervals he would produce would contain the value of  $\mu$ . The sample space to which the probability statement applies is the collection of all possible confidence intervals. But any real-life researcher draws only a few samples, typically just one. Whether or not the confidence interval he constructs captures the parameter he is interested in cannot be known by him. It may and then it may not. All he knows is that he has performed an action that would result, if he were to repeat it thousands of times, in the parameter being captured between the limits of his confidence interval 99% (or 95%, or any other percent he chooses) of the time. If his confidence coefficient is large enough, he cannot help but believe that this particular interval he has just calculated has captured the value of the parameter. (This is one good reason for keeping the confidence coefficient large—90, 95, 99, or even larger—as is typically done.) However, if he is rational, he will acknowledge the slight probability (.01, or .05, or .10, or any other value he chooses) that his belief is mistaken.

Summary of the construction of a confidence interval on  $\mu$  around  $\hat{X}_i$  for large samples when  $\sigma^2$  is known. Samples of size n, sufficiently large to insure the near normality of the sampling distribution of  $\hat{X}_i$ , are drawn from a population with unknown mean  $\mu$  and known variance  $\sigma^2$ . A confidence interval on  $\mu$  is to be constructed around  $\hat{X}_i$ . A confidence coefficient of  $1-\alpha$  is chosen. For example, if a confidence coefficient of 95 is desired,  $\alpha=.05$ . (This rather backward way of denoting the confidence coefficient, namely as  $1-\alpha$ , has a purpose that will become evident after more work in statistical inference. The notation is due partly to historical precedent, however. In Sec. 13.4, the origin of the  $\alpha$  notation will be presented.)

From Table B in Appendix A, the value of the z score above which  $100(\alpha/2)\%$  of the area under the unit normal curve lies is found. Denote this z score by  $z_{1-(a_j)}$ . The  $1-\alpha$  confidence interval around  $\bar{X}_i$  is given by the following formula:

$$\left(\bar{X}_{.} = z_{1-(\alpha/2)} \frac{\sigma}{\sqrt{n}}, \bar{X}_{.} + z_{1-(\alpha/2)} \frac{\sigma}{\sqrt{n}}\right).$$
 (12.13)

An expression equivalent to Eq. (12.13) is the following:

$$\bar{X}_{\cdot} \pm z_{1-(\pi/2)} \frac{\sigma}{\sqrt{n}}$$
 (12.14)

Equations (12.13) and (12.14) will be illustrated on the following data:

- n = 400, σ² = 36, and the confidence coefficient is arbitrarily chosen
  to equal .95; i.e., 1 α = .95, so α = .05. From Table B in
  Appendix A we see that 100(α/2)% = 2.5% of the area under the
  unit normal curve lies above a r score of 1.96. Hence, z<sub>sm</sub> = 1.96.
- 2. The mean of the sample of 400 observations equals 51.04.

The standard deviation of the sampling distribution of r for random samples of size n from a bivariate normal population with correlation coefficient a is

$$\sigma_r = \sqrt{\frac{1-\rho^2}{n-1}} \, \cdot$$

As  $\rho$  increases from 0 to +1, the sampling distribution of r becomes increasingly negatively skewed. As a decreases from 0 to -1, the sampling distribution becomes increasingly positively skewed.

Perhaps you can appreciate the impossible state of affairs that would exist if we tried to use the above sampling distribution information about r to construct confidence intervals on  $\rho$ . If you can, you will surely be impressed by the insightful solution to these problems that the English statistician Sir Ronald A. Fisher (1890-1962) produced in 1921. Fisher determined that a particular mathematical transformation of the sample correlation coefficient would yield a quantity with a sampling distribution that would have the same variance regardless of the value of p. This transformation is called Fisher's Z-transformation. The Z-transformation of any r is denoted by Z. the Z-transformation of the value of  $\rho$  is denoted by  $Z_{\rho}$ ,  $Z_{\tau}$  has the following formula: (12.15)

$$Z_r = \log_r \sqrt{(1+r)/(1-r)}$$
. (12.15)


If we want to find the Z-transformation of the value of  $\rho$ , we simply substitute p for r in Eq. (12.15).

$$Z_{\rho} = \log_{\rho} \sqrt{(1+\rho)/(1-\rho)}.$$

The mathematical symbolism in Eq. (12.15) may look baffling to you. Virtually no one ever computes Z, from Eq. (12.15), but uses a table instead; the calculations have been performed and tabulated for convenience in Table G in Appendix A. In Table G the value of Z, is given for values of r from 0 to  $\pm 1.000$  in steps of .001. If r is negative, simply give Z, a minus sign. Verify that a Z, of .418 corresponds to an r of .395, and that r = -.775 gives a  $Z_r$  of -1.033. The r to Z transformation is graphed in

Suppose we set out to build a sampling distribution of r by taking Fig. 12.13. thousands of random samples each of size n from a bivariate normal population with correlation  $\rho$  and by computing r for each. Instead of building up a frequency distribution of the r's, however, suppose a frequency distribution of the Zr's was built. How would such a frequency distribution look? The sampling distribution of the Z,'s would be nearly normal with mean  $Z_p$  and variance 1/(n-3). The sampling distribution would look like the distribution in Fig. 12.14.

Fisher's Z-transformation provides what is needed for a solution to the problem of placing confidence intervals around r. The standard deviation of



CHAP, 12

FIG. 12.13 Relationship between r and Fisher's Z-transformation of r. (Redrawn from Fig. 3-10 in Julian C. Stanley, Measurement in Today's Schools, 4th Ed., © 1964, by permission of Prentice-Hall, Inc., Englewood Cliffs, N.J.)

 $Z_r$  over repeated random samples is  $1/\sqrt{n} - 3$ , regardless of the value of  $\rho$ . Hence, 90% of the  $Z_r$ 's obtained in repeated random samples will lie within 1.64 standard deviations—a distance of  $1.64(1/\sqrt{n} - 3) - 07 Z_r$ ; 95% of the  $Z_r$ 's will lie within a distance of  $1.96(1/\sqrt{n} - 3)$  of  $Z_p$ ; etc. The distribution of  $Z_r$  is approximately normal regardless of the zize of n. Consequently, if we add and subtract some multiple of  $1/\sqrt{n} - 3$  from  $Z_r$  we will have a




FIG. 12.14 Sampling distribution of Fisher's Z-transformation of r for random samples of size n from a bivariate normal distribution with correlation coefficient p.

767

specified probability of capturing  $Z_a$  within these intervals. The following is another way of conceptualizing the problem. Since Z, is normally distributed with mean  $Z_n$  and standard deviation  $1/\sqrt{n-3}$ , then

$$\frac{Z_r - Z_p}{1/\sqrt{n-3}}$$

is normally distributed with mean 0 and standard deviation 1. Consequently,

prob 
$$\left[ -1.96 < \frac{Z_r - Z_\rho}{1/\sqrt{n-3}} < 1.96 \right] = .95,$$

OΓ

prob 
$$\left[ -2.58 < \frac{Z_r - Z_{\varrho}}{1/\sqrt{n-3}} < 2.58 \right] = .99.$$

Therefore, the interval

$$Z_r - 1.96 \frac{1}{\sqrt{n-3}}, Z_r + 1.96 \frac{1}{\sqrt{n-3}}$$
 (12.16)

captures  $Z_a$  with probability .95; and the interval

$$Z_r - 2.58 \frac{1}{\sqrt{n-3}}, Z_r + 2.58 \frac{1}{\sqrt{n-3}}$$
 (12.17)

captures Z, with probability .99.

To illustrate Eqs. (12.16) and (12.17) suppose that a sample of n = 84observations is drawn at random from what is believed to be a bivariate normal population and that r equals .245. A confidence interval with confidence coefficient .95 is desired.

First, r is transformed into Zr by reference to Table G in Appendix A:

$$Z_{.245} = .250.$$

Second, 1.96 times the standard error of  $Z_r$  is found:

$$1.96 \frac{1}{\sqrt{84-3}} = \frac{1.96}{9} = .218.$$

The 95% confidence interval on  $Z_{\rho}$  is found by substituting into Eq. (12.16). The lower and upper limits of the interval are

The lower and appears
$$Z_r - 1.96 \frac{1}{\sqrt{n-3}} = .250 - .218 = .032$$
and  $Z_r + 1.96 \frac{1}{\sqrt{n-3}} = .250 + .218 = .468$ ,

respectively.

The interval (.032, .468) was generated by a process having a probability of  $M_{\odot}$  of producing an interval that captures the Z-transformation of  $\rho_{\odot}$  we can interpret the confidence interval more intelligibly if everything is transformed back from Fisher Z scores into correlation coefficients. So we read Table G backwards and find the values of r which correspond to Z scores of .032 and .468. An r of .032 corresponds to a Z of .468. Therefore, the 95% confidence interval around an r of .436 extends from .032 to .436. We feel quite confident that the value of  $\rho_{\odot}$  the population correlation coefficient, is between .032 and .436.

#### 12.6 CONCLUSION

Our purpose in this chapter has been to present the theory and a portion of the practice of interval estimation. We have done so in considerable detail, because it is important that you comprehend the rationale of this highly useful inferential statistical technique. In subsequent chapters, you will come across numerous examples of the construction of confidence intervals. In each instance, some of the particulars of the calculation of the confidence interval will be different from those we have presented in this chapter. New theoretical distributions that describe the sampling distributions of some important statistics will have to be described. The solution to the important problem of placing a confidence interval around  $ar{X}$  when  $\sigma^2$  is unknown will be shown. We shall also see how it is possible to place a confidence interval around the difference between two sample estimates, e.g.,  $X_1$  and  $X_2$  (the means of two separate groups of observations on the same variable), for the purpose of estimating the difference between two parameters, e.g.,  $\mu_1$  and μ<sub>2</sub>. (This technique will provide a flexible tool for answering important questions about the superiority of one population over another.) But for almost every example presented in subsequent chapters, the basic rationale is the same. (The only exception to this will be found in Chapter 16.)

In Chapter 13 we introduce a second major branch of the subject of inferential statistics: "hypothesis testing," a statistical inferential methodology secences. We shall find that it is closely related to interval estimation. In though the close relationships are occasionally obscured to the eyes of the student of statistics, interval estimation. In the student of statistics, interval estimation and hypothesis testing are actually two sides of the same coin.

#### PROBLEMS AND EXERCISES

1. By using the table of random digits (Table A in Appendix A), draw a random sample of eight students from the following set of sixteen:

John	Al	Joan	Phil
Mary	Tom	Susan	Paul
	Maurice	Martha	Edith
Alice			Warn
Roh	Barbara	Jack	17 411

- 2. In which of the following pairs of terms do the two terms in the pair stand for exactly the same thing?
  - a. 1; the standard error of  $\vec{X}$ ; 2; the standard deviation of the random-sampling distribution of X.
  - b. 1:  $\sigma^2/n$ ; 2: the standard error of  $\bar{X}$ .
  - c. 1: the variance error of X; 2: the variance of the sampling distribution of
  - d. 1: the population variance  $\sigma^2$ ; 2: n times the variance error of  $\overline{X}$ .
  - e. 1: the mean of the sampling distribution of  $s_z$ ; 2:  $\sigma_z$ .
- 3. In general, the sampling distribution of s2 is positively skewed with a mean of  $\sigma_{\perp}^2$  (see Fig. 12.7). Recall the relationship between the mean and median in a positively skewed distribution. Is the probability of obtaining a value of s2 that exceeds of greater or less than .50?
- 4. A sample of size n is to be drawn from a population with mean 220 and variance 50. Complete the following table by calculating the variance error and standard error of X for various sample sizes:

n	01 7,101	$\sigma_{\tilde{x}_{i}}^{2}$	_	đ
a. 2				
b. 4				
c. 8				
d. 1	6			
e. 3				
f. 6	4			
g. 1	000			

h. 2000

- 5. When samples of size n are drawn from a normal population with mean  $\mu$  and variance  $\sigma^2$ , the variance error of  $\overline{X}$  is  $\sigma^2/n$  but the variance error of the sample median Md is 1.57 $\sigma^2/n$ . Suppose that for a particular problem,  $\mu = 100$ ,  $\sigma^2 = 25$ , and n=25, and X is used to estimate  $\mu$ ; the variance error of X is  $\sigma^2 | n = 1$ 25/25 = 1. If one chose to estimate  $\mu$  with Md, how large a sample would have to be taken so that the variance error of Md,  $\sigma_{Md}^2$ , would also equal 1?
- 6. A sample of size n is to be drawn at random from a population with mean  $\mu$ and variance  $\sigma^2$ . The sample size is sufficiently large that X can be assumed to have a normal sampling distribution. Determine the probabilities with which

X will be between the following pairs of points:

$$a_{i} u + \sigma l \sqrt{n}$$
 and  $u = \sigma l \sqrt{n}$ 

b. 
$$\mu + 1.64 \sigma / \sqrt{n}$$
 and  $\mu - 1.64 \sigma / \sqrt{n}$ 

c. 
$$\mu + 2.58\sigma | \sqrt{n}$$
 and  $\mu = 2.58\sigma | \sqrt{n}$ 

d. 
$$\mu + 0.675\sigma / \sqrt{n}$$
 and  $\mu - 0.675\sigma / \sqrt{n}$ 

$$n + 3al\sqrt{2}$$
 and  $n = 3al\sqrt{n}$ 

- e,  $\mu + 3\sigma / \sqrt{n}$  and  $\mu 3\sigma / \sqrt{n}$
- 7. A sample of 100 persons was randomly drawn from a population with variance 16 but with an unknown mean p. The value of X obtained was 106.75. Construct the 95% confidence interval on µ around X.
  - 8. Construct the 95% and 99% confidence intervals on p for the following cases: a. n = 28, r = +0.36 b. n = 12, r = -0.65 c. n = 300, r = +0.14
  - 9. Which one of the following sample statistics has the smaller variance over random samples of size n ("variance error")?

a. 
$$\bar{X}_1 = \frac{X_1 + \ldots + X_n}{n}$$
 b.  $\bar{X}_2 = \frac{X_1 + \ldots + X_n}{(n+1)}$ 

## STATISTICAL INFERENCE: HYPOTHESIS TESTING

### 13.1 INTRODUCTION

In Chapter 12 the statistical inferential technique known as interval estimation was developed for a few examples in detail. Interval estimation is just one part, though at present probably the most useful and important one, of the body of statistical inferential techniques. In this chapter we shall encounter a second statistical inferential technique: hypothesis testing. Hypothesis testing has become an almost omnipresent feature of empirical research in education and the behavioral sciences, Reported research can be read only with difficulty and partial comprehension if the reader is not trained in the theory and some of the techniques of statistical hypothesis testing. Empirical research is seldom executed in education and the behavioral sciences without the utilization of either interval estimation or, where it is appropriate, hypothesis testing. This state of affairs is partly a reflection of rigid and uncritical standards imposed by some anxious persons who administer research (as in the case of the academic adviser who demanded that his advisee use statistical inferential techniques on data from all 50 states which

exhausted the population of interest) and partly—the greater part, we hope an indication of the genuine utility and necessity for employing such techniques.

The number of new concepts to be introduced and comprehended in connection with hypothesis testing will make the discussion to follow a challenge. Fortunately, some of the concepts of interval estimation also play central roles in hypothesis testing. We shall find that the concepts of random samples, the sampling distributions of statistics, and probability values applying to statements are building blocks for hypothesis testing as well as for interval estimation. Hypothesis testing and interval estimation are carried out with different languages, but we shall see that they usually produce equivalent results or results that are easily converted from one technique to the other. The basic problem, however, remains "How does one infer properties of the population from observation of a sample?"

#### 13.2 SCIENTIFIC AND STATISTICAL HYPOTHESES

The history of statistical hypothesis testing began in the early eighteenth century. The earliest example of a formal statistical hypothesis test appears in a publication dated 1710 and written by John Arbuthnot (1667-1735); it is titled "An Argument for Divine Providence, Taken from the Constant Regularity Observed in the Births of Both Sexes." Noting that for 82 consecutive years the records showed a greater number of males born than females, Arbuthnot argued that the hypothesis that male and female births are equally likely (each with a probability of 1/2) was refuted by these data; for if the probability of a male birth were precisely 1/2, then the probability of 82 consecutive years in which more males than females were born would be infinitesimally small, (1/2)\*2 to be exact. Arbuthnot concluded that the greater proportion of male births than female births was an act of Divine Providence; the sacred institution of monogamy was being maintained since males were more likely to be killed in war or work before reaching adulthood. Arbuthnot's statistics were unimpeachable, but his theology failed to account for polygamous societies. Like Arbuthnot, the modern researcher is much concerned with the probabilistic consequences of various hypotheses.

A researcher is interested in studying the relationship between creativity and anxiety in fifth- and sixth-grade pupils (this entire example is based on a study by Ohnmacht, 1966). His search of the literature has disclosed one group of persons believing that the creative thinker should be less anxious than the uncreative thinker and another group believing that the two characteristics are not related in any way. Our researcher has not yet joined

either camp. He is undecided on this matter and intends to satisfy his curiosity with a small empirical study of his own. First he must decide how he will measure creativity and anxiety. Two tests that appear to have some validity for measuring the two characteristics are found: Getzels and Jackson's test "Uses of Things" is taken as a measure of creativity, and the "Children's Manifest Anxiety Scale" by Castenada, McCandless, and Palermo is taken as the measure of anxiety. There are over 20,000 fifth- and sixth-grade pupils available for study, but the resources available to our researcher make it possible for him to observe only about 200 of them. Being an excellent student of statistics, our researcher plans to draw a random sample of 200 pupils from the population of 20,000 so that he can draw statistical inferential conclusions about the population based on his sample observations. Each of the 200 pupils in the sample will be given the Uses of Things test and the Children's Manifest Anxiety Scale. The sample product-moment correlation coefficient between the measure of creativity and the measure of anxiety will be computed. Our researcher could proceed to establish a confidence interval around this sample r by the techniques outlined in Sec. 12.5. However, he has been trained in a somewhat different school of statistical thought, one in which decisions are paramount; this being so, he proceeds differently.

The decision that the researcher will supposedly make is a decision about the truth or falsity of a statistical hypothesis. There are at least two types of hypotheses it would be well to identify and distinguish: scientific hypotheses and statistical hypotheses. A scientific hypothesis is a suggested solution to a problem. It is an intelligent, informed, and educated guess. A scientific hypothesis is generally stated as a proposition. "It is an empirical proposition in the sense that it is testable by experience; experience is relevant to the question as to whether or not the hypothesis is true. . ." (Braithwaite, 1953). The formulation of a good scientific hypothesis is truly a creative act. On the other hand, a statistical hypothesis is merely a statement about an unknown parameter. For the next few pages we shall use H: (statement) to denote a statistical hypothesis. "H:  $\mu = 125$ " is a statistical hypothesis; it is an assertion that the unknown mean of a particular population is 125. Clearly, such a statement is either true or false. The decision "H:  $\mu = 125$ is false" is an example of the type of decision with which hypothesis testing is concerned. "H:  $\rho = 0$ , where  $\rho$  is the correlation coefficient in a bivariate normal distribution," is another example of a statistical hypothesis. "H:  $\sigma_1^2 = \sigma_2^{2n}$  is a statistical hypothesis stating that the variances of populations 1 and 2 are equal. How would you denote the hypothesis that the means of populations 1, 2, and 3 are all equal to each other?

It is important to distinguish scientific and statistical hypotheses. It is quite possible to test statistical hypotheses about very mundane matters that possess limited generality and not a whit of scientific importance. For

hypothesis tester, he views the statistical inferential problem of reasoning from the sample and r to the population and  $\rho$  as one of making a decision about a hypothesis that asserts  $\rho$  is a particular number. Partly out of habit, partly because of tradition, and partly because it is a sensible choice, this researcher establishes the hypothesis he wishes to test as  $H: \rho = 0$ , i.e., that the correlation between creativity and anxiety in the population is zero. This is his statistical hypothesis. On the basis of the observations he will make on a random sample of 200 pupils from the population, he will decide to accept his hypothesis as true or reject it as false. The techniques he will use to make a decision about the truth of the statistical hypothesis comprise what is called the hypothesis test.

What constitutes a legitimate and rational test of the hypothesis  $H: \rho = 0$ in this situation? Should the researcher compute r for his sample of 200 and decide that H is true if r is zero and decide that H is false if r is not zero? Obviously not: we know too much about the erratic behavior of sample estimates to agree to such a plan. It is quite possible for  $\rho$  to equal 0 in the population and for r to be substantially different from 0 in a sample of 200. In fact, it is not even an impossibility that a sample of 200 from a population with  $\rho = 0$  will yield an r of +1 or -1! It is extremely improbable, but it is certainly possible. This presents a perplexing problem. Even if  $\rho = 0$  in the population, any value of r from -1 to +1 is a "possibility" in a random sample of 200. Consequently, regardless of the value of r for the sample of 200, the researcher cannot with certainty conclude that o is or is not zero. This is an important principle that underlies all tests of statistical hypotheses, and we shall restate it: In testing any statistical hypothesis the researcher's decision that the hypothesis is true or that it is false is never made with certainty; he always runs a risk of making an incorrect decision. The essence of statistical hypothesis testing is that it is a means of controlling and assessing that risk. As we elaborate the example introduced, we shall learn the rationale of controlling and assessing the risk of deciding incorrectly about the truth of a hypothesis.

The next step after stating the hypothesis to be tested is to draw a sample from the population and make the observations that will reflect on the hypothesis. The researcher has drawn a random sample of 200 pupils, measured them on both the anxiety and creativity tests, and correlated the scores. The value of r for the sample was +.09.

The uncertainty in making a decision about  $H: \rho = 0$  arises from the phenomenon of sampling fluctuation, usually called sampling error. This is not a new idea to us in this text. Our entire discussion of inferential statistics has been concerned with the problem of how to handle estimation of a parameter by a sample value that is almost certainly in error to a greater or lesser degree. As before, we attack the inferential problems of hypothesis testing with the concept of a sampling distribution.




FIG. 13.1 Sampling distribution of r for samples of size 200 from a bivariate normal population with  $\rho \approx 0$ .


After stating the hypothesis, one must determine the sampling distribution of the estimator of the parameter about which the hypothesis is made. Furthermore, one finds that sampling distribution that would result if the hypothesis being tested were true. In our example, we must determine the sampling distribution of r for random samples of size 200 from a bivariate normal population in which  $\rho=0$ . Fortunately, this has been done before and the result is a convenient one. The sampling distribution of r for samples of size 200 under these circumstances is nearly normal, with mean 0 and standard deviation  $1/\sqrt{200-1}=.071$ .

Figure 13.1 is the sampling distribution of r when the hypothesis  $H: \rho = 0$  is true. This sampling distribution, when considered in connection with the actual sample r obtained (r = 0.900), will throw light on the possibility that the hypothesis being tested is true.

We have given up any hope of making either of the decisions "H is true" or "H is false" with certainty. Such certainty is not possible without knowing the entire bivariate population. Instead we must ask the question, "Is the decision 'H is true' plausible after observing an r of .09?" An equivalent question is "Is it reasonable to expect a value of r as deviant from zero as .09 when H is true?" If "yes" is the answer to these questions, then the researcher will decide to accept the hypothesis being tested; if "no" is the answer to the questions, H is rejected. But how do we give exact meaning to "plausible" and "reasonable"? This is an arbitrary matter. Suppose that in a certain locale it rains on 90% of the days. Is it "reasonable" to wake up one morning and announce for no reason whatsoever that today it will rain? Yes, it probably is. If on the other hand it rains on only 10% of the days, it is not very reasonable to go around saying it will rain every day. Ninety per cent makes it plausible to predict rain every day; 10% makes it implausible. These are crude definitions of "plausible" and "reasonable," but the arbitrary logic differs little from that which the statistician uses when he says it is unreasonable to assert at random that "today it will rain" if it rains only on 10%, 5%, or 1% of the days.

Suppose we agree that if an event has a probability of .05 of occurring at random it would be unreasonable or implausible to expect it to occur on the single trial of the event which you observe. We could have chosen the probability of .10 or .01 or .001. Our choice of .05 was arbitrary. It begins

FIG. 13.2 Sampling distribution of r for samples of size 200 from a bivariate normal population in which a = .20.



to become reasonable to expect to observe an event on a single trial when the probability of the event increases above .10, say.

If the hypothesis being tested,  $H: \rho = 0$ , is false, then  $\rho$  is either greater or less than 0. If  $\rho$  is not zero, we would expect to obtain a sample r which is somewhat above or somewhat below the major portion of the distribution in Fig. 13.1. For example, if  $\rho$  is .20, the sampling distribution in Fig. 13.2 would result from repeated sampling of the population.

Consequently if  $H: \rho = 0$  is true, the sampling distribution in Fig. 13.1 will hold. If  $H: \rho = 0$  is false, the distribution of r will lie generally higher on the scale, if  $\rho$  is above 0, or generally lower on the scale, if  $\rho$  is below 0, than the distribution in Fig. 13.1. Hence, a very large value of r, c.g., r = .50, is an unlikely event if the hypothesis  $H: \rho = 0$  is true, but it is much more likely to happen if  $\rho$  is above zero, c.g., if  $\rho = .40$  or .59. Similarly, a very small value of r, c.g., r = .60, is unlikely to occur if  $\rho = 0$  and somewhat more likely to occur if  $\rho = 0$  outlikely or irrelations, the values of r that make the truth of  $H: \rho = 0$  unlikely or implausible lie to the right of some point above zero and to the left of some point below zero in Fig. 13.1. Now we want to determine some such points exactly.

The standard deviation  $\sigma_r$  of the distribution in Fig. 13.1 is .071; thus, the probability that a value in the distribution exceeds  $(1.90)\sigma_r = .140$  is .025. The probability that a value is below -.140 is also .025. Therefore, the probability is .05 that a value sampled from the distribution in Fig. 13.1




FIG. 13.3 Sampling distribution in Fig. 13.1 with the extreme 5% (2.5% in each tail) of the area under the curve shaded.

will lie above .140 or below -.140. In less precise terms, it is unlikely to expect a value of r greater than .140 or less than -.140 to be sampled from the distribution in Fig. 13.3.

In repeated random samples of size 200 from a population in which  $\rho=0$ , a value of r above .140 or below -.140 will occur one time in twenty samples on the average, i.e., the probability is .05 that a random sample will produce an r greater than .140 or less than -.140. When we consider jointly the decisions our researcher can make about the hypothesis and the possible outcomes of selecting a sample and computing r, we find that there are four possibilities. Two possible decisions (after observing the data) are reasonable, and the other two are unreasonable, as the accompanying table shows:

The sample produced an r which was

-0.140 and 0.140 Above 0.140 or below -0.140

H was true

The researcher decided that H was false

Between ~0.140 and 0.140	Above U.140 or below -U.140
The researcher's decision was reasonable	The researcher's decision was unreasonable
The researcher's decision was unreasonable	The researcher's decision was reasonable

Let us see what is implied when a researcher decides  $H: \rho = 0$  is true after an r on a sample of 200 is observed that lies either above .140 or below -.140. In the table we have called such a decision in the light of such evidence "unreasonable." Why is it "unreasonable"? The decision is unreasonable because values of r deviating from 0 by more than .140 are not normally to be expected—they will occur in only 5% of all possible random samples—when  $H: \rho = 0$  is true. If one continues to maintain that the hypothesis  $H: \rho = 0$  is true after observing an r of .30 for a sample of 200, he is forced to admit that the event observed, namely an r of .30 or larger, is relatively rare. Naturally we expect to observe likely events, even though unlikely ones do occur. But common sense tells us to expect likely events and not unlikely ones. Consequently, if we are forced to acknowledge an event as unlikely in order to decide that  $H: \rho = 0$  is true, we find it far more reasonable to alter our decision. A more reasonable decision is that  $H: \rho = 0$ is false; for only values of p other than 0 make it reasonable to expect a sample r either below -.140 or above .140.

Consequently, the reasonable action is to decide H: p=0 is true if r for a sample of 200 lies between - .140 and .140, and to decide H: p=0 is false if r lies below - .140 or above .140. We call this our decision rule.

Will our decision rule ever lead us into error? Most certainly, the possibility of making an incorrect decision with this rule does exist. The heart of the problem of hypothesis testing is the formulation of such decision rules and the assessment of the probability that they will lead us into error.




FIG. 13.4 Representation of the area in the sampling distribution of r for samples of 200 when  $\rho = 0$  which lies more than 1.26 standard deviation units from the mean.

Suppose that in fact, but unknown to us,  $\rho$  is exactly 0. We have agreed to decide that  $H: \rho = 0$  is false whenever an r outside the interval -140 to .140 is obtained. In what percentage of an infinite number of random samples of size 200 will an r deviating from 0 by at least .140 be obtained when  $\rho$  is actually 0? In exactly 5% of the samples. Consequently, if  $H: \rho = 0$  is true, the decision rule we adopted would cause us to decide that  $H: \rho = 0$  was false in 5% of the samples—or with probability .05—when in fact the hypothesis was true.

The researcher investigating the relationship between creativity and anxiety found a correlation of .090 between the Uses of Things test and the Children's Manifest Anxiety Scale in a random sample of size 200. This value of r lies 1.26 standard-deviation units (.090/.071 = 1.26) above 0. If H:  $\rho = 0$  is true, our researcher has drawn an r which lies 1.26 standard deviations from the mean of the sampling distribution of r (see Fig. 13.4). How often would one expect to obtain an r for a sample of 200 that lies 1.26 standard deviations above or below the mean of this normal distribution? From Table B in Appendix A we see that 20.8% of the area under a normal curve lies more than 1.26 standard-deviation units from the mean. Therefore, a value of r deviating from 0 by at least .090 is to be expected in over 20% of the samples of size 200 from a population in which  $\rho = 0$ . It is not unlikely, then, to obtain an r of .090 from a population with  $\rho = 0$ . Consequently it would be unreasonable to conclude that H:  $\rho = 0$  was false on the basis of an r of .090 in a sample of 200.

## 13,4 TYPE 1 ERROR, LEVEL OF SIGNIFICANCE, AND CRITICAL REGION

In this section we shall summarize the points made in the preceding section and give the conventional names for many of the concepts presented there. From the discussion in Sec. 13.3 we shall reconstruct four steps:

- Step 1: A hypothesis to be tested is stated. In our example that hypothesis was  $H: \rho = 0$ . It has long been customary to call the hypothesis to be tested the null hypothesis. This convention arises from the fact that statistical hypothesis testing procedures arose within a philosophy of science that conceived of its role as gathering evidence in attempts to nullify hypotheses. We shall not use the term "null hypothesis" until later in this chapter, when it will be necessary to distinguish between different hypotheses.
  - Step 2: Assumptions are made that are necessary for determining the sampling distribution of the statistic that estimates the parameter about which something is hypothesized. The sampling distribution of this statistic is determined for the case in which the hypothesis of step 1 is true.
  - Step 3: A degree of risk of incorrectly concluding on the basis of sample evidence that H is false is adopted. This risk, stated as a probability, is denoted by x and is called the letel of significance of the hypothesis test (or, occasionally, the "size" of the test). This usage has made "significance test" synonymous with "hypothesis test." From the risk adopted, a set of values of the sample statistic is determined that will lead one to decide H is false if the sample yields such a value. This set of values is called the critical region.

For example, in the illustration in Sec. 13.3 it was decided that a risk of .05 was acceptable. In other words, since no decision about  $\rho$  could be made with certainty, it was considered acceptable to make the probability equal to .05 of deciding  $\mathcal{H}: \rho = 0$  was false when in fact it was true. The level of significance  $\alpha$  of the hypothesis test was thus taken to be .05.

By determining the point (.140) above which and the point (..140) below which 2.5% of the rs in repeated samples of size 200 from a population with  $\rho=0$  would fall, we found the two regions of "unlikely rs given a true hypothesis  $H: \rho=0$ ." These two regions, which constitute the most unlikely fs

of the sample r's one could obtain by sampling from a population with p=0, are the critical regions. The critical regions for the illustration in Sec. 13.3 are indicated in Figure 13.3. One portion of the critical region lies from .140 to +1.00, and the second portion lies from .140 to +1.00. A critical region is sometimes called a region of rejection because the occurrence of a sample value that lies in the critical region leads one to reject the hypothesis  $H_1 \circ p = 0$ .

Step 4: A single sample is drawn from the population, the value of a statistic is observed, and a decision about the truth of H is made. This is the final step in the testing of the hypothesis stated in step 1.

The sample data must lead us to make one of two decisions about H: "H is true" or "H is false." The former decision is spoken of as "accepting H," the latter as "rejecting H." From any sample it can never be concluded with certainty that H:  $\rho = 0$  is true or false; the best one can do is to make a decision about H that has a high probability of being true.

If H is true and our sample leads us to accept H, a correct decision is made. If H is true and our sample leads us to reject H, an incorrect decision is made. Such an incorrect decision is called a type I error or error of the first kind. (Later we shall meet a type II error.) A type I error is made when a true hypothesis H is rejected. It is, of course, impossible for a person to know whether his decision to reject H is correct or a type I error. To know this, it is necessary to know whether H is true or false; but if the truth is known about H, there is no need for inferential statistics. At best, one knows the probability—or proportion of times in the long run—of making a correct decision or a type I error.

In the hypothesis test of  $H: \rho = 0$ , a decision process was set up that would cause one to reject H erroneously five times in 100—or with probability .05—in a long series of similar hypothesis-testing situations if H were true. Hence, it is known that if H were true a type 1 error would occur with probability .05 if any sample r outside the interval -.140 to .140 was regarded as evidence that  $H: \rho = 0$  was false. If H is indeed true, what would be the probability of making a correct decision about it, i.e., accepting it? Since under these circumstances 95% of the r's for samples of size 200 will fall between -.140 and .140, the probability of accepting H when it is true is .95.

The size of the probability of a type I error can be controlled. We shall denote the probability of a type I error for any unspecified hypothesis-testing situation by  $\alpha$ . In the test of H:  $\rho=0$ ,  $\alpha$  was set equal to .05. We can make  $\alpha$  equal to such values as .20, .10, .01, .001, or even .125, if we choose.




FIG. 13.5 The critical region for testing the hypothesis that  $\rho = 0$  with an  $\alpha$  of .01(n = 200).

Since  $\alpha$  stands for the probability of making a certain type of incorrect decision, we would prefer to keep it small. It would probably rarely be acceptable to decide to accept or reject H with a plan that would commit a type I error with probability much greater than .10, i.e., with  $\alpha$  greater than .10. It is customary to let  $\alpha$  equal .05, .01, or .001. We shall now see how we could have tested H: p= 0 with an  $\alpha$  of .01.

If  $\rho=0$ , then the distribution of r for samples of size 200 is approximately normal with mean 0 and standard deviation .071, as we saw before. Both "large" and "small" values of r will cause us to question the truth of the assertion that  $\rho=0$ . So that we shall run a risk of exactly .01 of rejecting  $H: \rho=0$  if it is true, we must determine the two numbers between -1 and +1 that are exceeded by only 1% of the sampling distribution of r when  $\rho=0$ . These two halves of the critical region are depicted in Fig. 13.5.

It can be determined from the table of the unit normal distribution that the probability is .005 that a normally distributed variable will lie more than 2.58 standard deviations abore the mean. Similarly, the probability is .005 that a normally distributed variable will lie more than 2.58 standard deviations below the mean. Hence, if we establish the critical region from (2.58)(.071) = .184 to 1.00 and from -1.00 to (-2.58)(.071) = -1.84, we shall have a probability of only .01 of rejecting H: p = 0 if it is true.

It might be instructive to some who read this to see the above argument in its mathematical form. If  $H: \rho = 0$  is true, then

$$\frac{r-0}{\sigma_c} = \frac{r}{.071} \sim N(0,1),$$

where " $\sim N(0, 1)$ " means "is distributed normally, with population mean and population variance 1." Therefore,

prob 
$$\left(-2.58 < \frac{r}{.071} < 2.58\right) = .99$$
.

Multiplying the inequality in parentheses by .071 gives the following expression:

$$prob(-.184 < r < .184) = .99$$

Consequently, if the correlation coefficient for a sample of size 200 from a bivariate normal distribution is above .184 or below -.184, the null hypothesis H:  $\rho = 0$  can be rejected at the .01 level of significance. Suppose that for a sample of size 200, the value of r is .340. The obvious decision can be stated in several equivalent ways:

- 1. "Reject H;  $\rho = 0$  at the .01 level of significance."
- 2. "Reject H:  $\rho = 0$  with an  $\alpha$  of .01."
- 3. "Reject H:  $\rho = 0$  at the 1% level of significance."
- "Reject H: ρ = 0 with a probability of .01 of making a type I error."

The importance of the assumption that the sample comes from a bivariate normal distribution has been investigated by various researchers: Norris and Hjelm (1961), Netzger and Drasgow (1957), Binder (1959), Eurley (1958), LaForge (1958), and Milholland (1958). If you read one of these references, you would be well advised to read them all since no single reference on this topic presents a balanced picture.

#### 13.5 TYPE II ERROR, $\beta$ , AND POWER

Quite literally, so far in this chapter we have related only half the story of statistical hypothesis testing. In this section, we shall present the rest of our account.

The standard technique for testing the hypothesis  $H: \rho = 0$  is to select a level of significance  $\alpha$ , determine the critical values of r or  $z = (r-0)/\sigma_r$  (whichever you wish), draw a sample and compute r, and then accept or reject H. In the previous section, we showed how to measure the probability that H would be rejected when it was in fact true, i.e., the probability of a type I error. It was acknowledged that the decision "H is false" could be incorrect. Now we acknowledge that the decision to accept H, i.e., to conclude that "H is true," could also be incorrect. In other words, we could falsely accept H, e.g., conclude that  $\rho = 0$  when in fact  $\rho = 20$ . The error of accepting a false H is termed an error of the second kind or a type H error. Having acknowledged the possibility of committing a type H error, we now proceed to a discussion of the techniques for measuring the probability of an error of the second kind.

If the hypothesis  $H: \rho = 0$  is false, some other hypothesis—an alternative hypothesis—about the value of  $\rho$  must be true. We shall now make use of the term null hypothesis for the original hypothesis  $\rho = 0$  and the term alternative hypothesis to describe some other hypothesis we could make about the value of  $\rho$ , e.g., that  $\rho = .20$ . Henceforth, the null hypothesis vill be denoted by  $H_0$  and the alternative hypothesis by  $H_1$ . We can seldom specify a single alternative value of the parameter. Generally (though not always) the alternative hypothesis is composite, i.e., it specifies many possible values of the parameter, instead of simple, like the null hypothesis, in which a single value is hypothesized. The following illustrate a simple null hypothesis and a composite alternative:

null hypothesis: 
$$H_0$$
:  $\rho = 0$ ;  
alternative hypothesis:  $H_1$ :  $\rho \neq 0$ .

In the theory of hypothesis testing, it is held that one of two "states of instruct" may exist: either  $H_0$  is true or  $H_1$  is true, and it is agreed that after inspecting a sample, one of two decisions will be reached:  $H_0$  will be accepted (hence,  $H_1$  is rejected). The four possible combinations of these states of nature and decisions are illustrated here along with a description of the validity of the decision.

		State of Ho is true	nature H <sub>1</sub> is true
(Acc Decision R	Reject Ho	Type I error	Correct decision
	(Accept H <sub>1</sub> )	(Probability ≈ a)	(Probability = 1 ~ $\beta$ )
		Correct decision	Type II error
	(Accept H <sub>0</sub> )	(Probability = 1-a)	(Probability=3)

We have adopted the convention of calling the probability of committing a type I error a. The probability of committing a type II error, i.e., of accepting H, when H, is true, will be denoted by  $\beta$ . We will now look at an example of how  $\beta$  would be calculated.

Suppose that an investigator wishes to test the null hypothesis  $H_0\colon \rho=0$ . Also assume that the investigator has no particular reason to suppose that  $\rho$  is any more likely to be one nonzero value than any other. He can afford to draw a sample of n=200 persons and it is reasonable to assume that the variables observed have a bivariate normal distribution. Furthermore, he wishes to risk rejecting  $H_0\colon \rho=0$  when it is true only five times per 100; hence,  $\alpha=.05$ .




FIG. 13.6 Illustration of the power of the test of  $H_0$ :  $\rho = 0$  against  $H_1$ :  $\rho \neq 0$  for the case in which  $\rho = .20$  (n = 200 and  $\alpha = .05$ ).

The investigator would establish critical regions of r that lead to the rejection of  $H_0$  as was done in Fig. 13.3. A sample r of .140 or larger or -.140 or smaller would be taken as evidence of the falsity of  $H_0$ . We know, then, that if  $\rho$  is actually zero, there is one chance in twenty ( $\alpha = .05$ ) that the investigator will reject  $H_0$ .

However, what if  $\rho$  is really .20? In this case,  $H_0: \rho = 0$  should be rejected in favor of the conclusion that  $\rho$  is different from zero. But what is the probability that  $H_0$  will be rejected? This probability is the power of the test  $\rho = .20$ , and is depicted by the shaded area in Fig. 13.6.

The upper critical region for r is all values from .140 to 1.00. Hence, the power of the hypothesis test to reject  $H_0$  when  $\rho = .20$  is the area above .140 under the curve that represents the sampling distribution of r for samples of size 200 when p = .20. This area is approximately 82% of the total area under the curve on the right in Fig. 13.6. Thus the power is approximately .82. [Actually there also exists an infinitesimal chance that Ho will be rejected in favor of  $H_1$  when  $\rho = .20$  due to a sample r below -.140! We disregard this improbability here; however, see Kaiser (1960).] The area under the curve on the right in Fig. 13.6 (the sampling distribution of r when  $\rho = .20$ ) below .140 is a measure of the probability that r will fail to exceed the critical value even though  $H_0$  is false; this area measures  $\beta$ , the probability of a type II error. The area in question is about 18% of the total area under the curve. Hence,  $\beta$  is approximately .18. Since if  $\rho$  is actually .20 we must either commit a type II error or not, the probability of not committing the error, i.e., the power of the test, is given by  $1 - \beta = .82$ . Now try to convince yourself that if  $\rho$  were equal to -.20, the same hypothesis-testing procedure would run the same risk of a type II error and have the same power. .82, as when  $\rho = .20$ .

It will further extend the notions being developed here if we determine the power of the test of  $H_0$ : p=0 with  $\alpha=.05$  and n=200 when p=10 instead of 1.20. The critical regions of the test remain the same: -1.00 to -1.40 and 1.40 to 1.00. The sampling distribution of r for samples of size 200 is unchanged from previous discussions, and it appears along with the sampling distribution of r for n=200 when p=.10 in Fig. 13.7.




FIG. 13.7 Illustration of the power of the test of  $H_0$ :  $\rho = 0$  against  $H_1$ :  $\rho \neq 0$  for the case in which  $\rho = .10$  (n = 200 and  $\alpha = .05$ ).

Suppose one had chosen to test  $H_0\colon \rho=0$  against  $H_1\colon \rho\neq 0$  with an  $\alpha$  of .10 and a sample of 200 paired observations. By referring to Fig. 13.6, try to determine whether the power of this test is greater or less than the test with  $\alpha=.05$  when  $\rho=.20$ .

From an exact measure of the area under the curve on the right above the critical value of .140 in Fig. 13.7 it can be shown that the power of the test of  $H_0$ :  $\rho=0$  when  $\rho=.10$  is .29. Of course it follows that  $\beta=.71$ .

It seems almost never to be the case in research in education and psychology that the power of a hypothesis test for just one or two alternative values of the parameter is sufficient information. Generally one would want to determine the power of the test for several alternative values of the parameter, graph these values of the power against the values of the parameter, and then connect the points with a smooth line. The resulting curve provides an adequate determination of the power of the test against all alternative values of the parameter. The power curve for the test of  $H_0: \rho = 0$  against  $H_1: \rho \neq 0$  for n=200 and  $\alpha=.05$  appears as Fig. 13.8.

Note in Fig. 13.8 that the values of the power for  $\rho=\pm .20$  and  $\rho=\pm .10$  are .82 and .29, respectively, as they were calculated to be in Figs. 13.6 and 13.7. Notice also that the "power" is .05 when  $\rho=0$ ; that is, if  $H_0$  is true, there is a probability equal to the level of significance of rejecting

It is apparent in Fig. 13.8 that the power of the test increases toward 1 as the value of  $\rho$  departs from 0. This is comforting to know, but it is a contingency not under the control of the investigator since he does not "set" the true value of  $\rho$ . However, the sample size n and the level of significance  $\alpha$  are under his control, relatively speaking at least. For any given value increased (e.g., from 10 to 100) and also increases as n is increased (from .01 to .05, say).

The following can be said about hypothesis-testing procedures in general:

For a given value of the parameter being tested, e.g., p = .40, the
power of the test of H<sub>0</sub> increases as n, the sample size, increases.

2. For a given value of the parameter being tested, e.g.,  $\rho = .40$ , the power of the test of  $H_0$  increases as  $\alpha$ , the probability of rejecting a true null hypothesis, increases, e.g., from .01 to .05.

These two relationships are quite important since to some extent  $\alpha$  and n can be controlled by the investigator. It might be advisable in some circumstances to run a risk of a type I error as large as .10, i.e.,  $\alpha = .10$ , to insure a reasonable power for a test. The third relationship we shall state is much less under the control of the investigator:

3. For fixed values of  $\alpha$  and n, the power of the test of  $H_0$  increases as the true value of the parameter being tested deviates further from the value hypothesized for it in  $H_0$ . For example, if n=100 and  $\alpha=.01$ , the power of the test of  $H_0$ :  $\rho=0$  is greater when  $\rho$  actually equals. 40 or -40.

The popular notion among practicing researchers is that the statistician is a man who tells them "how large a sample to take." Presumably he derives this decision about sample size from studying cost per observation, costs of committing type I and type II errors, and the power of the test for different sample sizes and particular alternative values of the parameter about which a hypothesis is to be tested. The theory—known as the Neyman-Pearson hypothesis-testing theory—that gave us the notions of type II errors and power is very accommodating when these costs and specific alternative values of the parameter can be specified. However, in research in education and the social sciences it is rare that they can be specified with any confidence. We suspect that most statisticians consulting with researchers in these disciplines have had experiences similar to ours. We usually find ourselves advising




FIG. 13.8 Power curve for the test of  $H_0$ :  $\rho = 0$  against  $H_1$ :  $\rho \neq 0$  for n = 200 and  $\alpha = .05$ .

persons to draw the largest sample they can afford to take; then we determine whether the sample size they tell us they are capable of taking is unnecessarily large! If they can take a sample so large that the power of their test of  $H_0$  is .97, say, even when the true value of the parameter is only slightly different from the value specified in Ho, then they can be safely advised not to draw such a wastefully large sample. It may well be true that the power of their test would drop to only .90 if they took a sample half as large as the largest possible sample. If so, we would not hesitate to advise them to reduce the size of their sample. Thus, as our inquiries in education and the social sciences are presently constituted, the concept of the power of a test is more useful as a signal that too large a sample may be drawn than it is useful as the "determiner of sample size."

### 13.6 NONDIRECTIONAL AND DIRECTIONAL ALTERNATIVES: "TWO-TAILED VS. ONE-TAILED TESTS"

An alternative hypothesis,  $H_1$ , can be designated as either nondirectional or directional. The alternative  $H_1$ :  $\rho \neq 0$  is nondirectional in that it states only that  $\rho$  is not equal to zero and not in which direction (above or below) it deviates from zero. Consider the pair of hypotheses  $H_0$ :  $\rho = 0$  and  $H_1$ :  $\rho > 0$ 0. In this instance, the alternative hypothesis is directional; it is maintained by the investigator stating this pair of hypotheses that either ho equals zero to the investigator stating time pair or hypotheses that either  $\rho$  equals zero or else it is greater than zero. The investigator believes that  $\rho$  could not possibly be less than zero; he will gather evidence that will either convince him  $\rho$  is positive or that will lead him to continue to believe  $\rho$  is zero.

One consequence of stating the directional alternative  $H_1$ : ho > 0 to the null hypothesis is that now the critical region for rejection of  $H_0$  in favor of  $H_1$  is that value of r exceeded by 100(a)% of the area in the sampling distribution of r when  $\rho = 0$ . In other words, only large positive values of r will lead one to decide in favor of H, over Ho; hence the critical region for rejection of II. is in the right-hand tail of the sampling distribution of r for ρ = 0, as indicated in Fig. 13.9. In Fig. 13.9 the critical region extends from

A very small value of r, say  $\sim$  .40, certainly does not favor the hypothesis that  $\rho > 0$  over the hypothesis that  $\rho = 0$ . Since only the two conditions  $\rho \approx 0$  or  $\rho > 0$  are covered by the hypotheses, an  $\rho$  of -40 would imply the truth of  $H_0$  more than of  $H_1$ , even though it would be an extremely unlikely occurrence in a sample of n = 200 if p were zero. But such are the vagaries of directional hypothesis tests.

The fact that the critical region "lies in one tail" of the sampling dis-




FIG. 13.9 Illustration of critical region for testing  $H_0$ :  $\rho > 0$  against  $H_1$ :  $\rho > 0$  for n = 200 and  $\alpha = .05$ .

tribution of the statistic under the null hypothesis has made popular the phrase one-tailed test for a significance test of a directional hypothesis. This usage is somewhat ambiguous and potentially misleading. The important distinction is between nondirectional and directional alternative hypotheses; whether a test statistic has grown out of the history of statistics in such a manner that one or two tails of a sampling distribution lie above the critical values of a statistic is quite arbitrary. We shall see in Chapter 15, for example, that a nondirectional hypothesis about a set of population means is tested by referring a test statistic to one tail of the F-distribution.

One sometimes reads that the hypotheses tested in a directional situation are  $H_0$ :  $\rho < 0$  against  $H_1$ :  $\rho > 0$ . It is then maintained that sufficiently large sample r's favor  $H_1$ , and small or large negative r's favor  $H_2$ . Actually, logic does not support this particular statement of  $H_0$  and  $H_1$ .  $H_0$  must assert that  $\rho = 0$ , not that  $\rho < 0$ . Anything else leads to some interesting inconsistencies. Note that  $H_0$ :  $\rho > 0$  and  $H_1$ :  $\rho < 0$  are every bit as acceptable as  $H_0$ :  $\rho < 0$  and  $H_1$ :  $\rho > 0$ . However, for fixed n and  $\alpha$ , the probability of deciding in favor of  $H_0$ :  $\rho > 0$  when  $\rho$  is actually .10 is far greater than the probability of deciding in favor of  $H_1$ :  $\rho < 0$ . It is indeed an unacceptable state of affairs when the power of a test to decide that  $\rho$  is above zero when  $\rho$  is .10 depends greatly on the arbitrary choice between  $H_0$ :  $\rho > 0$ ,  $H_1$ :  $\rho < 0$ ,  $H_1$ :  $\rho < 0$ ,  $H_2$ :  $\rho < 0$ ,  $H_1$ :  $\rho < 0$ ,  $H_2$ :  $\rho < 0$ ,  $H_3$ :  $\rho < 0$ ,  $H_3$ :  $\rho < 0$ ,  $H_4$ 

A debate over the merits of testing directional versus nondirectional hypotheses in research in the behavioral sciences raged for a few years in the 1950's and 1960's. We shall not take the space to recapitulate the issues here. You will find the major issues presented in the following references: Burke (1953, 1954); Goldfried (1959); Hick (1952); Jones (1952, 1954); Kimmel (1957): Marks (1951, 1953); and Peizer (1967).

Our impression is that the potential for misusing directional hypothesis tests is great. To be perfectly legitimate, for example, one who hypothesizes that  $\rho=0$  against  $\rho>0$  must look the other way and refuse to budge from the belief that  $\rho$  is zero even if a sample of 1000 yields an r of -.99. We do not trust ourselves to be quite that orthodox, and we would not be astonished to observe others yielding to temptation.

#### 13.7 POSTSCRIPT

Among recent attempts to clear away the practicing researcher's confusions about statistical hypothesis testing, the most successful in our opinion is William Kruskal's contribution to the International Encyclopedia of the Social Sciences (1968) entitled "Tests of Significance." You could probably profitably read Kruskal's article when you have finished studying this chapter and Chapter 14. Also see Nunnally (1960), Grant (1962), and Wilson and Miller (1964).

### PROBLEMS AND EXERCISES

- 1. Give definitions of each of the following:
  - Null hypothesis, H<sub>0</sub>.
  - b. Alternative hypothesis, H1.
  - c. Type I error.
    - d. Type II error.
      e. Level of significance, a.
  - f. Power of a test,  $I = \beta$ .
  - g. Critical region.
  - a -----
- In which of the following instances can H<sub>0</sub>: p = 0 be rejected with certainty, i.e., in which instances is the reported sample r an impossibility given that p is zero?
   n = 10, r = .60
  - b. n = 100, r = 1.00
  - c. n = 1000, r = .50
  - 3. In each of the following instances, indicate whether a type I error, a type II

    error, or no error was committed by the researcher:

    H<sub>0</sub> H<sub>1</sub> True rates of a new type II

	True taue of p	Researcher's decision based on t
a. p = 0 p ≠ 0 b p = 0 p ≠ 0 c p + 0 p ≠ 0 d p = 0 p ≠ 0 4. The hypothese #	0 .40 0 50	Reject $H_0$ Reject $H_0$ Do not reject $H_0$ Do not reject $H_0$

- 4. The hypotheses II<sub>s</sub>: p = 0 and II<sub>s</sub>: p ≠ 0 were tested with a sample of n = 50 at the .01 leaf of significance. The sample p was sufficiently large that II<sub>s</sub> was repected in favor of II<sub>s</sub>. What is the probability that a type II error was committed? (Hint: Can one commit a type II error when one rejects II<sub>s</sub>?)
- A recearcher draws a sample of n = 200 paired observations from a bivariate normal distribution. He reasons correctly that if p = 0, then r will be distributed

approximately normally with a mean of zero and a standard deviation of .071. Further, he decides to reject  $H_0: p = 0$  if r is above .10 or below -.10. What is the probability that he will commit a type I error? (Hint: What percent of the area under a normal curve with mean zero and standard deviation .071 lies above .10 and below -.107)

- 6. In each of the following instances indicate whether the critical region for rejection of H<sub>θ</sub> lies in the upper (right) tail, lower (left) tail, or is divided between both tails of the sampling distribution of r for ρ = 0;
  - a.  $H_0$ :  $\rho = 0$ ,  $H_1$ :  $\rho \neq 0$
  - b.  $H_0$ :  $\rho = 0$ ,  $H_1$ :  $\rho > 0$
  - c.  $H_0$ :  $\rho = 0$ ,  $H_1$ :  $\rho < 0$
- 7. Researcher Rowe is testing H<sub>0</sub>: ρ = 0 at the α = .05 level with a sample of size n = 25, and he sets his critical values of r appropriately. Researcher Null is testing H<sub>0</sub>: ρ = 0 at the α = .05 level with a sample of n = 100 and he sets his critical values of r appropriately.
  - a. Does Rowe or Null have the larger probability of committing a type I error, or is this probability the same for both?
  - b. If ρ is actually .10, does Rowe or Null have the larger probability, β, of committing a type II error?
     c. Which researcher is performing a significance test that has greater power to
  - c. Which researcher is performing a significance test that has greater power to reject H<sub>0</sub> if ρ = -.20?
- 8. From Fig. 13.8, estimate the power of the test of  $H_0$ :  $\rho=0$  against  $H_1$ :  $\rho\neq 0$  for n=200 and  $\alpha=.05$  when:
  - a.  $\rho = .05$  b.  $\rho = .25$  c.  $\rho = -.25$  d.  $\rho = .40$
- 9. Refer to Fig. 13.7. Suppose that in testing H<sub>0</sub>: p = 0 against H<sub>1</sub>: p ≠ 0 with a sample of r = 200, the investigator sets his critical values at r = -30 and below and r = 3.0 and above. Thus he is adopting an extremely small a. If p is actually 1.0, approximately how large is the probability that he will commit a type II error, i.e., accept H<sub>0</sub> even though it is false?

# SELECTED INFERENTIAL TECHNIQUES

#### 14.1 INTRODUCTION

At one time or another someone has studied the inferential properties of almost every statistic known. Using either strictly mathematical techniques or, occasionally, empirical methods, statisticians have derived sampling distributions to fill most practical needs. In this chapter, the inferential properties of only the more frequently used statistics will be presented; thus the word "selected" in the title of this chapter is appropriate. Where possible, the techniques of both testing the significance of a statistic and constructing a confidence interval around it will be given. The statistics with which we shall deal fall into four classes: means, variances, correlation coefficients, and frequency data.

The discussion of the inferential properties of each statistic will take the following form: (a) statement of the null hypothesis  $H_0$  and the alternative hypothesis  $H_1$ —the alternative hypothesis will always be "nondirectional" so that modifications of critical values are necessary in the event a "one-sided" test is desired; (b) statement of the assumptions made in making the test;

١

(c) identification of the sample statistic employed in testing  $H_0$  and  $H_1$ ; (d) statement of the sampling distribution of the test statistic under both  $H_0$  and  $H_1$ ; (e) determination of critical values of the test; (f) construction of confidence intervals around the sample statistic; (g) an illustration; (h) special considerations, if any.

The first class of hypotheses we shall consider deals with population means.

### 14.2 INFERENCES ABOUT THE MEAN, μ, OF A POPULATION

a. The hypothesis to be tested is that the mean  $\mu$  of a population is equal to some real number a. The alternative hypothesis is, of course, that  $\mu$  is different from a:

$$H_0$$
:  $\mu = a$   
 $H_1$ :  $\mu \neq a$ 

b. It is assumed that the variable X has a normal distribution in the population sampled. One need not know the value of  $\sigma_2^2$ .

c. He is tested by means of the test statistic

$$t = \frac{\overline{X} - a}{s / \sqrt{n}},\tag{14.1}$$

where

$$s_x = \sqrt{\frac{\sum (X_i - \overline{X})^2}{n-1}}.$$

d. If  $H_0$ :  $\mu=a$  is true, t in Eq. (14.1) has the Student's t-distribution with n-1 if (degrees of freedom). When  $H_1$  is true, i.e., when  $\mu$  is actually equal to some value b that is different from a, then the sampling distribution of t in Eq. (14.1) has the shape and variability of Student's t-distribution with n-1 if but it has a mean approximately equal to  $(b-a)/(a_s/\sqrt{n})$ . For example, the distribution of  $t=(\bar{X}_1-0)/(s_s/\sqrt{n})$  is depicted in Fig. 14.1 both for the case in which  $\mu=0$  and  $\mu=2$ ; n and  $\sigma_z^2$  are both equal to 30.

e. The critical values for testing  $H_0$  at the  $\alpha$ -level of significance with the test statistic  $t = (\overline{X}, -a)/(s_d/\sqrt{n})$  are the  $100[1 - (\alpha/2)]$  percentile point in Student's t-distribution with n-1 df and the negative of this percentile point:

$$-1-(2/2)t_{n-1}$$
 and  $1-(2/2)t_{n-1}$ .




FIG. 14.1 Sampling distributions of  $t = (\bar{X}_1 - 0)/(s_s/\sqrt{n})$  for the case in which  $H_0: \mu = 0$  is true and the case in which  $H_1: \mu = 2$  is true. The values of n and  $\sigma_n^2$  are both 30 ( $\alpha = .05$ ).

A value of  $t=(\vec{X}_1-a)/(s_n/\sqrt{n})$  falling below the negative critical value or above the positive critical value constitutes evidence for rejecting  $H_0$ :  $\mu=a$  in favor of  $H_1$ :  $\mu\neq a$ .

f. The  $100(1-\alpha)\%$  confidence interval on  $\mu$  is constructed as follows:

$$\bar{X}_{\cdot} \pm \frac{5_x}{1-(\pi/2)}t_{n-1}\frac{5_x}{\sqrt{n}}$$
 (14.2)

g. In the 1930's a study was performed on the effects on intelligence of plants illegitimate children born of average mothers into foster homes. The mean IQ (as measured by the Kuhlmann revision of Bine's tests) of some 175 children so placed between the ages of 6 months and 1 year was about 115. Critics raised the question of whether the mean score in the population of all infants is actually 100 as was presumed. (Studies revealed the population mean μ to be substantially above 100.)

We wish to test the hypothesis  $H_0$  that the mean Kuhlmann IQ in the population of all infants in the United States is 100 against the alternative hypothesis that it is not:

$$H_0$$
:  $\mu \approx 100$   $H_1$ :  $\mu \neq 100$ .

The probability of falsely rejecting  $H_0$  will be set at .01.

A random sample of 25 infants yielded Kuhlmann IQ's with  $\bar{X} = 113.64$  and  $s_z = 12.40$ . Thus,

$$t = \frac{113.64 - 100}{12.40/\sqrt{25}} = 5.50.$$

The critical values for t are  $-m_0t_M = -2.797$  and  $m_0t_M = 2.797$ .

Hence, we see that  $H_0$  may be rejected in favor of  $H_1$  at the .01 level of significance. Using Eq. (14.2) we find the 99% confidence interval on  $\mu$  is

$$113.64 \pm 2.797 \frac{12.40}{\sqrt{25}} = 113.64 \pm 6.94 = (106.70, 120.58).$$

h. Although the assumption of a normal parent population is made in testing hypotheses about  $\mu$  when  $\sigma_{\nu}^2$  is not known, violations of the assumption have little effect upon either the level of significance or the power of the two-tailed t-test (see Srivastava, 1958). However, nonnormality of the population sampled can have serious effects on one-tailed t-tests of directional hypotheses.

Convenient tables for the calculation of the power of the 1-test can be found in Wine (1964, pp. 254-60).

## 14.3 INFERENCES ABOUT $\mu_1 - \mu_2$ USING INDEPENDENT SAMPLES

a. The hypothesis tested is that the difference between the means of two populations,  $\mu_1 - \mu_2$ , is equal to zero against the alternative hypothesis that it is different from zero:

$$H_0$$
;  $\mu_1 - \mu_2 = 0$   
 $H_1$ :  $\mu_1 - \mu_2 \neq 0$ .

b. It is assumed that  $X_1$  is normally distributed with mean  $\mu_1$  and variance  $\sigma_2^2$  and that  $X_2$  is normally distributed with mean  $\mu_2$  and the same variance  $\sigma_2^2$ . The assumption of equal variances in the two populations is referred to as the assumption of homogeneous variances or homoscedasticity (literally, "same spread"). Furthermore, it is assumed that a sample of size  $n_1$  is randomly drawn from population 1 and that an independent sample of size  $n_2$  is randomly drawn from population 2.

The major consequence of this assumption of independent samples is that the two sample means,  $\bar{X}_1$  and  $\bar{X}_2$ , will be perfectly uncorrelated across infinitely many pairs of samples. The independence assumption would be violated if, for example, sample 1 was a random sample of 10-year-old boys and sample 2 was composed of their sisters. The two means of brother-sister paired samples would correlate on most variables one might observe.

c. Ho is tested against H1 by means of the following test statistic:

$$t = \frac{\overline{X}_{.1} - \overline{X}_{.2}}{\sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}} \left(\frac{1}{n_1} + \frac{1}{n_2}\right)},$$
 (14.3)

where  $\vec{X}_{.1}$  and  $\vec{X}_{.2}$  are the means of the samples from populations 1 and 2, respectively,

 $s_1^2$  and  $s_2^2$  are the unbiased estimates from samples 1 and 2 of the common population variance  $\sigma_x^2$ , and

n, and ne are the sizes of samples 1 and 2.

The critical values against which a t of 2.34 are compared are

$$-\frac{1}{2}$$
 = -2.01 and  $\frac{1}{2}$  = 2.01.

Hence we see that  $H_0$ :  $\mu_1 - \mu_2 = 0$  can be rejected at the .05 level of significance. Indeed, a value of t deviating from 0 by more than 2.34 units has a probability t of approximately .03 if  $H_c$  is true.

The 95% confidence interval on  $\mu_1 - \mu_2$  can be constructed from Eq. (14.4):

$$(\bar{X}_{.1} - \bar{X}_{.2}) \pm {}_{.975}t_{48}s_{2.-1} = 1.65 \pm 2.01(.705) = (0.23, 3.07).$$

h. Violation of the assumption of normality in the *t*-test of  $H_0$ :  $\mu_1 - \mu_2 = 0$  has been shown to have only trivial effects on the level of significance and power of the test and hence should be no cause for concern (Boneau, 1960: Scheffé. 1959, chan. 10).

The effects of violation of the homogeneous variances assumption can be serious depending upon n<sub>1</sub> and n<sub>2</sub>. If n<sub>1</sub> and n<sub>3</sub> are equal, violation of the homogeneous variances assumption is unimportant and need not concern us (Box, 1954a, b; Scheffé, 1959, chap. 10). This fact is a compelling motive for selecting samples equal in size when possible in using the technique in this section. Whenever the variances of populations 1 and 2 are different and n<sub>1</sub> and n<sub>2</sub> are not equal, probabilities of type I and type II errors may be quite different from what one imagines them to be (see Sec. 15.13). When a study in which µ<sub>1</sub> — µ<sub>2</sub> is to be estimated cannot be designed so that n<sub>1</sub> = n<sub>3</sub>, and one suspects that the two populations have substantially different variances, recourse should be made to methods developed by Welch (1937) or Gronow (1951). The problem of testing the significance of the difference between two means when the population variances are unequal has been referred to as the Behrens-Fisher problem (see Fisher, 1959, pp. 93–97).

## 14.4 INFERENCES ABOUT $\mu_1 - \mu_2$ USING DEPENDENT SAMPLES

a. Population 1 has mean  $\mu_1$  and population 2 has mean  $\mu_2$ . The null hypothesis to be tested is the same as in Sec. 14.3:

$$H_0$$
:  $\mu_1 - \mu_2 = 0$   
 $H_1$ :  $\mu_1 - \mu_2 \neq 0$ .

b. It is assumed that samples 1 and 2 are randomly drawn from normal populations with the same variance σ<sup>2</sup><sub>π</sub>. In this instance the samples need not be independent, i.e., there may exist a correlation between X<sub>1</sub> and X<sub>2</sub> over repeated pairs of samples. The following are examples of dependent

samples: sample 1 is a sample of one-year-old infants and sample 2 consists of the fraternal twin-mates of the children in sample 1 is agroup of boys and sample 2 is the group of their sisters; sample 1 is the collection of scores on a reaction-time test made by a group of persons before administration of a drug and sample 2 is the collection of scores made by the same persons after taking the drug.

c. It will always be possible to "pair off" data from two dependent when the pairs may be defined by "brother-sister," "before-after," "twin 1, twin 2," "matched partner 1, matched partner 2," etc. Hence, data gathered from dependent samples will be in the form of n pairs of observations  $X_{t1}$  and  $X_{t2}$  for  $t=1,\ldots,n$ . This pairing of the data from dependent samples will be used to test the hypothesis that  $\mu_1-\mu_2=0$ . The hypothesis that  $X_1$  and  $X_2$  have the same mean, i.e., that  $\mu_1=\mu_2$ , is equivalent to the hypothesis that  $X_1-X_1$  has a mean of 0 in the population. The difference  $X_1-X_2$  between the normally distributed variables  $X_1$  and  $X_2$  is used in ormally distributed; thus the techniques in Sec. 14.2 can be employed to test the hypothesis that the n differences  $X_1-X_2$  can be considered a random sample from a normally distributed population with mean  $\mu_1-\mu_2$  equal to zero.

Denote  $X_{i1}-X_{i2}$  by  $d_{i1}$ , the difference between the paired observations from samples 1 and 2. The test statistic is

$$t = \frac{\dot{d}}{s_d / \sqrt{n}},\tag{14.5}$$

where

$$d = \sum_{i=1}^{n} (X_{i1} - X_{i2})/n = \sum_{i=1}^{n} d_i/n,$$

the average of the n difference scores, and

$$s_d = \sqrt{\sum_{i=1}^{n} \frac{(d_i - d_i)^2}{n-1}},$$

the standard deviation of the n difference scores  $X_{i1} - X_{i2}$ , and n is the number of pairs of observations.

- d. If  $H_i$ :  $\mu_1 \mu_2 = 0$  is true, then t in Eq. (14.5) will follow the Student's distribution with n-1 df. If  $H_i$ :  $\mu_i \mu_2 \neq 0$  is true, then t in Eq. (14.5) will have a distribution identical in shape to the Student's t-distribution with t-1 t-1, but the distribution will have a mean that is different from zero in a direction and by an amount depending partially on the size of t  $\mu_1 \mu_2$ .
  - e. The critical values for testing  $H_{\bullet}$  against  $H_{t}$  at the z-level of signifi-

cance by means of the t-statistic in Eq. (14.5) are as follows:

$$-1-(x/2)t_{n-1}$$
 and  $1-(x/2)t_{n-1}$ .

f. The  $100(1-\alpha)\%$  confidence interval on  $\mu_1 - \mu_2$  around d is constructed as follows:

$$d_{\cdot} \pm \frac{1}{1 - (x/2)} t_{n-1} \frac{s_d}{\sqrt{n}}. \tag{14.6}$$

g. Webster and Bereiter (1963) reported data on personality changes in 100 college women from the Freshman to Senior year. A 60-item personality scale was administered in the Freshman and Senior years to the same 100 women. The first set of 100 scores constitutes sample 1; the 100 Senior-year scores constitute sample 2. There are 100 pairs of "before-after" scores. From these data 100 difference scores are formed by subtracting Senior scores from Freshman scores:  $d_i = X_{i1} - X_{i2}$ . The mean and standard deviation of these 100 d scores are:

$$d_{\cdot} = \sum_{1}^{100} \frac{X_{11} - X_{12}}{100} = -7.02.$$

$$s_{d} = \sqrt{\frac{\sum_{1}^{100} (d_{i} - d_{i})^{2}}{\sum_{1}^{100} (d_{i} - d_{i})^{2}}} = 8.02.$$

We shall test  $H_0$ :  $\mu_1 - \mu_2 = 0$  at the .01 level of significance. The value of t in Eq. (14.5) is

$$t = \frac{d}{s_d/\sqrt{n}} = \frac{-7.02}{8.02/\sqrt{100}} = -8.75,$$

which lies far below the lower critical value of  $-..._{1882}f_{89} = -2.64$ . In fact, the probability p of obtaining a t of -8.75 or less given a true null hypothesis is substantially smaller than even .001. Thus we can confidently reject the null hypothesis that these two dependent samples of 100 observations each could have been randomly sampled from two normal populations with the same mean. There is overwhelming evidence that a "gain" takes place on the personality inventory from the Freshman to the Senior years.

The 99% confidence interval on  $\mu_1 - \mu_2$  can be found from Eq. (14.6):

$$d_{.\pm .995}t_{.99}\frac{s_{d}}{\sqrt{n}} = -7.02 \pm (2.64)\frac{8.02}{\sqrt{100}} = (-9.14, -4.90).$$

h. A common misapplication of inferential statistical techniques is to apply the techniques of Sec. 14.3 when the techniques of this section are

appropriate; in other words, researchers often have dependent samples, fail to recognize this fact, and inappropriately apply the t-test for independent groups to test the hypothesis that  $\mu_1-\mu_2=0$ . The difference between the independent and dependent groups' t-tests becomes apparent when one observes the standard errors of the difference between uncorrelated and correlated means. If  $X_1$  and  $\overline{X}_2$  are uncorrelated (estimated from independent groups), the standard error of the difference between the two means is

$$\sigma_{I,1-I,1} = \sqrt{\frac{\sigma_x^2}{n_1} + \frac{\sigma_x^2}{n_2}}$$

which equals

$$\sigma_{I_{x_1-I_{x_1}}} = \sqrt{\sigma_x^2 \frac{2}{n}} \tag{14.7}$$

when  $n_1=n_2$ . When  $X_1$  and  $X_2$  have a nonzero correlation coefficient of  $\rho_{12}$  (as they will have with two dependent samples), the standard error of  $\vec{X}_{.1}-\vec{X}_{.2}$  is

$$\sigma_{z_{-1}-z_{-1}} = \sqrt{\frac{\sigma_{z}^{2}}{n} + \frac{\sigma_{z}^{2}}{n} - 2\rho_{12}\frac{\sigma_{z}^{2}}{n}} = \sqrt{\sigma_{z}^{2}\left(\frac{2}{n} - \frac{2\rho_{12}}{n}\right)}.$$
 (14.8)

T. The t-statistic for testing  $\mu_1 - \mu_2 = 0$  with independent groups contains  $x_1 - X_4$  in the numerator and an estimate of Eq. (4.4.7) in the denominator. The t-statistic for dependent groups contains  $d_1 = X_1 - X_2$  in the numerator and an estimate of Eq. (14.8) in the denominator; note that

$$\frac{s_d}{\sqrt{n}} = \frac{s_{x_1 - x_2}}{\sqrt{n}} = \sqrt{\frac{s_{x_1 - x_2}^2}{n}} = \sqrt{\frac{s_{x_1}^2}{n} + \frac{s_{x_1}^2}{n} - 2r_{12}\frac{s_{x_1}s_{x_2}}{n}}.$$

If an inappropriate independent-groups t-test is performed with dependent groups for which  $X_t$  and  $X_t$  are substantially positively correlated, the standard error of  $X_t - X_t$  will be greatly overestimated and significant differences between the two means will be branded "nonsignificant." The opposite error, mistaking nonsignificant differences for significant ones, would frequently be made if the independent-groups t-test is applied to dependent groups in which  $X_t$  and  $X_t$  have a substantial negative correlation. Thus we see that the ability to recognize and distinguish independent and dependent samples is an important activity in applying certain inferential statistical techniques. Knowing which test to apply requires a familiarity with the phenomena being studied pass a ensitivity to the statistical problems attendant upon drawing dependent samples.

## I4.5 INFERENCES ABOUT THE VARIANCE, σ<sup>2</sup>, OF A POPULATION

Beginning with this section, we shall deal with testing hypotheses about population variances.

a. The hypothesis to be tested is that a population has a variance  $\sigma_x^2$  equal to some number a versus the hypothesis that  $\sigma_x^2$  is different from a:

$$H_0: \sigma_x^2 = a$$
  
 $H_1: \sigma_x^2 \neq a$ .

b. It must be assumed that the variable X has a normal distribution in the population and that a random sample of n observations has been selected from which  $\sigma_n^2$  will be estimated.

e. The test statistic for testing Ho against H1 is

$$\chi^2 = \frac{(n-1)s_x^2}{a},$$
 (14.9)

where

$$(n-1)s_x^2 = \frac{(n-1)\sum_{1}^{n}(X_i - \bar{X})^2}{n-1} = \sum_{1}^{n}(X_i - \bar{X})^2.$$

d. When  $H_0$  is true, the sampling distribution of  $x^2$  in Eq. (14.9) is the cayanar distribution with n-1 df, i.e.,  $\chi^2_{n-1}$ ; when  $H_1$  is true and  $\sigma^2_z$  is actually equal to some number b different from zero, the sampling distribution of  $(n-1)s^2/a$  will equal b/a times  $\chi^2_{n-1}$ . For example, the graphs of  $(n-1)s^2/a$  (10 are drawn in Fig. 14.2 for the case in which  $\sigma^2_z=10$ , i.e.,  $H_0$  is true, and  $\sigma^2_z=20$ , i.e.,  $H_1$  is true—n=9.

If, from a sample of size 9, a  $s_2^2$  of 21.40 was obtained, the value of the test statistic in Eq. (14.9) would be

$$\chi^2 = \frac{(n-1)s_x^2}{10} = \frac{8(21.40)}{10} = 17.12.$$

In Fig. 14.2 we see that a value of the test statistic as large as 17.12 or later is relatively improbable when  $\sigma_x^2 = 10$  but is quite reasonable when  $\sigma_x^2 = 20$  is true. On the basis of the evidence then, we are inclined to reject  $H_0$  and support  $H_1$ .




FIG. 14.2 Sampling distributions of  $(n-1)s_s^2/10$  when  $H_0$ :  $\sigma_s^1=10$  is true and when  $H_s$ :  $\sigma_s^1=20$  is true (n=9).

e. The critical values for testing  $H_0$  against  $H_1$  at the  $\alpha$ -level of significance are the  $\alpha/2$  and  $1-(\alpha/2)$  percentile points in the chi-square distribution with df=n-1, i.e.,  $\chi^2_{n-1}$ :

$$a/2\chi^2_{n-1}$$
,  $1-(x/2)\chi^2_{n-1}$ .

f. The 100(1  $-\alpha)\%$  confidence interval on the unknown value of  $\sigma_z^2$  is constructed as follows:

$$\frac{(n-1)s_x^2}{\frac{1-(x/2)\chi_{n-1}^2}{2}} < \sigma_x^2 < \frac{(n-1)s_x^2}{\frac{x/2\chi_{n-1}^2}{2}}.$$
 (14.10)

g. For several years the supervisor of curriculum and research in a large school system has observed a typical standard deviation of .80 yr. measured in grade placement units on a standardized test of arithmetic achievement given at the end of the third grade. Unlike previous years, this year saw the introduction of a programmed textbook for arithmetic study in the third grade. One of the highly touted features of programmed instruction is that it does a better job of accommodating individual differences in learning rate than do traditional methods. Hence, at the end of this year, third-grade students should show a different variance on the arithmetic achievement test than in past years. The curriculum supervisor selects a random sample of 25 pupils to whom he will administer the arithmetic achievement test. The data will be used to test the following hypotheses at the .10 level of significance:

$$H_0$$
:  $\sigma_x^2 = (.80)^2 = .64$   
 $H_1$ :  $\sigma_x^2 \neq .64$ .

The variance of the sample of 25 test scores was found to equal 1.14.

The value

$$\chi^2 = \frac{24(1.14)}{64} = 42.75$$

is compared with the critical values

$$_{.05}\chi_{.21}^2 = 13.85, \quad _{.95}\chi_{.21}^2 = 36.42,$$

and is seen to be significant at the .10 level. The probability p of a value of  $\chi^2$  equal to 42.75 or more given that  $H_0$  is true is approximately .01.

The 90% confidence interval on  $\sigma_2^2$  is found by substituting the sample data into Eq. (14.10):

$$\frac{24(1.14)}{36.47} < \sigma_x^2 < \frac{24(1.14)}{13.85} \rightarrow 0.75 < \sigma_x^2 < 1.98.$$

The conclusion seems obligatory that the population of third graders taught by programmed instruction has a greater variance on the arithmetic achievement test than in past years. Notice that inferential statistical methods have made it unnecessary to administer the test to any more than 25 pupils.

h. Unlike hypothesis tests about means using Student's r-distribution, the assumption of sampling from a normal population cannot be taken lightly when testing hypotheses about population variances (see Scheff, 1959, chap. 10). If the population is nonnormal—particularly if it departs substantially from mesokurtosis—the hypothesis test outlined above may be quite in error.

## 14.6 INFERENCES ABOUT $\sigma_1^2/\sigma_2^2$ USING INDEPENDENT SAMPLES

a. The problem at hand possesses greater practical significance than that of testing whether a population has a variance equal to some hypothesized value a. For now we are concerned with two populations (1 and 2), and we wish to test whether their variances  $\sigma_1^2$  and  $\sigma_2^2$  are the same or different;

$$H_0: \sigma_1^2 = \sigma_2^2$$
  
 $H_1: \sigma_1^2 \neq \sigma_2^2$ 

b. It is assumed that a sample of size  $n_1$  is drawn at random from a normal population with mean  $\mu_1$  and variance  $\sigma_1^2$ ; an independent random sample of size  $n_2$  is drawn from a second normal population with mean  $\mu_2$ 

SELECTED INFERENTIAL TECHNIQUES

and variance  $\sigma_2^2$ . The values of  $\mu_1$  and  $\mu_2$  are immaterial and of no interest in testing  $H_a$ .

c. The test statistic for testing Ho against H1 is the ratio of the two sample variances:

$$F = \frac{s_1^2}{s_2^2}. (14.11)$$

- d. When  $H_0$ :  $\sigma_1^2 = \sigma_2^2$  is true, the sampling distribution of  $F = s_1^2/s_2^2$  is the F-distribution with  $n_1-1$  and  $n_2-1$  df. When  $H_1$ :  $\sigma_1^2\neq\sigma_2^2$ , the distribution of  $s_1^2/s_2^2$  is equal to  $\sigma_1^2/\sigma_2^2$  times the F-distribution with  $n_1 - 1$  and  $n_2 - 1$  df. Thus if in reality  $\sigma_1^2/\sigma_2^2 = 2$ , the distribution of  $s_1^2/s_2^2$  will look like the F-distribution transformed by a multiplicative factor of 2.
- e. The critical values against which F in Eq. (14.11) is compared in testing  $H_0$  against  $H_1$  at the  $\alpha$ -level of significance are

$$a/2F_{n_1-1,n_2-2}$$
 and  $a/2F_{n_1-1,n_2-1}$ 

i.e., the  $100(\alpha/2)$  and  $100[1 - (\alpha/2)]$  percentile points in the F-distribution with  $n_1 - 1$  and  $n_2 - 1$  df. The upper percentile points in the F-distributions can be read directly from Table E in Appendix A. The lower percentile points are related as follows to the upper percentiles:

$$_{n/2}F_{n_1-1,n_2-1} = \frac{1}{1-(n/2)F_{n_2-1,n_2-1}}.$$
 (14.12)

f. The 100(1 - z)% confidence interval on the ratio of  $\sigma_1^2$  to  $\sigma_2^2$  is constructed as follows:

$$s^{-1}F_{n_1-1,n_2-1}\frac{s_1^2}{s_2^2} < \frac{\sigma_1^2}{\sigma_2^2} < \frac{\sigma_2^2}{1-(s/2)}F_{n_1-1,n_2-1}\frac{s_1^2}{s_2^2}.$$
 (14.13)

g. In a study by Sears (1940), children were given familiar arithmetic tasks, and upon completion, a randomly chosen half were told they had failed and the remaining half were told they did well. Each child was then asked to estimate the number of seconds it would take him to complete the next task. Observations were made by the experimenter of the difference between a child's goal (in seconds) on the task to be performed and his performance on the task just completed. It was felt that those who experienced failure on the preceding task might be erratic in establishing their goals.

The hypothesis to be tested at the .05 level of significance is that the rariance in the population of children's estimates of future performance is the same whether they have been told they failed or successfully completed

a preliminary task. Sears obtained the following data:

Group 1	Group 2
(told they were	(told they were
successful)	unsuccessful)
$n_1 = 12$ $s_1^2 = 8.16$	$n_2 = 12$ $s_2^2 = 90.45$

The value of the test statistic in Eq. (14.11) is

$$F = \frac{8.16}{90.45} = 0.090.$$

This value is compared with the critical values that are read approximately from Table E:

$$_{.975}F_{11,11} = 3.47$$
 and  $_{.925}F_{11,11} = \frac{1}{_{975}F_{11,11}} = \frac{1}{3.47} = 0.29$ .

Since F=0.09 falls below the lower critical value,  $H_0$ :  $\sigma_1^2=\sigma_2^2$  is rejected at the .05 level of significance.

The 95% confidence interval on  $\sigma_1^2/\sigma_2^2$  is constructed from Eq. (14.13):

$$(0.29) \frac{8.16}{90.45} < \frac{\sigma_1^2}{\sigma_2^2} < (3.47) \frac{8.16}{90.45} \to 0.03 < \frac{\sigma_1^2}{\sigma_2^2} < 0.31.$$

h. The choice of which sample to designate 1 and which to designate 2 is usually arbitrary but it must not be made after observing which sample variance is larger. To observe the larger sample variance and then designate  $s_1^2$  would effectively double the probability of a type I error over what that probability was believed to be. Determining which sample variance to designate  $s_1^2$  might well be done by the flip of a coin. If one's purpose is solely to construct an interval estimate of the ratio of the population variances, the above concern over designating  $s_1^2$  and  $s_2^2$  is immaterial.

The assumption that s<sub>1</sub> and s<sub>2</sub> are derived from independent samples from normal populations cannot be taken lightly—unlike the normality assumption underlying the t-tests of means. If substantial nonnormality is expected the significance test in Sec. 15.14 should be used.

In the past it was customary to test  $H_0: \sigma_1^2 = \sigma_2^2$  prior to performing a *t*-test of the hypothesis  $H_0: \mu_1 - \mu_2 = 0$ . The former hypothesis is a statement of the homogeneous variances assumption made in testing the latter hypothesis. One was advised in the textbooks of the time not to proceed with the simple *t*-test of Sec. 14.3 if  $s_1^2/s_2^2$  led to rejection of  $H_0: \sigma_1^2 = \sigma_1^2$ . Although such advice stemmed from an admirable concern for meeting the

SELECTED INFERENTIAL TECHNIQUES

and variance  $\sigma_2^2$ . The values of  $\mu_1$  and  $\mu_2$  are immaterial and of no interest in testing Ha.

c. The test statistic for testing H<sub>0</sub> against H<sub>1</sub> is the ratio of the two sample variances:

$$F = \frac{s_1^2}{s_2^2}. (14.11)$$

d. When  $H_0$ :  $\sigma_1^2 = \sigma_2^2$  is true, the sampling distribution of  $F = s_1^2/s_2^2$  is the F-distribution with  $n_1 - 1$  and  $n_2 - 1$  df. When  $H_1$ :  $\sigma_1^2 \neq \sigma_2^2$ , the distribution of  $s_1^2/s_2^2$  is equal to  $\sigma_1^2/\sigma_2^2$  times the F-distribution with  $n_1-1$  and  $n_2 - 1$  df. Thus if in reality  $\sigma_1^2/\sigma_2^2 = 2$ , the distribution of  $s_1^2/s_2^2$  will look like the F-distribution transformed by a multiplicative factor of 2.

e. The critical values against which F in Eq. (14.11) is compared in testing  $H_0$  against  $H_1$  at the  $\alpha$ -level of significance are

$$_{n/2}F_{n_1-1,n_2-2}$$
 and  $_{1-(n/2)}F_{n_1-1,n_2-1}$ 

i.e., the  $100(\alpha/2)$  and  $100[1-(\alpha/2)]$  percentile points in the F-distribution with  $n_1 - 1$  and  $n_2 - 1$  df. The upper percentile points in the F-distributions can be read directly from Table E in Appendix A. The lower percentile points are related as follows to the upper percentiles:

$$s_{2}F_{n_{1}-1,n_{2}-1} = \frac{1}{1-t_{2}(t)F_{n_{2}-1,n_{2}-1}}$$
 (14.12)

f. The 100(1 - 2)% confidence interval on the ratio of  $\sigma_1^2$  to  $\sigma_2^2$  is constructed as follows:

$$s_{2}F_{a_{1}-1,a_{2}-1}\frac{s_{2}^{2}}{s_{2}^{2}} < \frac{\sigma_{1}^{2}}{\sigma_{2}^{2}} < \frac{\sigma_{1}^{2}}{\sigma_{2}^{2}} < \frac{1-(s_{2}/2)}{s_{1}}F_{a_{1}-1,a_{2}-1}\frac{s_{1}^{2}}{s_{2}^{2}}.$$
 (14.13)

g. In a study by Sears (1940), children were given familiar arithmetic tasks, and upon completion, a randomly chosen half were told they had failed and the remaining half were told they did well. Each child was then asked to estimate the number of seconds it would take him to complete the next task. Observations were made by the experimenter of the difference between a child's goal (in seconds) on the task to be performed and his performance on the task just completed. It was felt that those who experienced failure on the preceding task might be erratic in establishing their goals.

The hypothesis to be tested at the .05 level of significance is that the rariance in the population of children's estimates of future performance is the same whether they have been told they failed or successfully completed a preliminary task. Sears obtained the following data:

Group 1	Group 2
(told they were	(told they were
successful)	unsuccessful)
$n_1 = 12$ $s_1^2 = 8.16$	$n_2 = 12$ $s_2^2 = 90.45$

The value of the test statistic in Eq. (14.11) is

$$F = \frac{8.16}{90.45} = 0.090.$$

This value is compared with the critical values that are read approximately from Table E:

$$_{.975}F_{11,11} = 3.47$$
 and  $_{.025}F_{11,11} = \frac{1}{_{.975}F_{11,11}} = \frac{1}{3.47} = 0.29$ .

Since F=0.09 falls below the lower critical value,  $H_0$ :  $\sigma_1^2=\sigma_2^2$  is rejected at the .05 level of significance.

The 95% confidence interval on  $\sigma_1^2/\sigma_2^2$  is constructed from Eq. (14.13):

$$(0.29) \frac{8.16}{90.45} < \frac{\sigma_1^2}{\sigma_2^2} < (3.47) \frac{8.16}{90.45} \rightarrow 0.03 < \frac{\sigma_1^2}{\sigma_2^2} < 0.31.$$

h. The choice of which sample to designate 1 and which to designate 2 is usually arbitrary but it must not be made after observing which sample variance is larger. To observe the larger sample variance and then designate  $s_1^2$  would effectively double the probability of a type I error over what that probability was believed to be. Determining which sample variance to designate  $s_1^2$  might well be done by the flip of a coin. If one's purpose is solely to construct an interval estimate of the ratio of the population variances, the above concern over designating  $s_1^2$  and  $s_2^2$  is immaterial.

The assumption that  $s_1^2$  and  $s_2^2$  are derived from *independent* samples from *normal* populations cannot be taken lightly—unlike the normality assumption underlying the *t*-tests of means. If substantial nonnormality is expected the significance test in Sec. 15.14 should be used.

In the past it was customary to test  $H_0$ :  $\sigma_1^2 = \sigma_2^2$  prior to performing a t-test of the hypothesis  $H_0$ :  $\mu_1 - \mu_2 = 0$ . The former hypothesis is a statement of the homogeneous variances assumption made in testing the latter ment of the homogeneous variances assumption made in testing the latter hypothesis. One was advised in the textbooks of the time not to proceed hypothesis. One was advised in the textbooks of the time not to proceed hypothesis. One was advised in  $s_1^2 s_2^2 = s_1^2$ . Ed to rejection of  $H_0$ :  $\sigma_1^2 = \sigma_1^2$ , with the simple t-test of Sec. 14.3 if  $s_1^2/s_2^2 = 1$  led to rejection of  $H_0$ :  $\sigma_1^2 = \sigma_1^2$ . Although such advice stemmed from an admirable concern for meeting the

assumptions of the tests employed, it proved to be generally poor advice. In particular, the preliminary test of the assumption of homogeneous variances can be largely invalidated by nonnormality of the populations; but this same nonnormality is of no consequence to the validity of the t-test of  $\mu_1 - \mu_2 = 0$ . In fact, if  $n_1 = n_1$  there is no reason to be concerned about violation of the assumption of homogeneous variances. The only circumstance in which one might advisedly test  $H_0: \sigma_1^2 = \sigma_1^2$  prior to testing  $H_0: \mu_1 - \mu_2 = 0$  is when there is good evidence that the populations are normally distributed and  $n_1$  and  $n_2$  are quite unequal. It may be possible to find a simple transformation of the observations that will give them a more nearly normal distribution. The subject of "normalizing transformation" is dealt with in detail in P. O. Johnson's Statistical Methods in Research (1949, chap. 7).

## 14.7 INFERENCES ABOUT offort USING DEPENDENT SAMPLES

a. As in Sec. 14.6, the null hypothesis being tested here is that two populations have the same variance:

$$H_0: \sigma_1^2 = \sigma_2^2$$
  
 $H_1: \sigma_1^2 \neq \sigma_2^2$ .

b. It is assumed that two possibly dependent samples are drawn, one of size n from a normal population with variance  $\sigma_1^*$  and the other of the same size n from a normal population with variance  $\sigma_2^*$ . The values of  $\mu_1$  and  $\mu_2$  are not of interest.

The nature of dependent samples was discussed in Sec. 14.4 on testing differences between means.

c. The test statistic used in testing Ho against H1 is

$$t = \frac{s_1^2 - s_1^2}{\sqrt{\frac{4s_1^2s_2^2}{n_1 - s_1^2}}}$$
(14.14)

where si and si are the variances of samples 1 and 2, respectively,

n is the number of pairs of observations, pairing each observation of sample 1 with a single observation in sample 2, and

fit is the correlation coefficient calculated on the n paired observations.

As was pointed out in Sec. 14.4 it will generally be possible to "pair off" the observations in two dependent samples into n pairs ("before-after," "brother-sister," "husband-wife").

- 307
- **d.** When  $H_0: \sigma_1^2 = \sigma_2^2$  is true the sampling distribution of t in Eq. (14.14) is Student's t-distribution with n-2 df.
- e. The critical values for testing  $H_0$  against  $H_1$  at the  $\alpha$ -level of significance are:

- f. Constructing a confidence interval on  $\sigma_1^2 \sigma_2^2$  or  $\sigma_2^2/\sigma_2^2$  when estimates of the variances are obtained from dependent samples presents difficulties beyond the scope of this textbook.
- g. Lord (1963b) reported data gathered by William E. Coffman on the performance of a sample of 95 students on the Stanford Achievement Test in the seventh and eighth grades. One might wonder whether, in the population of students sampled, performance is more uniform (less variable) in the seventh or the eighth grade.

Sample 1 is the set of scores on the Stanford Achievement Test earned by the 95 students in grade seven, and sample 2 is the set of scores earned by the same 95 students in grade eight. Thus the two samples are not independent. The following data were obtained to test  $H_0$ :  $\sigma_1^2 = \sigma_2^2$  at the .10 level of significance:

Sample 1 (grade 7) (grade 8)

$$n = 95$$
  $n = 95$   $s_1^2 = 134.56$   $s_2^2 = 201.64$ 

$$r_{12} = .876$$

The value of  $r_{12} = .876$  is the product-moment correlation coefficient between the 95 students' performance in the seventh and eighth grades. From Eq. (14.14),

$$t = \frac{134.56 - 201.64}{\sqrt{\frac{4(134.56)(201.64)}{95 - 2}(1 - .876^2)}} = -4.07.$$

The critical values with which the obtained t of -4.07 is compared are

$$-a_{st}t_{sq} \approx -1.66$$
 and  $a_{ss}t_{sq} = 1.66$ .

Thus we see that evidence exists to conclude that in the populations sampled—seventh-grade students and eighth-grade students—the variances of and of are different. The probability of obtaining a value of t as discrepant from zero as that obtained is less than .001 if of is truly equal to of. Per-

308

formance on the Stanford Achievement Test is more variable among eighthgrade than among seventh-grade students.

#### 14.8 INFERENCES ABOUT p. THE POPULATION PRODUCT-MOMENT CORRELATION COEFFICIENT

a. The hypothesis being tested is that the product-moment correlation coefficient between variables X and Y, i.e.,  $\rho_{xy}$ , is equal to some value a-between -1 and +1, of course-in the population sampled:

$$H_0$$
:  $\rho_{xy} = a$   
 $H_1$ :  $\rho_{xy} \neq a$ .

b. It is assumed that a random sample of n paired observations  $(X_i, Y_i)$ is drawn from a bivariate normal population (see Sec. 6.7) in which the correlation of X and Y is  $\rho_{xy}$ . The means and variances for the bivariate normal population are not of interest.

c. To test  $H_0$  against  $H_1$ , one first calculates  $r_{xy}$ , the sample productmoment correlation coefficient. The value of  $r_{xy}$  is then transformed by means of Fisher's Z-transformation of r (see Table G in Appendix A) into  $Z_r$ . The test statistic for testing the null hypothesis that  $\rho_{rr} = a$  is

$$z = \frac{Z_r - Z_o}{16\sqrt{r}},$$
 (14.15)

where Z, is the Z-transformed value corresponding to the sample value of rz, Z<sub>a</sub> is the Z-transformed value corresponding to a, the value hypothesized for  $\rho_{xy}$  in  $H_0$ , and n is the sample size.

d. When  $H_0$  is true, i.e.,  $\rho_{xy}$  is equal to a, then z in Eq. (14.15) is normally distributed with mean 0 and standard deviation 1, i.e., z has the unit normal distribution. If, on the other hand,  $H_1$  is true and  $ho_{xy} = b$  that is different from a, then z in Eq. (14.15) has a normal distribution with standard deviation 1 and centering around a mean of  $Z_0 - Z_0$ . For example, if  $H_0$ ,  $\rho_{xy} = 0$  is true, the distribution of  $z = (Z_y - 0)/(1/\sqrt{n-3})$  for samples of size 12 is represented by the curve on the left in Fig. 14.3. If  $H_1: \rho_n = .60$  is true and n = 12, the sampling distribution of  $z = (Z_r 0)/(1/\sqrt{n-3})$  is the curve on the right in Fig. 14.3.




FIG. 14.3 Sampling distributions of  $z = (Z_r - 0)/(1/\sqrt{n-3})$  for samples of size 12 when  $\rho_{re} = 0$  and when  $\rho_{re} = .60$ .

e. The critical values against which z in Eq. (14.15) is compared in testing  $H_0$  against  $H_1$  at the x-level of significance are the  $100(\pi/2)$  and 100[1 - (g/2)] necentiles in the unit normal distribution.

For example, if  $H_0$  is tested at the .05 level, the critical values are -1.96 and +1.96.

f. Confidence intervals are constructed on  $\rho_{xy}$  by first constructing a confidence interval on  $Z_\rho$  around  $Z_r$  and transforming the upper and lower limits on this interval back to the scale of r by reading Table G in reverse.

The first step in building the  $100(1-\alpha)\%$  confidence interval on  $\rho_{xy}$  is to calculate

$$Z_r \pm \frac{1}{1-(\pi/2)^2} \sqrt{\frac{1}{n-3}},$$
 (14.16)

where  $Z_n$  and n are as defined in Eq. (14.15) above, and

 $\frac{1}{1-(\alpha/2)^2}$  is the  $100[1-(\alpha/2)]$  percentile in the unit normal distribution.

Equation (14.16) will determine two points on the Z-transformation scale. Table G is entered with these two values and the two corresponding values of  $r_{xy}$  are determined. These two values of  $r_{xy}$  constitute the  $100(1-\alpha)\%$ , confidence interval on  $\rho_{xy}$ .

g. Forehand and Libby (1962) gathered data on the correlation between "supervisors' ratings of innovative behavior," X, and scores on tests of divergent thinking ("flexibility" and "implications"—known in the vernacular as tests of "creativity"), Y, on n=60 governmental administrators. An  $r_{r_0}$  of .30 was obtained.

We shall test the null hypothesis with  $\alpha = .05$  that  $\rho_{xy}$  is zero:

$$H_0$$
:  $\rho_{xy} = 0$   
 $H_1$ :  $\rho_{xy} \neq 0$ .

From Table G, the Z-transformed value of an r of .30 is .310, i.e.,  $Z_{20} = .310$ . Of course,  $Z_0 = 0$ . Thus, the value of z in Eq. (14.15) is

$$z = \frac{.310 - 0}{1/\sqrt{60 - 3}} = .310(\sqrt{57}) = 2.34.$$

The obtained value of z is compared with

and

310

$$_{.025}z = -1.96$$
  
 $_{.025}z = 1.96$ .

Since 2.34 exceeds 1.96,  $H_0$  can be rejected and  $H_1$  accepted at the .05 level of significance. The 95% confidence interval on  $\rho_{xy}$  is constructed as follows:

$$Z_{.30} \pm (1.96) \frac{1}{\sqrt{60}} = .310 \pm .259 = (.051, .569).$$

From Table G, the values of rev corresponding to Z-values of .051 and .569 are found. These values are approximately .05 and .51, respectively. Thus the 95% confidence interval on  $\rho_{xy}$  is (.05, .51).

h. In the vast majority of instances in practice the value of  $\rho_{xy}$  hypothesized in  $H_0$  is 0. Hence,  $(Z_r - 0)/(1/\sqrt{n-3}) = Z_r(\sqrt{n-3})$  is referred to a table of the unit normal distribution. An alternative test statistic exists that exactly describes the sampling distribution of  $r_{xy}$  when  $\rho_{xy}$  is 0. (The hypothesis test outlined above is an approximate one, but one which is so close to the exact test for even small n that the fact that it is an approximation should cause no concern.) When  $\rho_{xx}$  is 0,

$$t = \frac{r_{sv}}{\sqrt{(1 - r_{ro}^2)/(n - 2)}}$$

has the Student I-distribution with n-2 df. This fact was used to construct a table of critical values of  $r_{xy}$  to facilitate testing  $H_0$ :  $\rho_{xy}=0$ . Table I in Appendix A lists the values which  $r_{xy}$  must exceed in absolute value to constitute evidence for rejection of  $H_0$ :  $\rho_{xy}=0$ . For example, with  $\pi=60$ , a sample r<sub>sp</sub> must lie above approximately .250 or below -..250 before  $H_{\bullet}$ ,  $\rho_{zz}=0$  can be rejected at the .05 level of significance. (Table I is entered for particular values of x and df = n - 2.) Hence, a sample  $r_{ex}$  of .30 leads to rejection, as was seen above. As a further example of how to read Table I, the critical values of  $r_{ss}$  for testing  $\rho_{rs}=0$  at the .01 level of significance with a sample of size 12 are .708 and -.708. An  $r_{er}$  of -.45, say, for n=12, would not lie sufficiently far from 0 to allow one to reject  $H_0$ :  $\rho_{xy}=0$ .

For a study of the effects of violation of the assumption of bivariate normality, see Norris and Hjelm (1961) and Carroll (1961).

# I4.9 INFERENCES ABOUT $\rho_1-\rho_2$ USING INDEPENDENT SAMPLES

a. In this instance, inferences concern the possible difference between the correlation  $\rho_1$  of X and Y in population 1 and the correlation  $\rho_2$  of the same two variables in population 2. For example, are aptitude X and achievement Y more highly correlated for boys (population 1) than for girls (population 2)? The null hypothesis is generally that  $\rho_1 = \rho_2$ ; the alternative hypothesis is that they are unequal:

$$H_0$$
:  $\rho_1 = \rho_2$   
 $H_1$ :  $\rho_1 \neq \rho_2$ 

- b. It is assumed that a random sample of size  $n_1$  is drawn from a bivariate-normal population 1 in which the product-moment correlation coefficient is  $\rho_1$  and that an independent random sample of size  $n_2$  is drawn from a bivariate-normal population 2 with correlation  $\rho_2$ .
- c. Again the inferential statistical problems are handled by means of Fisher's Z-transformation of r. Samples 1 and 2 are drawn from populations 1 and 2, respectively. The two sample correlation coefficients,  $r_1$  and  $r_2$ , are calculated and then transformed to  $Z_{r_2}$  and  $Z_{r_2}$  by means of Table G. The test statistic for testing  $H_0: p_1 = p_2$  is:

$$z = \frac{Z_{r_1} - Z_{r_2}}{\sqrt{\frac{1}{n_1 - 3} + \frac{1}{n_2 - 3}}}.$$
 (14.17)

- d. If in fact  $\rho_1$  equals  $\rho_2$ , then z in Eq. (14.17) has a normal distribution with mean 0 and standard deviation 1 over repeated pairs of independent random samples. If  $\rho_1$  and  $\rho_2$  actually differ, then the mean of the sampling distribution of z in Eq. (14.17) will shift away from 0—it will become  $Z_{\rho_1}$ —but the standard deviation will remain equal to 1.
- e. To test  $H_0$ :  $\rho_1=\rho_2$  against  $H_1$ :  $\rho_1\neq\rho_2$  at the  $\alpha$ -level of significance, the single calculated value of z in Eq. (14.17) is compared with the  $100(\alpha/2)$  and  $100[1-(\alpha/2)]$  percentiles in the unit normal distribution:

$$a/2z$$
 and  $a-(a/2)z$ .

f. Confidence intervals on  $\rho_1 - \rho_2$  are constructed by way of confidence intervals on the Z-transformation of  $\rho_1$  and  $\rho_2$ . To find the  $100(1-\alpha)\%$ 

confidence interval on  $\rho_1 - \rho_2$ , one first calculates

$$(Z_{r_1} - Z_{r_2}) \pm \frac{1}{1 - (r_2/2)^2} \sqrt{\frac{1}{n_1 - 3} + \frac{1}{n_2 - 3}}.$$
 (14.18)

The two values on the Z scale obtained from Eq. (14.18) are then transformed to the r scale by means of Table G. The resulting two values of r form the  $100(1-\alpha)\%$  confidence interval on  $\rho_1-\rho_2$ . The problems of attaching meaning to a value for  $\rho_1-\rho_2$  will be discussed under (h) below.

g. The techniques of this section will be illustrated on data based loosely on studies by Hinton (1939) and Dispensa (1938). In a sample of 200 children ranging in age from six to 15, the correlation between intelligence (Stanford-Binet) X and basal metabolic rate Y was  $r_1 = .71$ . In a sample of 78 adults ranging in age from 18 to 25, intelligence and basal metabolic rate correlated  $r_2 = .28$ . We shall test  $H_0: p_1 = p_2$  at the .01 level of significance. Hence, the critical values for x in Eq. (14.17) are -2.58 and 2.58.

$$Z_{.71} = .887$$
  $Z_{.23} = .288$ .

$$z = \frac{Z_{71} - Z_{51}}{\sqrt{\frac{1}{n_1 - 3} + \frac{1}{n_2 - 3}}} = \frac{.887 - .288}{\sqrt{\frac{1}{200 - 3} + \frac{1}{78 - 3}}} = \frac{.599}{.136} = 4.40.$$

We see that  $H_a$ :  $\rho_1=\rho_2$  can be rejected at the .01 level of significance. The 99% confidence interval on  $\rho_1-\rho_2$  is found from Eq. (14.18) as follows:

.887 - .288 
$$\pm 2.58 \sqrt{\frac{1}{200 - 3} + \frac{1}{78 - 3}} = .599 \pm .350 = (.249, .949).$$

Transforming these two Z-values back to the r-scale yields the 99% confidence interval on  $\rho_1 = \rho_2$ : .24, .74.

h. While testing the null hypothesis that  $\rho_1=\rho_2$  makes perfectly good sense, merely estimating the difference between the two coefficients with  $r_1-r_2$  and setting a confidence interval around this sample difference may lack meaning. Suppose, for example, that  $\rho_1-\rho_2$  is 20. Any one of the following conditions on  $\rho_1$  and  $\rho_2$  could hold: (1)  $\rho_1=.90$  and  $\rho_2=.70$ : each case the difference between  $\rho_2$  and  $\rho_3$  is the same, .20, the meaning that each case the difference between  $\rho_2$  and  $\rho_3$  is the same, .20, the meaning that attaches to the three conditions is quite different. In (1) the .20 difference represents a much larger difference in the ability to predict one variable from the other than the .20 difference in (2). Moreover, the .20 difference between .10 and -.10 represents no difference in the ability to predict one variable from the other. Thus, the unadorned difference between  $\rho_1$  and  $\rho_2$ 

is of little interest, although testing hypotheses about that difference or merely noting the values of  $r_1$  and  $r_2$  are both meaningful approaches.

Hypothesis testing and inferential techniques exist for making inferences about a set of J correlation coefficients for variables X and Y when J independent estimates,  $r_{xy}$ , are available. For example, one might test the hypothesis that intelligence X and age at which a child begins to speak Y are equally highly correlated for the three populations of first-, second-, and third-born children. For a presentation of appropriate inferential techniques see Marascuilo (1966).

# 14.10 INFERENCES ABOUT Pzy - Pzz USING DEPENDENT SAMPLES

a. The null hypothesis to be tested is that a variable X has the same correlation with two other variables, Y and Z, against the alternative hypothesis that  $\rho_{xx}$  and  $\rho_{xx}$  are not equal:

$$H_0$$
:  $\rho_{xy} = \rho_{xx}$   
 $H_1$ :  $\rho_{xy} \neq \rho_{xx}$ 

This situation would arise if one wished to predict "academic success" (as measured by a grade-point average) and had two potential predictors, Y and Z. If for financial reasons only one predictor (Y or Z) could be used, it would be wise to gather data, estimate  $r_{xy}$  and  $r_{xz}$ , and then test the hypothesis that the observed difference between the two sample r's represented a true difference between  $r_{xy}$  and  $r_{xz}$ .

- b. It is assumed that there exist three bivariate-normal populations, one for each of the pairs of variables X and Y, X and Z, and Y and Z. A single random sample of n persons is drawn from which the three correlation coefficients  $r_{zy}$ ,  $r_{zx}$ , and  $r_{yx}$  are calculated. Obviously, these three estimates of correlation are not independent.
  - c. The test statistic for testing  $H_0$  against  $H_1$  is:

$$z = \frac{\sqrt{n}(r_{xy} - r_{xz})}{\sqrt{(1 - r_{xy}^2)^2 + (1 - r_{xx}^2)^2 - 2r_{xz}^2}}$$

$$-(2r_{yx} - r_{xy}r_{xx})(1 - r_{xy}^2 - r_{xx}^3 - r_{yx}^3)$$
(14.19)

where n is the sample size,  $r_{ev}$  is the sample correlation of X and Y,  $r_{zz}$  is the sample correlation of X and Z, and  $r_{vv}$  is the sample correlation of Y and Z.

- 14
- d. If  $H_0$  is true, i.e.,  $\rho_{xy} = \rho_{xy}$ , then z in Eq. (14.19) has a sampling distribution over repeated samples of size n that is closely approximated by the normal distribution with mean 0 and standard deviation 1 (see Olkin and Siotani, 1964, and Olkin, 1967). When  $H_1$  is true, the mean of the sampling distribution of z in Eq. (14.19) shifts away from zero but the standard deviation remains approximately equal to 1.
  - e. As might be expected, the null hypothesis  $H_0$ :  $\rho_{xy} = \rho_{xz}$  is tested at the  $\alpha$ -level of significance by comparing the observed value of z in Eq. (14.19) with the  $100(\alpha/2)$  and  $100[1-(\alpha/2)]$  percentile points in the unit normal distribution; the two critical values for the hypothesis test are

- f. Interval estimation of the difference  $\rho_{xx} \rho_{xx}$  is difficult to justify for the same reasons discussed at the end of Sec. 14.9. However, if such estimation can ever be justified, appropriate techniques may be found in Olkin (1967, p. 113).
- g. Suppose that success in college X (as measured by grade-point average at the end of the first year) can be predicted either from test battery Y or test battery Z. Only one test battery can be employed because of the short available testing time. It is desired to test the hypothesis  $H_0$ :  $\rho_{xy} = \rho_{xx}$  at the .05 level of significance,

For a random sample of 100 freshmen who were given both test batteries as high-school seniors, the three possible correlation coefficients among X, Y, and Z are:

$$r_{xv} = .56$$
,  $r_{xx} = .43$ ,  $r_{yx} = .52$ .

The value of z in Eq. (14.19) is found to be

$$z = \frac{\sqrt{100(.56 - .43)}}{\sqrt{(1 - .56^3)^2 + (1 - .43^3)^2 - 2(.52)^3}} - \frac{10(.13)}{\sqrt{.659^7}} = \frac{10.00}{.3122} = 1.60.$$

The obtained value of z falls far below the required critical value of 1.96. This he hypothesis that batteries Y and Z are alike with respect to accuracy in predicting X cannot be rejected. Unless a large sample would alter this conclusion, the decision to administer Y or Z in high school could not be made on the basis of relative predictive accuracies.

## 14.11 INFERENCES ABOUT OTHER CORRELATION COFFEIGURES

In this section, significance testing procedures for the correlation coefficients presented in Chapter 9 will be discussed. The presentation of inferential techniques will be greatly simplified since space does not permit detailed treatments such as those in the previous sections of this chapter. Procedures will be presented with which the null hypothesis of no correlation between variables X and Y can be tested against a nondirectional alternative hypothesis for most of the coefficients in Chapter 9. Where possible, a test statistic will be defined that has a known or approximately known sampling distribution given that the two variables being correlated—whether they are measured dichotomously, with ranks, or otherwise—are unrelated, i.e., have a coefficient of zero in the population. In each instance, the significance testing procedure will be illustrated with hypothetical data.

#### The Phi Coefficient, &

Suppose that a random sample of n persons is drawn from a population in which two dichotomously scored variables, X and Y, have a population phi coefficient of zero. For large values of n (20 or greater, say) the sampling distribution of  $\sqrt{n} \phi$ , where  $\phi$  is the sample phi coefficient, is approximately normal with mean 0 and standard deviation 1. (When the population phi coefficient is different from zero, the sampling distribution of  $\sqrt{n} \phi$  becomes skewed and centers around a mean different from zero by an amount that increases as the population value of  $\phi$  deviates further from 0.)

A random sample of n=25 persons is observed on two dichotomous variables: X is the variable "sex" scored 0—female and 1—male; Y is the variable "attrition" scored 0—"dropped out of school" and 1—"remained in school." The sample value of  $\phi$  is equal to -.41. The test statistic is found as follows:

$$z = \sqrt{n} \ \phi = \sqrt{25} \ (-.41) = 5(-.41) = -2.05.$$

This value of z lies just below the 2.5th percentile in the unit normal distribution ( $_{esz} = -1.96$ ). Hence, the hypothesis that sex and attrition from high school are unrelated in the population sampled can be rejected at the .05 level of significance. (It is interesting to note that a sample  $\phi$  lying as far from zero as -30 would not have led to rejection of the null hypothesis at the .05 level for a sample of 25 persons:  $\sqrt{n} \phi = -1.50 > -1.96$ .)

## Spearman's Rank-Correlation Coefficient, r.

The sampling distribution of r, given zero correlation between two sets of ranks X and Y in the population cannot be characterized in terms of any well-known statistical distributions for n less than approximately 10. For n > 10, say, the sampling distribution of r, when the population Spearman rank-correlation coefficient is zero is related approximately to Student's t-distribution. In fact, for n > 10

$$t = \frac{r_s}{\sqrt{(1 - r_s^2)/(n - 2)}}$$
(14.20)

is approximately distributed as Student's *t*-distribution with n-2 df when the population value of r, is zero.

Suppose, for example, that in a sample of n=22, the value of r, is .38. The value of t is

$$t = \frac{.38}{\sqrt{\frac{1 - .38^2}{22 - 2}}} = \frac{.38}{.207} = 1.84.$$

To test the hypothesis of no correlation in the population, at the .01 level, the above value of t=1.84 is compared with  $\pm_{.000}t_{10}=\pm2.845$ . Since t lies between the .5th and 99.5th percentiles in the t-distribution with 20 dt, the null hypothesis of no correlation in the population cannot be rejected at the .01 level.

#### Kendali's au

The significance of a sample value of Kendall's  $\tau$  is most conveniently tested in terms of one of the components in its calculation. As was seen in Sec. 9.3, for a sample of size n,  $\tau$  is found from the formula

$$\tau = \frac{S}{n(n-1)/2},$$

where S = P - Q is the difference between the total number of "agreements"

P and the total number of "inversions" Q in the two arrays of ranks (see Sec. 9.3 for definitions of "agreements" and "inversions").

The sampling distribution of S is more conveniently studied than that of r. When n is greater than or equal to 10, the sampling distribution of S is nearly normal when X and Y are uncorrelated in the population; the standard deviation of S is approximately

$$\sqrt{n(n-1)(2n+5)/18}$$

The sampling distribution of S can be made more nearly normal by a simple "continuity correction" by which S is transformed into a quantity that will be denoted by  $S^*$ :

If S is negative, 
$$S^* = S + 1$$
.  
If S is positive,  $S^* = S - 1$ .

(In some textbooks, one is directed to calculate  $S^*$  by |S|-1. This procedure is incorrect and approximately doubles the probability of a type I error.)

It follows that

$$z = \frac{S^*}{\sqrt{n(n-1)(2n+5)/18}}$$
 (14.21)

closely follows the normal distribution with mean 0 and standard deviation l when n>10 and the null hypothesis of no relationship between the two variables is true.

Suppose that n = 10 and S = 9 so that  $\tau = 9/[10(9)/2] = .20$ . Since S is positive.  $S^* = S - 1 = 9 - 1 = 8$ . The value of z is

$$z = \frac{8}{\sqrt{10(10-1)[(2\cdot10)+5]/18}} = \frac{8}{\sqrt{125}} = 0.72.$$

The probability is greater than .40 that a z score will lie either above +0.72 or below -0.72. Hence we see that a  $\tau=.20$  for a sample of size 10 gives no evidence that would lead one to conclude the value of  $\tau$  is different from zero in the population sampled.

For values of n from 4 to 10, Kendall (1962) tabulated certain characteristics of the sampling distribution of S. Table L of Appendix A lists the probabilities of obtaining values of S equal to or greater than certain tabulated values when  $n=4,\ldots,10$  and the population value of  $\tau$  is zero. For example, when n=8 and the population value of  $\tau$  is zero, we see from Table L that the probability of obtaining a value of S equal to 14 or greater is .054. S attains a value of T less with probability .054. Thus a value of T less with an T of T is zero, with an T of T or T of T or T

## The Point-Biserial Correlation Coefficient, r.,

By inspection of the formula for  $r_{pb}$  it can be seen that  $r_{pb}$  is equal to zero if and only if  $\bar{X}_{.1} = \bar{X}_{.0}$ :

$$r_{pb} = \frac{\overline{X}_1 - \overline{X}_0}{s_x} \sqrt{\frac{n_1 n_0}{n(n-1)}}.$$

Testing the null hypothesis that a population value of the point-biserial correlation coefficient is zero is closely equivalent to testing the hypothesis that in the population those persons scoring 1 on the dichotomous variable have a mean equal to the population mean of the persons scoring 0 on the dichotomy, i.e.,  $H_0$ :  $\mu_1 = \mu_0$ . If  $r_{sb}$  is zero in the population sampled, then

$$t = \frac{r_{ph}}{\sqrt{(1 - r_{ph}^2)/(n - 2)}}$$
 (14.22)

is approximately distributed as Student's t-distribution with n-2 df.

Suppose that in a sample of size n = 18 the value of  $r_{pb}$  is .56. The value of t is found as follows:

$$t = \frac{.56}{\sqrt{\frac{1 - (.56)^2}{18 - 2}}} = \frac{.56}{\sqrt{.6864}} = 2.70.$$

A t value of 2.70 exceeds the 99th percentile in Student's t-distribution with 16 df. Hence, it appears unlikely that the point-biserial correlation coefficient is zero in the population sampled.

## The Tetrachoric Correlation Coefficient, r.,

When the null hypothesis of no relationship between X and Y (two normally distributed variables forced into dichotomies) is true, the sampling distribution of the sample tetrachoric correlation coefficient,  $r_{tt}$ , is approximately normal for n greater than about 20 with mean 0 and standard deviation

$$\sigma_{r_{tot}} = \sqrt{\frac{p_x p_y q_z q_y}{n}} \frac{1}{u_z u}, \qquad (14.23)$$

where n is the sample size,

 $P_s = n_s/n$ , the proportion of persons scoring 1 on the dichotomously measured X variable,

 $P_n = n_0 / n_0$ , the proportion of persons scoring 1 on the dichotomously  $P_n = n_0 / n_0$ . It is the measured Y variable,

 $u_x$  is the ordinate above the z-score on the unit normal curve above which  $p_x$  proportion of the area lies (see Table B), and

 $u_v$  is the ordinate above the z-score on the unit normal curve above which  $p_v$  proportion of the area lies.

Thus, for moderately large and large n

$$z = \frac{r_{tel}}{\sigma_{rel}}$$

can be referred to the unit normal distribution to test the hypothesis that the population tetrachoric correlation coefficient is zero.

Suppose the data in Fig. 14.4 are gathered in studying the relationship between two variables X and Y that are believed to be normally distributed but can only be measured dichotomously.

The value of (bc)/(ad) is (21)(19)/(4)(6) = 399/24 = 16.625. In Table H in Appendix A 16.625 is found to correspond to an  $r_{t,t}$  of .81.

The standard error of rea is estimated from the following data:

$$p_x = \frac{2.5}{5.0} = .50$$
  $q_x = .50$   
 $p_y = \frac{2.3}{5.0} = .46$   $q_y = .54$ .

From Table B, the ordinate on the unit normal curve at the point above which  $p_a = .50$  of the area lies is  $u_c = .3989$ ; the ordinate on the curve at the point above which  $p_v = .46$  of the area lies is  $u_p = .3970$ . Thus the value of z is

$$z = \frac{.81}{\sqrt{\frac{(.50)(.46)(.50)(.54)}{50}} \frac{1}{(.3989)(.3970)}} = 3.64.$$

A z score of 3.64 far exceeds even the 99.5th percentile in the unit normal distribution. We must conclude that the population tetrachoric correlation coefficient is nonzero.

## The Biserial Correlation Coefficient, rs...

The exact sampling distribution of  $r_{bis}$  is not known. It was never derived by Pearson, and the only substantial finding was that of Soper (1914), who derived the standard deviation of  $r_{bis}$  for large samples. McNamara and 320

Dunlap (1934) argued that when the population biserial correlation coefficient is zero, then for large samples  $r_{str}$  should be approximately normally distributed with mean 0 and standard deviation

$$\sigma_{r_{0in}} \doteq \frac{\sqrt{n_1 n_0}}{u n \sqrt{n}},\tag{14.24}$$

where n is the sample size,

 $n_1$  is the number of persons scoring 1 on the dichotomous variable  $(n_0 = n - n_1)$ , and

u is the ordinate on the unit normal curve at the point above which n<sub>1</sub>/n proportion of the area under the curve lies.

When the population biserial correlation coefficient departs from zero, the value of  $\sigma_{r_{kt}}$ , is diminished by  $1/\sqrt{n}$  times the square of the population value of  $r_{kt}$ . When the population value of  $r_{kt}$ , is different from zero, the sampling distribution of  $r_{kt}$  becomes nonnormal, being skewed toward zero.

Empirical sampling studies by Lord (1963a) and Baker (1965) have shown that the above large-sample estimate of the standard error of  $r_{th}$ , is quite nearly exact; in Baker's case this was true even for samples of size 15.

Suppose that for a sample of size 36 in which  $n_1 = 16$  and  $n_0 = 20$ , the value of  $r_{bis}$  is -.145. The value of  $\sigma_{r_{bis}}$  when the population biserial coefficient is zero is

$$\sigma_{r_{bla}} = \frac{\sqrt{16 \cdot 20}}{(3951)36\sqrt{36}} = .210.$$

If the population value of  $r_{bis}$  is zero,  $r_{bis}|\sigma_{r_{bis}}$  will be approximately normally distributed with mean 0 and standard deviation 1 over repeated random samples of size n.

$$t = \frac{r_{bis}}{\sigma_{r_{bis}}} = \frac{-.145}{.210} = -.69.$$

A z value of -.69 appears to be a not unusual observation to obtain in sampling from a unit normal distribution. Hence, evidence does not exist to allow rejection of the null hypothesis of a population biserial correlation coefficient of zero.

## The Rank-Biserial Correlation Coefficient, r.

For a discussion of significance tests for the rank-biserial correlation coefficient  $r_{rb}$  see Cureton (1956) and Glass (1966b).

## The Partial Correlation Coefficient, r....

The partial correlation coefficient between X and Y with variable Z "held constant" was developed in Sec. 9.4. There it was seen that the correlation of the errors of estimate when both X and Y are predicted individually from Z is

$$r_{xy\cdot t} = \frac{r_{xy} - r_{xx}r_{yx}}{\sqrt{(1 - r_{xx}^2)(1 - r_{yx}^2)}}$$

The population partial correlation of X and Y with Z "partialed out" can be denoted by

$$\rho_{xy-t} = \frac{\rho_{xy} - \rho_{xt}\rho_{yx}}{\sqrt{(1 - \rho_{xy}^2)(1 - \rho_{yy}^2)}}$$

When  $\rho_{xy}$  equals zero, the test statistic

$$t = \frac{r_{xy\cdot z}}{\sqrt{(1 - r_{xy\cdot z}^2)/(n - 3)}}$$
(14.25)

has Student's t-distribution with n-3 df. Suppose that in a sample of n=12, the value of  $r_{\pi y 2}$  is .80. Then

$$t = \frac{.80}{\sqrt{(1 - .80^2)/(12 - 3)}} = 4.00.$$

The 99.5th percentile in the r-distribution with 9 degrees of freedom is 3.250. The null hypothesis of a zero population partial correlation coefficient can be rejected at the .01 level of significance.

#### 14.12 INFERENCES ABOUT P, THE POPULATION PROPORTION

a. We shall now be concerned with the proportion of "units" (persons, families, schools, counties, etc.) in a population that possess some characteristic (blue eyes, a late-model car, a "modern math" curriculum, a children's diagnostic center, etc.). The proportion of the units in the population that possess the characteristic in question will be denoted by P, which equals the number of the units possessing the characteristic divided by the total number of the units in the population. For example, of the 60,000 pupils in a particular school system, 3000 are of Spanish-American descent; hence, P = 3000/

60,000 == .05 when the characteristic being observed is "Spanish-American vs. non-Spanish-American descent."

The hypothesis to be tested is that in a relatively large population, the proportion P possessing a particular characteristic is equal to a value a that lies between 0 and 1.0, inclusive:

$$H_0: P = a$$
  
 $H_1: P \neq a$ 

b. For purposes of testing  $H_0$  against  $H_1$  it need only be assumed that a random sample of size n is drawn from the population.

c. Within the sample of n units, the number f who possess the characteristic in question is observed. The sample proportion p is the ratio of f to n:

$$p=\frac{f}{n}$$
.

The statistic p is an estimator of P. In fact, if we think of a dichotomously measured variable X that equals I when the unit observed possesses the characteristic in question and I0 when it does not, then I9 is I1 to I2 as I3 is to I4. For example, suppose that "male gender" is the characteristic being measured. A child's score is denoted by  $X_i$ , which equals I1 if the child is male and I3 if female. If in a sample of I2 children there are I3 males ("I's scorers"), then

$$\sum_{i=1}^n X_i = X_1 + \ldots + X_n = f.$$

Hence,

 $\tilde{X}_{\cdot} = \frac{\sum X}{n} = \frac{f}{n} = p.$ 

The population mean of X will be P. Thus p possesses all the properties as an estimator of P which X possesses as an estimator of  $\mu$ .

In this context it is also interesting to note that the dichotomously scored X variable has a rariance in the population that equals

$$\sigma_x^2 = E(X - P)^2 = P(1 - P),$$

where P is the population proportion. This fact will be used below.

d. Suppose that the null hypothesis is true so that P equals a. You probably recall that the sampling distribution of X, has a mean  $\mu$  and standard deviation  $\sigma_a / \sqrt{n}$ . The same is true of p. For samples of size n, the sampling distribution of p has a mean of a—the population proportion—and a standard

deviation of  $\sigma_a / \sqrt{n} = \sqrt{a(1-a)/n}$ :

$$E(p) = a$$

$$\sigma_p = \sqrt{\frac{a(1-a)}{a}}.$$

The question of the shape of the sampling distribution of p is answered by appealing to the central limit theorem. When n is sufficiently large, the sampling distribution of p is normal with mean P and standard deviation  $\sqrt{P(1-P)/n}$ . "Sufficiently large" can be taken to mean that before p can be considered normally distributed, the value of nP or n(1-P), whichever is smaller, must be greater than 5. This rule-of-thumb depends on the sample size n and the unknown population proportion P. Suppose, for example, that an anthropologist wishes to make an inference about the proportion of left-handed persons in an African tribe. (P is as small as .02 in some African tribes, but P is greater than .10 in the United States.) The anthropologist would do well to draw a sample as large as 500 to guard against the possibility that P is as small as .01-nP = 500(.01) = 5.

If the value of P is a, i.e., if Ho is true, and if na is greater than 5, then

$$z = \frac{p - a}{\sqrt{a(1 - a)/n}}$$
 (14.26)

is approximately normally distributed over random samples of size n with mean 0 and standard deviation 1.

If P is some value b different from a, p will, of course, be approximately normally distributed with mean b and standard deviation  $\sqrt{b(1-b)/n}$ , provided nb or n(1-b), whichever is smaller, is greater than about 5.

e. To test  $H_0$  against  $H_1$  at the  $\alpha$ -level of significance, the value of z in Eq. (14.26) is compared with the  $100(\alpha/2)$  and  $100[1 - (\alpha/2)]$  percentiles in the unit normal distribution:

$$a/2^{Z}$$
  $1-(a/2)^{Z}$ 

f. The 95% and 99% confidence intervals on P around p can be found most easily by reference to Table P in Appendix A. Suppose, for example, that in a sample of n = 100, the value of p is .60. The 95% confidence interval on P ranges approximately from .51 to .71. For large samples, the probability is only approximately  $1-\alpha$  that  $p\pm_{1-(a/2)}z\sqrt{p(1-p)/n}$ will include P between its limits.

g. The superintendent of a school district wishes to take a poll one month before a city election to determine the chances of the proposed school bond receiving a majority of the votes that will be cast. The hypothesis to be tested at the .01 level of significance is that P, the proportion of the 25,000 registered voters favoring the school bond, is .50; the alternative hypothesis is that P is not .50. A random sample of n=100 registered voters is draw; upon questioning, f=42 voters indicated that they favored the school bond.

The value of p is f/n = .42. The value of z in Eq. (14.26) is

$$z = \frac{p - a}{\sqrt{a(1 - a)/n}} = \frac{.42 - .50}{\sqrt{.50(.50)/100}} = -1.60.$$

The z value of -1.60 is compared with the critical values  $_{.003}z = -2.58$  and  $_{.003}z = -2.58$ . It cannot be concluded that p = .42 is significantly different from .50 at the .01 level. In fact, even if the .05 level of significance had been adopted, evidence would not exist for rejecting  $H_0$ : P = .50 since the critical values of z are -1.64 and 1.64.

The 99% confidence interval on P is found (from Table P) to extend from approximately .28 to .55.

#### 14.13 INFERENCES ABOUT P<sub>1</sub> -- P<sub>2</sub> USING INDEPENDENT SAMPLES

a. There are two large populations 1 and 2 in which the proportions of persons (or more generally, "units") possessing a certain characteristic are  $P_1$  and  $P_2$ . The hypothesis to be tested is that  $P_1 = P_2$  against the alternative hypothesis that  $P_1 \neq P_2$ :

$$H_0: P_1 = P_2$$
  
 $H_1: P_1 \neq P_2$ 

For example, one might test the null hypothesis that the proportion  $P_1$  of students receiving vocational counseling who plan to attend college is equal to the proportion  $P_2$  of students not receiving vocational counseling who plan to attend college.

- b. All that need be assumed is that a random sample of size  $n_1$  is drawn from population 1 and that an independent random sample of size  $n_2$  is drawn from population 2.
- c. The number of persons in the sample from population 1 possessing the characteristic being observed is  $f_1$  and the proportion is  $p_1 = f_1/n_1$ . In the sample from population 2,  $f_2$  possess the characteristic in question (planning to attend college, say) and the proportion is  $p_2 = f_2/n_2$ . The

following test statistic is defined:

$$z = \frac{p_1 - p_2}{\sqrt{\left(\frac{f_1 + f_2}{n_1 + n_2}\right)\left(1 - \frac{f_1 + f_2}{n_1 + n_2}\right)\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}.$$
 (14.27)

The quantity  $(f_1 + f_2)/(n_1 + n_2)$  is the proportion of persons in both samples 1 and 2 that possess the characteristic of interest. If the null hypothesis is true, then  $P_1$  and  $P_2$  equal a common value P that is estimated by  $(f_1 + f_2)/(n_1 + n_2)$ . And  $(f_1 + f_2)/(n_1 + n_3)$  multiplied by I minus the same quantity is an estimate of the variance of X, a dichotomously scored variable with mean P. Hence, z in Eq. (14.27) bears a resemblance to

$$t = (\bar{X}_{.1} - \bar{X}_{.2}) / \sqrt{s_x^2 \left(\frac{1}{n_1} + \frac{1}{n_2}\right)}$$

that would be used to test the hypothesis that  $\mu_1 = \mu_2$ .

- d. If  $H_0$ :  $P_1 = P_2$  is true and if, for both populations 1 and 2,  $n_t P_1$  [or  $n_t (1 \sim P_1)$  whichever is smaller] and  $n_t P_2$  [or  $n_t (1 \sim P_2)$  whichever is smaller] are greater than 5, then z in Eq. (14.27) has a normal distribution with mean 0 and standard deviation 1 over repeated pairs of independent samples.
- e. To test  $H_0$  against  $H_1$  at the  $\alpha$ -level of significance, the single calculated value of z in Eq. (14.27) is compared with  $\alpha/2^{\alpha}$  and  $\alpha/2$  in Eq. (14.27) percentiles in the unit normal distribution.
- f. For large values of  $n_1$  and  $n_2$  (both equal to 100 or more, perhaps), the  $100(1-\alpha)\%$  confidence interval on  $P_1-P_2$  is given approximately by

$$(p_1 - p_2) \pm {}_{1 - (a/2)} \mathcal{I} \sqrt{\left(\frac{f_1 + f_2}{n_1 + n_2}\right) \left(1 - \frac{f_1 + f_2}{n_1 + n_2}\right) \left(\frac{1}{n_1} + \frac{1}{n_2}\right)}. \quad (14.28)$$

g. A group of 200 students is randomly divided into two groups of 100 students each. Students in sample I are required to study instructional materials in which the concept of transitivity of the relationship "taller than" is first stated verbally then followed by several examples. In the instructional materials for sample 2, the examples are given first and are followed by the verbal statement of the concept. Underlying these two samples are hypothetical populations I and 2 of students who could have been selected to participate in the experiment. After studying the instructional materials, students in both samples are given one test item to determine whether they have mastered the transitivity concept. We wish to test whether the proportions P<sub>3</sub> and P<sub>2</sub> of students in the hypothetical populations who

326

master the concept (as evidenced by correctly answering the test item) are

equal.

Suppose we choose  $\alpha$  to be .05 and that at the end of the experiment  $f_1 = 68$  of the students in sample 1 mastered the concept and  $f_2 = 54$  of the students in sample 2 mastered it. The value of z in Eq. (14.27) is

$$z = \frac{(68/100) - (54/100)}{\sqrt{\left(\frac{68 + 54}{100 + 100}\right)\left(1 - \frac{68 + 54}{100 + 100}\right)\left(\frac{1}{100} + \frac{1}{100}\right)}} = 2.03.$$

A z value of 2.03 exceeds the critical value of  $_{.978}z=1.96$  and, thus, constitutes evidence to reject  $H_6$ :  $P_1=P_2$ . We can conclude in this hypothetical example that stating the concept first and then presenting examples of it is superior to the reverse order.

h. Appropriate techniques for making inferences concerning a set of J population proportions (in this section J=2) exist. For example, one could be interested in the null hypothesis that in a large urban school system the proportion of Negroes, Puerto Ricans, and Orientals (J=3) who leave school before graduation is the same. For a discussion of these techniques, see Marascuilo (1966) first and then Goodman (1964).

## 14.14 INFERENCES ABOUT P<sub>1</sub> — P<sub>2</sub> USING DEPENDENT SAMPLES

- a. The hypotheses to be tested are identical to those in Sec. 14.13, namely,  $H_0: P_1 = P_2$  against  $H_1: P_1 \neq P_2$ .
- b. Two random samples both of size n are drawn from populations 1 and 2, respectively. In contrast to the procedures in Sec. 14.13, we do not require the two samples to be independent. Thus samples 1 and 2 could comprase matched pairs, twin mates, "before" and "after" observations, etc. The most frequent application of the procedures in this section are to problems in which samples 1 and 2 are the same group of persons observed at two different points in time. Thus, it has sometimes been referred to as a significance test of "change," but we shall see that this interpretation is not entirely correct.
  - c. As with all techniques involving dependent samples, it is possible to establish "pairs" of observations, one member of each pair from sample 1 and the other member from sample 2. Without losing any generality, we shall consider samples 1 and 2 to be observations of n persons at times 1 and




FIG. 14.5

2, respectively. The number of persons scoring 1 on the dichotomous variable, i.e., having the characteristic being observed, in sample 1 is  $f_2$  and  $p_2 = f_1/n$ . In sample 2,  $f_2$  persons have the characteristic, and  $p_2 = f_2/n$ . It is also necessary to determine the number of pairs of observations in which both members of the pair (one from sample 1 and one from sample 2) score 1, i.e., possess the attribute being observed. Such data can be tabulated in a  $2 \times 2$  contingency table as shown in Fig. 14.5.

For example, b might be the number of persons out of n who possess the characteristic being observed at both times 1 and 2.

The following test statistic can be used to test  $H_0$  against  $H_1$ :

$$z = \frac{d-a}{\sqrt{1+a}}. (14.29)$$

d. When  $H_0: P_1 = P_2$  is true, z in Eq. (14.29) is approximately normally distributed with mean 0 and standard deviation 1, provided that d+a is as large as 10.

e. The critical values against which z in Eq. (14.29) is compared in testing  $H_0$  at the  $\alpha$ -level of significance are a/2z and 1-(a/2)z.

f. The question of placing a confidence interval on  $P_1 - P_2$  will not be discussed.

g. A sample of n=60 persons is asked to indicate whether or not they approve of capital punishment both before and after being exposed to a persuasive lecture on the abolition of capital punishment. The sample of 60 pre-lecture responses to the question on capital punishment constitutes sample 1; the 60 post-lecture responses constitute sample 2. The data obtained are tabulated in Fig. 14.6.

As an example of how the table in Fig. 14.6 is interpreted, one sees that 26 persons approved of capital punishment before hearing the lecture and disapproved after the lecture.

The null hypothesis that  $P_1 = P_2$  will be tested at the .05 level of signifi-

28

cance. The value of z in Eq. (14.29) is

$$z = \frac{d-a}{\sqrt{d+a}} = \frac{10-26}{\sqrt{10+26}} = \frac{-16}{6} = -2.67.$$

A z value of -2.67 lies considerably below the critical value of .025 = -1.96. (Note that d+a is far greater than 10.) Hence, the hypothesis that in the populations sampled (persons who had not heard an abolition of capital punishment lecture and persons who had) the proportions endorsing capital punishment are equal can be rejected at the .05 level of significance.

		Sample 1			
		Pre-			
	_	Approve	Disapprove		
	다 를 Disapprove	a=26	b=8	34	
	S Sample S Approve	c=16	d ≈ 10	26	
FIG. 14.6		42	18	n ≈ 60	

h. The test outlined above must not be considered equivalent to testing the significance of "change." Notice that in the table in Fig. 14.7 the overwhelming changes taking place are apparent even though the test statistic in Eq. (14.29) is equal to zero.

$$z = \frac{d-a}{\sqrt{d+a}} = 0.$$

The significance test of this section is a test of the significance of the difference between  $p_i=f_i ln$  and  $p_z=f_i ln$  and not of a hypothesis about "change." Change is evaluated by the significance test presented here only as it may be reflected in the difference between  $P_i$  and  $P_i$ .

The techniques presented in this section are due to McNemar (1947).

Time 1

Agree Disagree

Disagree

40 20 60 = f.

Time 2

Agree D 40 40

40 60 = f. 
$$(100 = n)$$

FIG. 14.7

# 14.15 INFERENCES ABOUT THE INDEPENDENCE OF CLASSIFICATIONS IN A CONTINGENCY TABLE

Often observations are made that permit nominal measurement of each unit (person, group, etc.) with respect to two modes of classification. For example, students can be classified both with respect to sex (male-female) and academic major (humanities, social sciences, physical sciences, engineering, or education). Schools can be classified with respect to whether they are public or private and whether they offer an industrial arts curriculum or not.

Suppose that 120 persons were classified with respect to their political affiliation and their sex. The frequencies of the various combinations of attributes could be tabulated in a contingency table as shown in Fig. 14.8. As an example of how the table is interpreted, one sees that 14 of the 120 persons sampled are female Democrats.

In general, we can consider a contingency table with I rows and J columns. We shall denote the frequency of observations in the cell at the intersection of the ith row and jth column by  $f_{ii}$ . For example, in Fig. 14.9 the number of persons in the cell in row 1 and column 1 is  $f_{1i}$ .

Note that in Fig. 14.8 the value of  $f_{11} = 29$ ,  $f_{12} = 36$ ,  $f_{1.} = 80$ ,  $f_{3} = 17$ , and  $f_{11} = 120$ .

a. The null hypothesis to be tested is that the two modes of classification upon which the contingency table is based are independent. This notion of independence requires some elaboration.

nucremaence requires some cases of studying the relationship between sex Consider again the problem of studying the relationship between sex and political affiliation in the population of all adults in a particular community. We shall denote the proportions of males in the population by  $P_{-1}$  and the proportion of females by  $P_{-2}$ . The proportions of Democrats,

	Political affiliation			
	Democrat	Republican	Independent	
Male Sex Female	29	36	15	80
	14	24	2	40
	43	60	17	120

FIG. 148 Contingency table showing the relationship between sex and political affiliation.

	1	, C1a	ssificatio	n 2	J	Row totals
١,	fss		fy	\	h	4
:	   :		:		<u>:</u>	]:
Classification i	fn		$t_{ij}$		$I_{i,j}$	<u>}</u> f <sub>0</sub> ,
1 .	:	T -	:		:	]:
	f <sub>f1</sub>	1	r <sub>n</sub>		f <sub>1,3</sub>	14.
Column totals	f3		$t_d$		1,,	f = n FIG. 14.9

Republicans, and Independents in the population will be denoted by  $P_1$ ,  $P_2$ , and  $P_3$ , respectively. If a sample of size n is drawn strictly at random from the population, these proportions can be interpreted as probabilities of obtaining males, females, or any one of the three political affiliations.

It was seen in Chapter 10 that the probability of the joint occurrence of two independent events is the product of their separate probabilities. For example, if sex and political affiliation are independent of each other, then the probability that an adult randomly selected from the population is a female Democrat is  $P_aP_a$ , if the probability of randomly drawing a male Republican is  $P_aP_a$ , if the two classifications are independent. Testing the null hypothesis of independence is equivalent to testing the hypothesis that the probability of drawing a person who falls into cell ij of the contingency table is equal to the product of the probability that the person belongs to any cell in row i and the probability that he belongs to any cell in column j:

$$H_0: P_{ij} = P_i P_{.i}$$
, for all values of i and j.

The alternative hypothesis,  $H_{10}$  is that  $P_{10} = P_1 P_2$  is not true for at least one of the H cells in the contingency table,

b. All that is assumed is that a random sample of size n is drawn from the population in question.

c. The following test statistic is used to test  $H_0$  against  $H_1$ :

$$\chi^{2} = n \left[ \sum_{i=1}^{I} \sum_{j=1}^{J} \frac{f_{ij}^{2}}{f_{i}f_{j}} - 1 \right], \tag{14.30}$$

where  $f_{ij}$  is the number of observations in the (ij)th cell of the contingency table,

f, is the number of observations in the ith row of the table,

 $f_j$  is the number of observations in the jth column of the table, and  $n = f_{ij}$ , the total number of observations.

d. When the null hypothesis of independence of the two classifications is true, the statistic  $\chi^2$  in Eq. (14.30) has a chi-square distribution with (I-1)(J-1) degrees of freedom over repeated random samples of size n from the population. For example, if sex and political affiliation are independent in the population from which the sample in Fig. 14.8 was drawn, then  $\chi^2$  for the data in that figure should appear to be a typical observation from a chi-square distribution with (I-1)(J-1)=(2-1)(3-1)=2 degrees of freedom.

If  $H_0$  is false,  $\chi^2$  in Eq. (14.30) will tend to be larger than the  $\chi^2_{(I-1)(J-1)}$  distribution. In other words, nonindependence can be expected to produce large values of  $\chi^2$  in Eq. (14.30).

- e. To test  $H_0$  at the  $\alpha$ -level of significance, the single computed value of  $\chi^2$  in Eq. (14.30) is compared with the  $100(1-\alpha)$  percentile point in the chi-square distribution with (I-1)(I-1) degrees of freedom, i.e., in the distribution  $\chi^2_{(I-D)(I-D)}$ . This percentile point is denoted by  $_{1-\alpha}\chi^2_{(I-D)(I-D)}$ . Selected percentile points in the chi-square distributions are presented in Table C of Appendix A.
- f. The hypothesis tested is one involving several parameters (population proportions) and the question of interval estimation of some function of these does not arise.
- g. The data in Fig. 14.8 will be used as an illustration. A random sample of n=120 is drawn from the population of adults in a particular community. Each person is classified with respect to sex and political affiliation. The hypothesis of independence of the two modes of classification will be tested at the .01 level of significance.

Substituting the data in Fig. 14.8 into the formula for  $z^2$  in Eq. (14.30) yields the following value:

$$\chi^2 = 120 \left( \frac{29^2}{80 \cdot 43} + \frac{36^2}{80 \cdot 60} + \frac{15^2}{80 \cdot 17} + \frac{14^2}{40 \cdot 43} + \frac{24^2}{40 \cdot 60} + \frac{2^2}{40 \cdot 17} - 1 \right)$$

= 4.776.

The obtained value of  $\chi^2 = 4.776$  is compared with the critical value of  $\kappa k_L^2 = 9.210$ . The null hypothesis of independence cannot be rejected at the .01 level. However, the probability of obtaining a value of  $\chi^2$  as large as 4.776 or larger when the null hypothesis is true is slightly less than .10. The evidence for an association other than chance between sex and political affiliation is rather weak.

We shall consider a second illustration in which I=J=2. A sample was drawn at random from the population of first-year graduate students in several large universities by the Committee on the Undergraduate Program

 $f_{.1} = 176$   $f_{.2} = 83$  |  $f_{..} = 259$  FIG. 14.

in Mathematics of the Mathematical Association of America. The n=259 students in either psychology or sociology were classified as having either "no credit" or "some credit" in undergraduate mathematics. The contingency table shown in Fig. 14.10 was obtained.

The null hypothesis to be tested is that there is no association between academic area and undergraduate mathematics training. In a  $2 \times 2$  contingency table, the formula for  $\chi^2$  in Eq. (14.30) can be simplified considerably:

$$\chi^2 = \frac{n(f_{11}f_{22} - f_{12}f_{21})^2}{f_{11}f_{21}f_{11}f_{12}}.$$
 (14.31)

The value of  $\chi^2$  with (I-1)(I-1)=(2-1)(2-1)=1 degree of freedom for the data on mathematics training is found by Eq. (14.31) as follows:

$$\chi^2 = \frac{259(25 \cdot 49 - 34 \cdot 151)^2}{59 \cdot 200 \cdot 176 \cdot 83} = 22.96.$$

From Table C in Appendix A it is seen that the probability of obtaining a 2<sup>\*</sup> as large as 22.96 or larger is less than .001 if academic area (psychology vs. sociology) and undergraduate mathematics training (no credit vs. some credit) are independent. We can conclude that there is a tendency for first-year psychology graduate students to have more undergraduate mathematics training than first-year sociology graduates students.

h. The discussion in this section of the chi-square test of independence of the classifications in a contingency table is sketchy in the extreme. Space permits only the mention of several important considerations in the application of this technique and the presentation of references in which they are examined.

When the number of observations in any cell of a  $2 \times 2$  contingency table is small (less than 10), use of a correction to the formula for  $\chi^2$  in Eq. (14.31) due to Yates (1934) is advisable. The purpose of Yates's continuity  $\chi^2$  to the chi-send stribution of with one degree of freedom. For discussions of Yates's correction in textbooks in applied statistics see McNemar (1962), Ferguson (1966), or Hays (1963), among others.

There exists a conceptually distinct interpretation of the chi-square test of this section that is appropriate when the persons in the different rows (or columns) of the contingency table can be regarded as sampled from separate and distinct populations. The chi-square contingency table test presented above may then be regarded as a test of homogeneous populations. For a discussion of the chi-square test of homogeneous populations, see Keeping (1962) and Guenther (1965).

Techniques exist whereby the hypothesis of independence of three modes of classification in a three-way contingency table may be tested. See Tate and Clelland (1957).

The problem of determining which subset of cells in a contingency table contributes to a significant  $\chi^2$  statistic has been dealt with by Marascuilo (1966).

Finally, a valuable series of papers on the use and misuse of the chisquare contingency table test appears in the Psychological Bulletin: Lewis and Burke (1949), Edwards (1950), Lewis and Burke (1950), Pastore (1950), Peters (1950). Burke (1951).

# 14.16 RELATIONSHIP BETWEEN INTERVAL ESTIMATION AND HYPOTHESIS TESTING

A relationship exists in most instances between methods of interval estimation and hypothesis testing that allows one to determine from inspection of a 100(1 - α)% confidence interval what the results would be of a hypothesis test at the α-level of significance. For example, if the 95% confidence interval around  $\bar{X}$  on  $\mu$  includes 0, then the hypothesis  $H_0$ :  $\mu = 0$  cannot be rejected at the .05 level. If the  $100(1-\alpha)\%$  confidence interval around r on  $\rho$ includes 0, then the hypothesis  $H_a$ :  $\rho = 0$  cannot be rejected at the  $\alpha$ -level of significance. If the  $100(1-\alpha)\%$  confidence interval around  $\vec{X}_{.1}-\vec{X}_{.2}$  does not include 0, then the hypothesis  $H_0$ :  $\mu_1 - \mu_2 = 0$  can be rejected at the  $\alpha$ -level. In general, all values along a particular  $100(1-\alpha)\%$  confidence interval would lead to acceptance at the a-level of the null hypothesis that the parameter being estimated was equal to any one of those values. Conversely, any value outside the confidence interval would lead to rejection of the hypothesis that the parameter in question was equal to any one of those values. For example, if the 99% confidence interval around a particular  $\bar{X}$  on  $\mu$  extended from -4.86 to 8.41, then the same data would lead to rejection of any hypothesis that  $\mu$  was any number less than -4.86 or greater than 8.41; if it had been hypothesized that  $\mu$  was 0, the data would lead to acceptance of this hypothesis at the .01 level.

Suppose that the 95% confidence interval on  $\mu$  is being established

334

SELECTED INFERENTIAL TECHNIQUES

around  $\bar{X}$ , by means of Eq. (14.2). Suppose further that the entire confidence interval is above zero. Hence, the lower limit of the confidence interval, namely  $\vec{X}_1 = \frac{1}{1975}t_{n-1}s_x/\sqrt{n}$ , is greater than zero:

$$\vec{X}_{\cdot} - \frac{s_x}{\sqrt{n}} > 0.$$

It follows that

$$\bar{X}_{\cdot} > .575 t_{n-1} \frac{s_r}{\sqrt{n}}$$

and that

$$\frac{\overline{X}_{\cdot}}{s_{\cdot}/\sqrt{n}} > .975t_{n-1}.$$

The above inequality states that the test statistic [see Eq. (14.1)] for testing the hypothesis that  $\mu=0$  exceeds the 97.5 percentile in the t-distribution with n-1 df. Hence, the hypothesis that  $\mu=0$  can be rejected at the 1-.95=.05 level. Thus, the confidence interval contained the information necessary to make a hypothesis test as well.

#### PROBLEMS AND EXERCISES

1. Herman (1967) found Minnesota Teacher Attitude Inventory scores for a sample of 14 athletes and 28 nonathletes. His findings are summarized below:

Athletes	Nonathletes		
$n_1 = 14$ $X_{.1} = 116.00$ $s_1 = 31.11$	$n_2 = 28$ $X_{.2} = 119.54$		

Test the null hypothesis at the .01 level that in the populations of athletes and nonathletes sampled  $\mu_1$  equals  $\mu_2$ .

2. Samuels (1967) performed an experiment to determine if the presence of pictures facilitated or interfered with young children's learning of words. Twenty pre-first-grade children were randomly assigned to either learn words which were illustrated with simple pictures or learn the same words without pictures. After several learning trials each child was tested on his knowledge of the words taught. The number of correct responses on the test trials for each group had the following means and standard deviations: Nounter

Nonpicture group	Picture group		
$n = 10$ $X_{.1} = 19.20$ $s_1 = 7.93$	$n = 10$ $X_{.2} = 11.30$ $s_2 = 5.79$		

Test the null hypothesis at the .05 level of significance that the two groups can be considered to be random samples from two populations with the same mean.

- 3. Herman (1967) studied the relationship between grade-point averages (X) in professional education courses and scores on the Minnesota Teacher Attitude Inventory (Y) on a sample of 42 prospective physical education teachers. An r<sub>xy</sub> of +.19 was obtained. Construct the 99% confidence interval around this value of r<sub>xy</sub> on p<sub>xy</sub>. Does the interval include zero?
- Yamamoto (1967) correlated IQ as measured by the Lorge-Thorndike Intelligence Test and creativity as measured by the Minnesota Tests of Creative Thinking on a sample of n = 75 inth-grade pupils. An r<sub>xy</sub> of 1.2 was obtained.
  - a. Test the hypothesis  $H_0$ :  $\rho_{m} = 0$  at the .05 level of significance.
  - b. Yamamoto also obtained a correlation between IQ and creativity of -.01 for n = 84 eleventh-grade pupils. Test the hypothesis that the population correlation coefficients between IQ and creativity are equal in the populations of ninth-grade and eleventh-grade pupils sampled. Test this hypothesis at the .05 level of significance.
- 5. Wallen and Campbell (1967) intercorrelated scores on the Miller Analogies Test (X), the Wide-Range Vocabulary Test (Y), and the Lorge-Thorndike Picture Reasoning Test (Z) on a sample of 60 graduate students. The following sample values were obtained:

$$r_{max} = .58$$
  $r_{max} = .17$   $r_{max} = .10$ 

Test the hypothesis at the .05 level of significance that  $\rho_{xy}$  equals  $\rho_{xx}$ , i.e., the hypothesis that the Miller Analogies Test is as closely related to a vocabulary test as it is to a nonverbal reasoning test. Interpret the results

6. Thalberg (1967) intercorrelated intelligence (X), reading rate (Y), and reading comprehension (Z) in a sample of n = 80 college students. The following correlation coefficients were obtained:

$$\frac{X}{X}$$
  $\frac{Y}{X}$   $r_{xy} = -.034$   $r_{xz} = .422$   $r_{yz} = -.385$ 

Test the null hypothesis at the .05 level of significance that  $\rho_{xy} = \rho_{xz}$ , i.e., that intelligence is correlated with both reading rate and reading comprehension to the same degree.

- 7. North and Buchanan (1967) asked a sample of 69 teachers (29 Caucasian and 40 Negro) whether they perceived "poor children" to be generally responsible or irresponsible. A phi-coefficient of correlation was calculated between the race of the teacher, X, and the teacher's perception of "poor children," X. A value of \$\phi = 2.4\$ seemed to indicate a weak relationship between the variables; Caucasian teachers tended to perceive "poor children" as irresponsible more than did Negro teachers. Using the techniques of Sec. 14.11, test the hypothesis at the .10 level of significance that in the population of teachers sampled "teacher's race" and "perception of "poor children" as irresponsible" are uncorrelated.
- Stennet (1967) sampled kindergarten pupils from rural Minnesota. In a random sample of n<sub>1</sub> = 873 boys, the proportion who were absent from school 20 or

## 15

## THE ONE-FACTOR ANALYSIS OF VARIANCE— FIXED EFFECTS

#### 15.1 LAYOUT OF DATA

As an example of a one-factor experiment, assume that a researcher wants to know if four methods of studying a topic differ in their efficiency. He plans to let 10 students study the English monetary system by outlining the topic. A different 10 students will study by writing a summary of the topic; 10 other students will study a programmed textbook of the same material; the remaining 10 students will study the programmed material on a teaching machine. The four study conditions (treatments) represent increasing observable activity by the learner. The one factor in this experiment is "learner activity"; the factor is a concept that relates the four conditions. The treatments of the experiment are different types or amounts of that thing called the factor. A factor in an experiment might be "size of type," and the three treatments comprising this factor could be a passage printed in 6 point, 8 point, and 10 point type. The researcher wants to know if at least wo of the study conditions produce a different amount of achievement (as frieasured by a multiple-choice test on English money). 378

Treatment				
_1_	2	3	4	
X1,1	X1,2	X1,3	X <sub>1,4</sub>	
X2,1	X2,2	X2,3	X2,4	
		•		
1.	•	•	• )	
١.	•	•	٠ ا	
X <sub>10,1</sub>	X10,2	X10.3	X10.4	

FIG. 15.1 Layout of data from an experiment comparing four levels of learner activity.

The data, multiple-choice test scores of the 40 students, can be tabulated as in Fig. 15.1.

You may wonder why the researcher doesn't simply look at the sums of the four treatments and see if any two are different. He does not do this because his attention doesn't focus only on the scores obtained for these particular persons at this particular time. What good would his results be if all he could say was that this group of 10 students did better than this other group on September 11 at this place? Although he has data on these 40 students taken at one particular time, if his experiment is to contribute any knowledge to science, his interest must focus on the population of students from which these 40 are only a sample and on the population of experiments of which this is only one. The researcher seeks to answer the question, "Could I expect these same results if I had chosen a different 40 subjects and run the experiment at different time under slightly different conditions?"

#### 15,2 A MODEL FOR THE DATA

To answer this question is to make an inference to a population of students and a population of performances of the experiment. The researcher uses the methods of inferential statistics to answer the question. These methods are powerful tools for experimentation. Like many "servants" of man, however, they ask that certain concessions be made to them. The methods presented in this chapter are appropriate in making inferences to a hypothetical population of trials of the experiment if the population of scores, from which the 40 scores obtained are considered to have been sampled, has a certain form and the sampling is done in a certain way. We may at times speak as though the researcher samples a person when in fact the person's score on the test is that thing assumed to have been sampled from a certain population. As an example of the type of demands inferential statistical methods will make, the researcher must assume that he has drawn the 40 obtained scores at random (i.e., with equal and independent probabilities of



being chosen) from four (one for each level of the treatment) normal distributions of scores with equal variances of. These assumptions may appear quite restrictive; it will be pointed out later (Sec. 15.13) that certain violations of the assumptions have little effect on the results of the statistical analysis. Nevertheless, the assumptions should be remembered;

- 1. The scores were sampled at random
- 2. from normal populations
- 3. with equal variances σ<sup>2</sup>,
- 4. and the different samples (four in our example) are independent.

Notice that no assumption is made about the mean  $\mu$  of the normal populations. It will be seen that the researcher's question about the efficiency of the four methods will become a question about the four means,  $\mu_1, \ldots, \mu_k$ , of the four normal populations,

We must now phrase the question about the efficiencies of the four methods in more precise terms. Toward this end, we assume that any one of the 40 scores can be represented by a linear model, the sum of components (none of which has to be squared, cubed, etc.). The linear model is a decomposition of a score  $X_{ij}$  into a sum of terms that have a certain meaning to the statistician. We shall "tentatively entertain" the following linear model for the  $X_{ij}$ 's:

$$X_{ij} = \mu + \alpha_j + e_{ij}, \tag{15.1}$$

where Xii is the ith score in the jth group,

 $\mu$  is a term (equal to the average of the four population means) that is constant for all 40 scores and reflects the overall elevation of the scores.

- $\alpha_j$  is a constant for all 10 scores in group j and reflects the elevation (or depression) in these 10 scores that results from the persons' being treated as they were (all 10 persons in group j were treated alike), and
- alike), and e<sub>ij</sub> is the "error" in the linear model. It is what remains to make up a person's score after  $\mu$  and  $\alpha_i$  have been taken out.

You must remember that  $\mu$ ,  $\alpha_1$ , and  $e_{i1}$  are numbers. If in an experiment  $\mu$  is 12,  $\alpha_2$  is -2, and  $e_{12}$  is 1, then  $X_{12}$  is 12-2+1=11.

You might expect that interest focuses on the  $\alpha_2$ 's. The term  $\mu$  is of little interest, and little can be done about the  $\alpha_2$ 's; but  $\alpha_2$  reflects what effect level j of the treatment had on the scores. The researcher's original question, "Do the four methods differ?" can now be stated more precisely: its into it tue that  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4$ ? He wishes to know if he can safely say that the four levels of the treatment do not have the same effect. The methods to be developed will be appropriate for testing the hypothesis,

called the *null hypothesis* (because there are *no* differences), that  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4$ .

One needs a method of testing this hypothesis because such things as the  $e_{ij}$ 's exist. If there were no error in the linear model, the researcher could simply look at the sums of the scores in the four groups to see if they were different and thus test his null hypothesis. But the ear's are there and they must be dealt with. How do the e, 's arise? They come about in various ways. First, persons or whatever is measured are inherently different even when they are treated alike. Of the 10 students who outline the English money material, some will score higher on the test than others simply because they are more intelligent. No amount of control over the conditions of study would result in all 10 students achieving the same degree of knowledge, Second, errors arise when an attempt is made to measure the students. These errors are due to unreliability of the measuring instrument, a multiplechoice test in this instance. A group of persons will not all earn the same test score today that they will tomorrow even if there were no forgetting. Third, errors arise from all manner of uncontrolled happenings during the experiment. A student may break his pencil and have to obtain a new one; someone may feel ill and not do well; someone else may not like the researcher's look and act uncooperatively. Additional error arises if the postulated linear model is not correct; it's possible that no linear model will give an exact description of the data.

All of these sources of error combine to make the results of the experiment today different from what would have been obtained yesterday or tomorrow on different groups of 40 students. The results of the experiment run at any one time will undoubtedly show differences between the four groups. It remains to be established, however, that the groups would be different if the experiment had been run on a different randomly chosen group of students under different conditions.

Let's go back to the original data and see how the null hypothesis that  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4$  can be tested. First we must develop some machinery.

The sum of the scores under level one is denoted by

$$\sum_{i=1}^{10} X_{i1}$$
.

The sum of the scores divided by 10, the number of scores summed, is the arithmetic mean of group one. The mean will be denoted as follows:

$$\overline{X}_{,1} := \frac{\sum\limits_{\ell=1}^{20} X_{i1}}{10}$$
.

As you'll recall the bar above the X indicates a mean; the dot in place of i means that the mean was found by summing over i. How do you find  $X_2$ ?

We also need a grand mean, the mean of all 40 scores. To find the grand mean, we sum all 40 scores and divide by 40. The grand mean is denoted by

$$\bar{X}_{..} = \frac{\sum_{i=1}^{4} \sum_{i=1}^{10} X_{ij}}{40}.$$

Once again, the bar above X indicates a *mean*; the two dots in place of the i and j indicate that both i and j have been summed over.

If one takes the difference between each score in group one and  $X_{1}$ , squares this difference, and sums the squared differences, this sum has the form

$$\sum_{i=1}^{10} (X_{i1} - \bar{X}_{,1})^2.$$

This sum divided by 9, (n-1), is the sample variance for group one and is denoted by  $s_1^*$ :

$$s_1^2 = \sum_{i=1}^{10} \frac{(X_{i1} - \bar{X}_{i1})^2}{9}$$
.

 $s_1^2$  is an unbiased estimator of  $\sigma_1^2$ , the variance of the population underlying group one. Since  $\sigma^2$  is also the variance of the three other populations,  $s_1^2$  is an unbiased estimator of this common variance  $\sigma^2$ .  $s_2^2$ ,  $s_3^2$ , and  $s_4^2$  are also unbiased estimators of  $\sigma^2$ . What does  $s_1^2$  stand for?

$$s_2^2 = \sum_{i=1}^{10} \frac{(X_{i2} - \bar{X}_{,2})^2}{9}$$
.

Now we have the machinery necessary to continue. Remember, we want to know if we can safely say that it is not true that  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4$ .

#### 15,3 ESTIMATES OF THE TERMS IN THE MODEL

What one actually observes in an experiment are the 40  $X_{ii}$ 's. The numbers  $\mu$ ,  $\alpha$ , and  $e_{ii}$  are unknown and unobservable; but certain estimates of them are possible. The statistician has found a way to estimate the components of the linear model that has very useful and desirable properties. He obtains what are called least-squares estimates of  $\mu$ ,  $\alpha$ , and  $e_{ii}$ . Denote these least-squares estimates, by  $\hat{\mu}$ ,  $\hat{\mu}$ , and  $\hat{\mu}$ . In the process of obtaining the least-squares estimates, the parameters of greatest interest,  $\alpha_1, \ldots, \alpha_g$ , are assumed to sum to zero, i.e.,  $\alpha_1 + \cdots + \alpha_r = 0$ . This is an entirely reasonable sort of restriction to adopt since we have conocived of the  $\alpha$ /5 as "elevators" or "depressors" above or below a general level that is embodied

in the parameter  $\mu$ . By placing this restriction on the  $\alpha_j$ 's, the statistician almost predestines that the  $\alpha_j$ 's will have the properties of deviations from a mean  $\mu$ , since the sum of deviations around a mean is zero. The statistician has found that

$$\hat{\mu} = \bar{X}_{..},$$

$$\hat{\alpha}_j = \bar{X}_{.j} - \bar{X}_{..},$$

$$\hat{\epsilon}_{ij} = X_{ij} - \bar{X}_{ij}.$$

The estimates are made to fit the observed data in the sense that

$$X_{it} = \hat{\mu} + \hat{\alpha}_t + \hat{e}_{tt}$$

Notice that

$$X_{ij} = \vec{X}_{ij} + (\vec{X}_{,i} - \vec{X}_{,i}) + (X_{ij} - \vec{X}_{,i}).$$
 (15.2)

The first term on the right is  $\hat{\mu}$ , the second term is  $\hat{\alpha}_j$ , and the last term is  $\hat{e}_{ij}$ .

#### J5.4

#### SUMS OF SQUARES

The motivation for the steps that follow won't be clear until much later in this chapter; let's proceed to do some arbitrary-looking manipulations on Eq. (15.2).

First, subtract the grand mean X, from each side of Eq. (15.2):

$$X_{ij} \sim \overline{X}_{ij} = (\overline{X}_{.j} \sim \overline{X}_{.j}) + (X_{ij} - \overline{X}_{.j}).$$

Second, square each side of the above equation and sum both sides over j and i:

$$\sum_{i=1}^{4} \sum_{j=1}^{10} (X_{ij} - \bar{X}_{j})^{2} = \sum_{j=1}^{4} \sum_{j=1}^{10} [(\bar{X}_{ij} - \bar{X}_{j}) + (X_{ij} - \bar{X}_{j})]^{2}.$$
 (15.3)

The quantity on the left in Eq. (15.3) is called the total sum of squares. For the entire set of numbers obtained in an experiment, the sum of the squared deviation of each number from the grand mean is the total sum of squares.

Let's consider the right side of Eq. (15.3). Notice that if we let a stand for  $(X_i - \bar{X}_i)$  and b stand for  $(X_i - \bar{X}_i)$  then the right side involves the quantity  $(a+b)^2$ . You already know that  $(a+b)^2 = a^2 + 2ab + b^2$ . So it's easy to show that

The right side of this equation is the same as

$$\sum_{j=1}^{4} \sum_{i=1}^{10} (\bar{X}_{,j} - \bar{X}_{,i})^2 + 2 \sum_{j=1}^{4} \sum_{i=1}^{10} (\bar{X}_{,j} - \bar{X}_{,i}) (X_{ij} - \bar{X}_{,i}) + \sum_{j=1}^{4} \sum_{i=1}^{10} (X_{ij} - \bar{X}_{,i})^2.$$
(15.4)

Consider for a moment only the middle term in Eq. (15.4). Since the quantity  $\overline{X}_{ij} - \overline{X}_{ii}$  has no i subscript, the  $\sum$  sign for i can be moved past it; thus we may write the second term as follows:

$$2\sum_{j=1}^{4} (\bar{X}_{,j} - \bar{X}_{,i}) \sum_{j=1}^{10} (X_{ij} - \bar{X}_{,j}). \tag{15.5}$$

Say for the moment that we are in group one and that we want to find

$$\sum_{i=1}^{10} (X_{i1} - \bar{X}_{,1}).$$

For this group,  $\bar{X}_{1}$  is a constant; using Rule 2 of Sec. 2.5, then, we can show that

$$\sum_{i=1}^{10} (X_{i1} - \overline{X}_{i1}) = \sum_{i=1}^{10} X_{i1} - 10\overline{X}_{i1} = \sum_{i=1}^{10} X_{i1} - \frac{10\sum_{i=1}^{10} X_{i1}}{10} = 0.$$

Going back to Eq. (15.5), we now see that

$$2 \sum_{j=1}^{4} (\bar{X}_{,j} - \bar{X}_{,j}) \sum_{i=1}^{10} (X_{,i} - \bar{X}_{,i})$$

$$= 2[(\bar{X}_{,i} - \bar{X}_{,i}) \cdot 0 + \dots + (\bar{X}_{,i} - \bar{X}_{,i}) \cdot 0] = 0.$$

Since the middle term of Eq. (15.4) is exactly zero—this is not an assumption, we have proved it, and it will be true for any set of numbers—we have shown that

$$\sum_{j=1}^{4} \sum_{i=1}^{10} (X_{ij} - \bar{X}_{..})^2 = \sum_{j=1}^{4} \sum_{i=1}^{10} (\bar{X}_{.j} - \bar{X}_{..})^2 + \sum_{j=1}^{4} \sum_{i=1}^{10} (X_{ij} - \bar{X}_{.j})^2. \quad (15.6)$$

Notice that in the first term on the right side of Eq. (15.6) no subscript i appears. The quantity  $(X_A - \bar{X}_A)^2$  is the same for all i from one to 10 when j has been fixed. For example, for the first, second, ..., tenth scores in group one,  $(\bar{X}_A - \bar{X}_A)^2$  is the same. Therefore,

$$\sum_{j=1}^{4} \sum_{i=1}^{10} (\bar{X}_{,i} - \bar{X}_{,i})^2 = \sum_{j=1}^{4} 10 \cdot (\bar{X}_{,j} - \bar{X}_{,i})^2.$$

Finally, we have broken the total sum of squares into two parts:

$$\sum_{j=1}^{4} \sum_{i=1}^{10} (X_{ij} - \bar{X}_j)^2 = \sum_{j=1}^{4} 10 \cdot (\bar{X}_j - \bar{X}_j)^2 + \sum_{i=1}^{4} \sum_{i=1}^{10} (X_{ij} - \bar{X}_i)^2.$$

$$SS_{tot} = SS_{tot} + SS_{within}$$

The total sum of squares  $(SS_{tot})$  has been partitioned (analyzed) into two additive components, the sum of squares between groups  $(SS_{tot})$  and the sum of squares within groups  $(SS_{tot})$ .  $SS_{tot}$  reflects variation in the scores obtained. This total variation has been analyzed into two components,  $SS_{tot}$  and  $SS_{within}$ . Thus the name "analysis of variance," (abbreviated ANOVA). You will soon see how this analysis is used to test the null hypothesis that  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4$ .

# IS.5 RESTATEMENT OF THE NULL HYPOTHESIS IN TERMS OF POPULATION MEANS

Before proceeding, let's phrase the null hypothesis that  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4$  in an equivalent but slightly different form. The estimator of  $\alpha_1$  was taken to be  $\hat{\alpha}_1 = X_1 - X_0$ . Since  $\hat{\alpha}_1$  is an unbiased estimator of  $\alpha_1$ , the expectation E (long-term average) of  $\hat{\alpha}_1$  is  $\alpha_2$ :

$$E(\hat{\alpha}_1) = E(\vec{X}_{.1} - \vec{X}_{..}) = E(\vec{X}_{.1}) - E(\vec{X}_{..}) = \mu_1 - \mu = \alpha_1.$$

Similarly,  $\alpha_2 = \mu_2 - \mu$ ,  $\alpha_3 = \mu_3 - \mu$ , and  $\alpha_4 = \mu_4 - \mu$ .

The null hypothesis can be written as

$$(\mu_1 - \mu) = (\mu_2 - \mu) = (\mu_3 - \mu) = (\mu_4 - \mu).$$

Add  $\mu$  to each of the four terms and you will see that this is obviously the same as saying that  $\mu_1 = \mu_2 = \mu_3 = \mu_4$ . Hence, the null hypothesis that  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4$  in the linear model  $X_{ij} = \mu + \alpha_j + e_{ij}$  is the same as the hypothesis that the means of the normal populations from which the samples are drawn are all equal, i.e.,

$$u_1 = u_2 = u_3 = u_4$$

We may state the null hypothesis in either form. A third equivalent form, when  $n_1 = n_2 = n_3 = n_4$  is

$$H_0: \sum_{j=1}^4 (\mu_j - \bar{\mu}_j)^2 = 0,$$

where  $\bar{\mu}$  is the average of  $\mu_1, \ldots, \mu_J$  and equals  $\mu$ .

#### 15.6 DEGREES OF FREEDOM

Some more machinery is needed before we can show how the null hypothesis can be tested.

We must associate with each quantity in the partitioning of the total sum of squares  $(SS_{tot}, SS_{bet}, SS_{within})$  an integer called the degrees of freedom.

"Degrees of freedom" is a name borrowed from the physical sciences where it denotes a characteristic of the movement of an object. If an object is free to move in a straight line only, it has one degree of freedom; an object that is free to move through any point in a plane, such as a bowling ball rolling down an alley, has two degrees of freedom; a ball in a handball court has three degrees of freedom; it can go from back to front, side to side, and floor to ceiling. You may be surprised to learn that the techniques we group under the rubric "analysis of variance" or ANOVA (the partitioning of SStot and testing the null hypothesis) have a geometric interpretation. The name "degrees of freedom" enters into analysis of variance (ANOVA) by way of its geometric interpretation.

First, let us discuss the degrees of freedom associated with SSpet-Consider the definition of SS,...:

$$SS_{bet} = \sum_{j=1}^{4} 10 \cdot (\bar{X}_{.j} - \bar{X}_{.j})^2$$

The four group means,  $X_1, \ldots, X_d$ , are related to  $X_n$  by the equation:

$$\frac{\bar{X}_{,1} + \bar{X}_{,2} + \bar{X}_{,3} + \bar{X}_{,4}}{4} = \bar{X}_{,1}. \tag{15.7}$$

If  $\vec{X}_{..}=6$ ,  $\vec{X}_{.1}=3$ ,  $\vec{X}_{.2}=4$ , and  $\vec{X}_{.3}=8$ , what must  $\vec{X}_{.4}$  be?  $\vec{X}_{.4}$  must be 9. If  $X_{..} = 4$ ,  $X_{.2} = 3$ ,  $X_{.3} = 6$ , and  $X_{.4} = 4$ , what must  $X_{.1}$  be?  $X_{.1}$  must be 3. If  $X_n$  is a given number, then we are free to assign any values whatsoever to three of the group means, but having done so the last group mean is determined—it must have a value that satisfies Eq. (15.7). The degrees of freedom for  $SS_{bet}$  is (4-1)=3, one less than the number of group means, i.e., J-1. We will abbreviate "degrees of freedom for  $SS_{bet}$ " to  $df_{bet}$ .

Consider now the degrees of freedom associated with SS within-

$$SS_{within} = \sum_{j=1}^{4} \sum_{i=1}^{10} (X_{ij} - \bar{X}_{,j})^2$$
.

For group one, the computation of  $SS_{within}$  involves

$$(X_{11} - X_{.1})^2 + (X_{21} - X_{.1})^2 + \dots + (X_{10,1} - X_{.1})^2$$

How are  $X_{11}, X_{21}, \ldots, X_{10,1}$  and  $\overline{X}_{.1}$  related?

$$\frac{X_{11}+X_{21}+\ldots+X_{10}}{10}=\bar{X}_{.1}.$$

If  $X_{.1}$  were a fixed number, say 12.40, to how many of the 10 quantities  $X_{11}$  through  $X_{10,1}$  could you assign any number you wished before you have to assign a value or values to make  $X_{.1}$  be 12.40? The answer is nine.

$$(X_{12} - \overline{X}_{.2})^2 + (X_{22} - \overline{X}_{.2})^2 + \dots + (X_{10,2} - \overline{X}_{.2})^2$$

is used in computing  $SS_{within}$ . Since  $(X_{12} + X_{22} + \ldots + X_{10,2})/10 = X_{2}$ , if  $X_{.2}$  is fixed, then nine of the 10 single numbers can be freely assigned before the tenth number must be assigned that gives the proper  $X_{.2}$ . There are nine degrees of freedom for each of the four groups in the computation of  $SS_{within}$ ; hence the degrees of freedom for  $SS_{within}$  equals

$$(10-1) + (10-1) + (10-1) + (10-1) = 40-4 = 36,$$

the total number of observations less the number of groups, i.e., Jn-J=J(n-1). Space can be saved if "degrees of freedom for  $SS_{within}$ " is denoted by  $df_{within}$ .

SStot equals

$$(X_{11} - \bar{X}_{..})^2 + (X_{21} - \bar{X}_{..})^2 + ... + (X_{10,4} - \bar{X}_{..})^2;$$

41 terms are involved, and they are related as follows:

$$\frac{X_{11}+X_{21}+\ldots+X_{10,4}}{40}=\vec{X}_{..}.$$

Of the 40 quantities on the left of the equation, 39 can be assigned numbers without restriction before the fortieth one must be given that number that yields the preassigned  $\bar{X}_n$ . The degrees of freedom for SS<sub>tot</sub> equals 40-1=39, the total number of observations minus 1, i.e., Jn-1.

## 15.7 MEAN SQUARES

A sum of squares (SS) divided by its degrees of freedom (df) is called a mean square (MS). In the one-factor ANOVA, only two mean squares will be of interest: the mean square between,  $MS_s$ , which equals  $SS_s|df_s$ , and the mean square within,  $MS_s$ , which equals  $SS_s|df_s$ . A "mean square total" will not be defined because it would prove useless for the purpose of testing the null hypothesis. However, an important relationship to keep in mind is  $SS_s + SS_s = SS_s$ ; another is  $df_s + df_w = df_t = Jn - 1$ .

# 15.8 EXPECTATIONS OF MS, AND MS,

At this point, hopefully, you'll start finding the answers to some of the questions which must have arisen in your mind while studying the last few

In the remainder of this chapter "between" will be abbreviated to b, "within" to w, and "total" to t. Thus we shall write SS<sub>s</sub>, SS<sub>s</sub>, SS<sub>s</sub>, df<sub>s</sub>, df<sub>s</sub>, MS<sub>s</sub>, and MS<sub>s</sub>.

pages. Why partition  $SS_i$  into  $SS_a$  and  $SS_a$ ? Why define degrees of freedom and mean squares? To test the null hypothesis  $H_a$ , of course; but by looking at the expectations of  $MS_a$  and  $MS_a$ , you'll begin to see how the machinery developed so far relates to  $H_a$ .

The expectation of  $MS_w$  means the long-term average of  $MS_w$  from experiment to experiment. The expectation of  $MS_w$  will be denoted by  $E(MS_w)$ . If in our learner-activity experiment we were to repeat the experiment an indefinitely large number of times and each time compute  $MS_w$ . then the average of all of these  $MS_w$  would be  $E(MS_w)$ . We can conceive of doing this even though it would be impossible to do so physically.  $E(MS_w)$  is the average of all of the  $MS_w$  in that population of experiments about which we wish to make an inference from the data obtained this one time

We can write  $E(MS_{\bullet})$  in terms of a characteristic of the normal populations from which the scores obtained in the experiment are a random sample.

First, let's look at MS, in a slightly different way.

$$\begin{split} MS_w &= \frac{SS_w}{df_w} = \frac{\sum_{i=1}^{\frac{1}{2}} (X_{i1} - \bar{X}_{,i})^2}{4 \cdot (10 - 1)} \\ &= \frac{1}{4} \left[ \frac{\sum_{i=1}^{\frac{1}{2}} (X_{i1} - \bar{X}_{,i})^2}{9} + \frac{\sum_{i=1}^{\frac{1}{2}} (X_{i2} - \bar{X}_{,2})^3}{9} + \frac{\sum_{i=1}^{\frac{1}{2}} (X_{i3} - \bar{X}_{,i})^2}{9} + \frac{\sum_{i=1}^{\frac{1}{2}} (X_{i4} - \bar{X}_{,i})^2}{9} \right]. \end{split}$$

 $\sin(\sum_{n=1}^{2n}(X_n-X_n)^n/9)$  the sample variance of group one? We shall denote it by  $x_1^n$  and the sample variances of groups two, three, and four by  $x_2^n$ ,  $x_3^n$ , and  $x_3^n$ , respectively.

Therefore,  $MS_{\omega} = \{(s_1^* + s_2^* + s_3^* + s_3^*), i.e., MS_{\omega}$  is the average of the sample variances of all the groups. We stated earlier that  $E(s_1^*)$  was  $a_1^*$ , the variance of the normal population from which the scores in group one were sampled. Since all of the normal populations had the same variance,  $a_1^*$ , we can argue that

$$E(MS_w) = \frac{1}{4}E(s_1^2 + \dots + s_t^2) = \frac{1}{4}[E(s_1^2) + \dots + E(s_t^2)]$$
  
=  $\frac{1}{4}(\sigma^2 + \sigma^2 + \sigma^2) = \sigma^2$ 

The expectation of MS, is at.

The size of  $MS_{\phi}$  does not depend on the means of the populations underlying the groups in the experiment.  $MS_{\phi}$  is "mean-free," reflecting only the variability among the measures within groups. Such variability is

349

around each group mean, rather than around the grand mean of all the groups. Whether all of the groups are samples from the same normal population or all the normal populations have different means, E(MS,) will be the same.  $\sigma^{2}$  \*

The same cannot be said for  $E(MS_b)$ , however. We shall show that if all of the normal populations underlying the samples in the experiment have the same mean (we've already assumed they have the same variance), then  $E(MS_b)$  is  $\sigma^2$ . If, however, at least two of the population means are different. then  $E(MS_b)$  will be larger than  $\sigma^2$ . If all the population means are equal. then the null hypothesis that  $\mu_1 = \mu_2 = \mu_3 = \mu_4$  is true. If at least two population means are different, e.g.,  $\mu_1 = \mu_2$  but  $\mu_3 \neq \mu_4$  or  $\mu_1 \neq \mu_2 \neq \mu_4$  $\mu_3 \neq \mu_4$  (these are two of many possible examples), then the null hypothesis Ho is false.

We shall state without proof that

$$E(MS_b) = \sigma^2 + \frac{n \sum_{j=1}^{J} (\mu_j - \mu)^2}{I - 1},$$
 (15.8)

where  $\sigma^2$  is the variance in each population,

n is the number of subjects in each group (10 in our example),

J is the number of groups (four in our example),

μ, is the population mean of the jth population, and u is the mean of the J population means.

Let us say for example that all four population means were 6.45. (This is a case where  $H_0$  is true.) Then  $\mu$  would be 6.45 and

$$\sum_{i=1}^{J} (\mu_i - \mu)^2 \approx 0.$$

Take an example where  $H_0$  is not true and satisfy yourself that  $E(MS_n)$ is greater than σ2.

To summarize:

1. If Ha is true, then

$$E(MS_w) = \sigma^2$$
 and  $E(MS_b) = \sigma^2$ .

\* Convince yourself that this is indeed true. Start with

$$E(MS_w) = E[(SS_{w_1} + SS_{w_2} + ... + SS_{w_j})/J(n-1)].$$

For each  $SS_{v_i}$  substitute its equivalent,  $(n-1)MS_{v_i}$ . Then distribute the expectations and simplify. Further, note that the definition of  $SS_{\pi_j}$  is  $\sum_{i=1}^{n} (X_{ii} - X_{ij})^n$ , which does not involve the overall mean,  $X_{ii}$ . 2. If Ho is false, then

$$E(MS_w) = \sigma^2 \quad \text{and} \quad E(MS_b) = \sigma^2 + \frac{n\sum\limits_{j=1}^J (\mu_j - \mu_j)^2}{J - 1} \; .$$

which is greater than  $\sigma^2$ .

In any experiment,  $MS_a$  and  $MS_b$  are known, but the values of their expectations are not. By comparing  $MS_b$  to  $MS_a$ , one can determine if  $H_0$  is plausible. If  $MS_b$  is very large relative to  $MS_a$ , then it is likely that  $H_0$  is false. But the test of the null hypothesis is not as simple as this. It can be shown that  $MS_b$  and  $MS_b$  are independent of one another (this is one reason for the original assumption of normal populations).

Suppose for the moment that you draw four samples at random of 10 scores each from a normal population. This is exactly what would be done if in our example experiment four conditions were equally efficient (in terms of the average score they produced). The 40 scores sampled could be placed in a table such as Table 15.1, and MS, and MS, could be computed. Remember this is a case of Ho being true. Would MS, and MS, be equal? No. Their expectations would be equal; but from one sampling to the next, MS, might be a little bigger than MS, this time, and maybe a lot smaller next time. The sample value of MS, and MS, will fluctuate independently from one sampling of 40 scores to the next (the experiment that was actually run is considered to be one case of sampling 40 scores). The cases that vex the researcher are those in which  $MS_b$  is large relative to  $MS_a$ . He would like to regard such an occurrence as evidence that  $H_0$  is false. (If  $H_0$  is false, he would expect MS, to be larger than MS...) But how can he be sure that a large value of  $MS_b$  relative to  $MS_a$  didn't occur simply by random fluctuations in scores sampled from the same normal population? The researcher can never be certain, but we shall see in the next section how he can control the proportion of times he will incorrectly conclude that  $H_0$  is false.

# 15.9 SOME DISTRIBUTION THEORY

We saw in Chapter 11 that a chi-square variable with one degree of freedom has the form

$$\frac{(X-\mu)^2}{\sigma^2} \sim \chi_1^2,$$

i.e., the squared normal deviate is distributed as chi square with 1 degree of

Recall that X is a normally distributed variable with population mean  $\mu$  and variance  $\sigma^2$ .

Suppose one randomly samples n scores from a normal distribution with mean and variance  $\mu$  and  $\sigma^2$ . Since chi-square variables have the special additive property noted in Chapter 11, the quantity

$$\frac{(X_1 - \mu)^2}{\sigma^2} + \frac{(X_2 - \mu)^2}{\sigma^2} + \ldots + \frac{(X_n - \mu)^2}{\sigma^2} \sim \chi_n^2$$

The sum of the squared z scores has a chi-square distribution with n degrees of freedom, i.e., this sum computed on repeated samples of n scores has a known frequency distribution  $z_n^2$ .

We shall not prove the assertion; however, it is true (see, e.g., Wilks, 1962) that if  $X_1, \dots, X_n$  are n independent observations from a normal population with variance  $\sigma_1^n$ , then

$$\frac{(X_1 - \bar{X})^2}{\sigma^2} + \frac{(X_2 - \bar{X})^2}{\sigma^2} + \ldots + \frac{(X_n - \bar{X})^2}{\sigma^2} \sim \chi_{n-1}^2.$$

Therefore, 
$$\sum_{i=1}^{n} (X_i - \bar{X}_i)^2 / \sigma^2 \sim \chi_{n-1}^2$$
. Also,  $\sum_{i=1}^{n} (X_{i,i} - \bar{X}_{i,i})^2 / \sigma^2 \sim \chi_{n-1}^2$ .

Notice that  $\bar{X}$ , the sample mean of the n observations, has replaced  $\mu$ , the mean of the population from which the observations were selected. Also, the chi-square variable has n-1 instead of n degrees of freedom.

For the experiment at the beginning of the chapter we defined SSw:

$$SS_w = \sum_{i=1}^{10} (X_{i1} - \bar{X}_{i1})^2 + \sum_{i=1}^{10} (X_{i2} - \bar{X}_{i2})^2 + \dots + \sum_{i=1}^{10} (X_{i4} - \bar{X}_{i4})^2.$$

If we divided the first quantity on the right of this equation by  $\sigma^2$ , it would have the distribution  $\chi_1^2$  (chi-square with 9 degrees of freedom). This is so because we assumed the scores in group one were randomly drawn from a normal population with variance  $\sigma^2$ . The same can be said for the other three quantities on the right of the equation. Since all four quantities are distributed as  $\chi_2^2$ , then their sum divided by  $\sigma^2$  is distributed as  $\chi_{10+2+3+9}^2$ , or  $\chi_2^2$  (chi-square with 36 degrees of freedom). Hence, for our example

$$\frac{SS_w}{3} \sim \chi_{36}^2$$

We can divide SS, by 36 to obtain MS, then

$$\frac{MS_{10}}{\sigma^2} \sim \frac{\chi_{16}^2}{36}.$$

This fact will be saved for future use.

In general, how is the mean  $\bar{X}$ , of a group distributed? That is, what frequency distribution would result if we sampled n scores at random from a population, computed  $\bar{X}$ , then recorded only this one score,  $\bar{X}$ ? What

would the distribution of these scores be like if this process of sampling, computing  $\vec{X}$ , and recording it would continue indefinitely? What would be the mean, variance, and shape of this distribution?

If the means,  $\bar{X}$ 's, are based on n scores randomly drawn from a normal population, they will:

- 1. be normally distributed themselves,
- have mean μ, and
- have variance σ<sup>2</sup>/n.

Using the above facts, we can see that  $(\bar{X}_1 - \mu)/\sqrt{\sigma^2/n}$  has a mean of zero and a variance of 1; hence, it is a z score. Consequently

$$\frac{(\bar{X}_{.} - \mu)^{2}}{\sigma^{2}/n} = \frac{n(\bar{X}_{.} - \mu)^{2}}{\sigma^{2}} \sim \chi_{1}^{2}.$$

If the null hypothesis is true, then

$$\frac{\sum_{j=1}^{4} 10(\bar{X}_{,j} - \bar{X}_{,j})^2}{\sigma^2} \sim \chi_3^2,$$
 (15.9)

because under that condition  $E(X_a) = E(X_a) = E(X_a) = E(X_d) =$ 

$$\frac{\left[\sum_{j=1}^{4} 10(\bar{X}_{,j} - \bar{X}_{,j})^{2}\right]/3}{\sigma^{2}} = \frac{MS_{b}}{\sigma^{2}} \sim \frac{\chi_{3}^{2}}{3}.$$

provided Ho, the null hypothesis, is true.

You'll remember from Chapter II that the ratio of two chi-square variables each of which is divided by its degrees of freedom has an F-distribution.

$$\frac{\left[\sum_{i=1}^{4} 10(\bar{X}_{ii} - \bar{X}_{i})^{2}\right]/3\sigma^{2}}{\left[\sum_{i=1}^{4} \sum_{i=1}^{10} (X_{ii} - \bar{X}_{i})^{2}\right]/36\sigma^{2}} = \frac{\left[\sum_{i=1}^{4} 10(\bar{X}_{ij} - \bar{X}_{j})^{2}\right]/3}{\left[\sum_{i=1}^{4} \sum_{i=1}^{10} (X_{ii} - \bar{X}_{i})^{2}\right]/36} \sim F_{1.34}.$$

Notice that since of appears in both numerator and denominator it cancels

You should recognize that the numerator is MS, and the denominator

is MS... To summarize then, if the null hypothesis is true.

$$\frac{MS_b}{MS} \sim F_{3,36},$$

i.e., the ratio of the mean-square between,  $MS_b$ , to the mean-square within,  $MS_w$ , has an F-distribution with 3 and 36 degrees of freedom when  $\mu_1 = \mu_2 = \mu_3 = \mu_4$ .


The ratio  $MS_b/MS_w$  is called the *F-ratio*. The *F*-ratio is the statistic that will be used in the last steps of the test of  $H_0$ .

# 15.10 THE F-TEST OF THE NULL HYPOTHESIS: RATIONALE AND PROCEDURE

To coalesce the distribution theory that has been developed, consider the act of repeatedly sampling at random four groups of 10 scores each from the same normal distribution (note that under these conditions the null hypothesis is true). If each time this were done the F-ratio  $MS_b/MS_c$  were calculated and its value recorded, the frequency distribution of these F-ratios (many, many of them) would look like the mathematical curve  $F_{3.98}$ . This knowledge is extremely important, for with it the statistician can calculate the percent of F-ratios that will exceed 11.35 or 6.12; or he can determine the number that will be exceeded only by 5% or 1% of the F-ratios. He does this assuming  $H_0$  is true. The F-ratios obtained in this procedure of repeatedly sampling, calculating, and recording would all be greater than zero and the largest would be infinitely large, theoretically.

What if the sampling of the four groups of 10 scores each was not done from the same normal population, i.e., what if sampling was done under the condition of a false null hypothesis? We saw earlier that under these conditions one expects  $MS_v$  to be larger than it would be if the null hypothesis were true. However,  $MS_v$  would have the same expectation,  $\sigma^2$ . When the null hypothesis is false,  $MS_v$  does not have a  $\chi^2$  distribution and the ratios  $MS_v MS_v$  obtained from the sampling do not have an F-distribution. We do know, however, what effect this sampling under conditions of a false null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the distribution of the F-ratios. They will be on the null hypothesis has on the first hypothesis has on the first hypothesis has one had not have an expectation of the first hypothesis has not have an expectation of the first had hypothesis had not have an expectation of the first had hypothesis had not have an expectation of the first hypothesis had hypothesis hypothesis had hypothesis hypothesis had hypothesis hypothesi

Suppose we randomly sampled repeatedly four groups of 10 scores each from one normal population and calculated the F-ratio each time. The plot of the F-ratios obtained would look like the curve  $F_{3,36}$  in Fig. 15.2. Now suppose we drew repeatedly two groups of 10 scores each from a normal



(curve  $F_{3.14}$ ) and when  $H_{9}$  is false (curve  $F^{9}$ ).

population with mean  $\mu_1$ , and two more groups of 10 scores from a normal population with a different mean  $\mu_2$ . The null hypothesis is false, but we wouldn't in practice know this because  $\mu_1$  and  $\mu_2$  are unknown characteristics of populations. The distribution of the F-ratios obtained would look something like curve  $F^*$  in Fig. 15.2. Notice that in general the F-ratios shift to the right (are larger) when sampling is done under conditions of a false null hypothesis instead of a true null hypothesis. Even so, some of the values in the curve  $F_{3.20}$  are larger than some of the values in curve  $F^*$ .

Is an F-ratio greater than 3.25 more likely to occur if one is sampling under conditions of a true or false null hypothesis? Compare the areas under the two curves to the right of the point 3.25 and see which area is larger (i.e., shows a greater probability of yielding a value of F greater than 3.25).

The statistician can find the percentile points of the curve  $F_{3,28}$ . He can calculate the value of F such that only 5% or 1% of the values in the curve  $F_{3,28}$  rear larger, i.e., he can find the 95th and 99 herecretiles of the curve  $F_{3,28}$ . The 95th percentile in the curve  $F_{3,28}$  is 2.86; the 99th percentile is 4.38. Table E of Appendix A shows the 75th, 90th, 95th, 97.5th, 99th, 99.5th, and 99.9th percentiles for various F-distributions. As we saw in for the numerator and the degrees of freedom for the denominator of the F-ratio. To find the 95th percentile in the curve  $F_{2,28}$  find the cell at the intersection of column 2 and row 40 for that portion of the table headed 95th percentile. The value of  ${}_{3}F_{3,28}$  is 3.23.

What if our experiment was run and an F-ratio of 6.51 was obtained with 3 and 36 degrees of freedom? Since the 95th percentile of the curve  $F_{3.86}$ , is 2.86, and under the null hypothesis the F-ratios form the curve  $F_{3.86}$ , if the null hypothesis were true, something very unlikely would have occurred. Less than five times in 100 a value as large or larger than 2.86 would be obtained if the null hypothesis were true. If the null hypothesis were false,

then large values of F are more likely to be observed. The researcher agrees to reason thus:

If the value of the F-ratio obtained would occur less than five times in 100 (i.e., if it is greater than the 95th percentile of  $F_{3,20}$ ) if the null hypothesis were true, then l'Il conclude that the null hypothesis is false. It seems more likely that such a value indicates a false null hypothesis since large values are more likely if the null hypothesis is false.

Choosing the 95th percentile of the curve  $F_{3.26}$  as the point on which the decision about  $H_6$  hinges is rather arbitrary. One could have chosen the 90th, 99th or 99.9th percentile point. What if the 50th percentile point had been chosen? That is, what if one had agreed to reject the null hypothesis if the F-ratio was greater than the 50th percentile of  $F_{3.26}$ ? If the null hypothesis were true, one would have a probability of  $\frac{1}{2}$  of rejecting  $H_6$  (calling it false). If researchers agreed on the 50th percentile point to make their decisions, half of them would be concluding that one method is better than another when the methods were actually no different. Scientists want to guard against these mistakes, type I errors; so they agree to conclude that the null hypothesis is false when values equal to or greater than the value of F obtained have a small probability of occurring when the null hypothesis is true. A "small probability" means .10, .05, or .01. These values correspond to using the 90th, 95th, and 99th percentile points, respectively.

Do not make the mistake of thinking that if the F-ratio obtained in our experiment were 6.51 the null hypothesis is certainly false. Such assertions are not possible in our situation. A researcher makes a conclusion of the form "I reject  $H_0$  as a true statement about the means of the populations if V resampled "or "I fall to reject  $H_0$  as . . ." He is never certain of the truth of his conclusion. He does know, however, that his conclusions will be correct a certain percent (90%, 95%, 99%, etc.) of the times he makes them given the truth or falsity of  $H_0$ .

Suppose that the four learner activity levels were all alike with respect to the amount achieved under each. Suppose also that our researcher ran the same experiment, 40 subjects into four groups, a very large number of times. Each time he would compute an F-ratio and draw a conclusion about  $H_0$ . He is not aware that  $H_0$  is true. The curve  $F_{3.26}$  that the obtained F-ratios would follow has a known form. We know, for example, that 5% of the area under the curve lies to the right of the point 2.86. Our researcher has agreed that if the F-ratio he obtains exceeds 2.86, he will say that  $H_0$  is false. He will expect 5% of the F-ratios he obtains to be greater than 2.86; is false, he will be expected to make the mistake of rejecting  $H_0$  when it is really true 5% of the time.

Notice that the value (2.86) the researcher chooses, such that any F-ratio exceeding that number is regarded by him as evidence for the falsity  $H_0$ , is arbitrary. If he had chosen 4.38, the 99th percentile point in the

356

distribution  $F_{3.36}$ , and if  $H_0$  is true, then on the average one time in 100 he will obtain an F-ratio exceeding 4.38 and conclude incorrectly that  $H_0$  is false. The probability that the researcher will reject  $H_0$  as the true state of nature when  $H_0$  is actually true is called  $\alpha$  (alpha). The size of  $\alpha$  is under the researcher's control; he can make it as large or as small as he wishes by his selection of a number from the F-table that will govern his decision.

It is customary to assume (as we saw in Sec. 13.5) that there are two states of nature that can exist relative to our ANOVA model: Ho can be true, or it can be false. A researcher agrees to make one of two conclusions after inspecting his data: Reject Ho as explaining my situation, or do not reject (continue to entertain) Ho as an explanation of my situation. Four outcomes of an experiment are possible, as shown in the accompanying table

	State of nature	
	Ho is true	$H_0$ is false
Reject H <sub>O</sub> Do not reject H <sub>O</sub>	Type I error	No error is made
	No error is mode	Type II error

If the researcher does not find evidence for the falsity of  $H_0$  and  $H_0$  is true (this situation is the lower left-hand cell in the table), then his thoughts will be in accord with the true state of nature: no error in his conclusion has been made. If the researcher rejects  $H_0$  when in reality  $H_0$  is false (upper right-hand cell), he has not committed an error in making his conclusion. However, if he rejects Ho when it is true, we say he has made a "type I error" or an "error of the first kind." If  $H_0$  is actually false, and the researcher fails to reject  $H_0$ , we say he has committed a "type II error" or an "error of the second kind." These names are merely a convenience; that one is a type I error and the other a type II error is of no significance.

The probability of committing a type I error is equal to  $\alpha$ . The researcher can control the size of  $\alpha$ , thus making his chances of incorrectly rejecting  $H_0$  very large ( $\alpha=.25$  or .30) or very small ( $\alpha=.01$  or .001). It is customary to assign a value to  $\alpha$  of .05 or .01; although .05 and .01 may be appropriate for industrial or agricultural research in which the ANOVA first enjoyed popularity, there is little reason to advocate their use exclusively in educational and psychological research. The value of  $\alpha$  a researcher wishes to use should depend on certain aspects of his particular analysis. Values of z = .15 or .10 might be justifiable if a small number of subjects are included in the experiment. As will be shown later, the size of  $\alpha$  is related in an indirect way to the probability of making a type II error (failing to reject a false null hypothesis). The probability of failing to reject a false




FIG. 15.3 Illustration of the probabilities of type I and type II errors for J=4, n=10, and a particular alternative hypothesis.

null hypothesis is under the control of the researcher to some extent.  $\beta$  depends on the number of subjects in the experiment and the value of  $\alpha$  chosen, among other things. We shall denote the probability of correctly rejecting  $H_0$  when it is false by some given amount by  $1-\beta$ . So the probability of type II error is  $\beta$ .  $1-\beta$  is the probability of rejecting  $H_0$  when it is false; it is called the "power" of the test. The larger  $\alpha$ , the larger it is false; it is called the "power" of the test. The larger  $\alpha$ , the larger  $1-\beta$ . The value of  $1-\beta$  is quite important to the researcher. All too often, people completely disregard "power." They invest large amounts of time and money in an experiment in which their probability of discovering differences between the treatments is only .2, say. That is, even if differences of a given magnitude exist, the researcher has two chances in 10 of rejecting  $H_0$  ("discovering" that differences exist). There's a high probability that he won't find what he's looking for even when it is there.

It may happen that if a researcher set  $\alpha$  at .05 the corresponding value of  $1-\beta$  would be .20; but if  $\alpha$  were .10,  $1-\beta$  would be .50. Obviously, the prudent course of action is to accept a greater risk of making a type I error to increase the probability,  $1-\beta$ , of finding differences between the treatments.

The definitions of  $\alpha$  and  $\beta$  and their relationship are depicted in Fig. 15.3. The calculation of the power of the F-test of  $H_0$  will be illustrated in Sec. 15.15.

# 15.11 THE ONE-WAY ANOVA WITH n OBSERVATIONS PER CELL (SUMMARY)

An experiment with one treatment factor and J levels of that factor is run. Under each of the J levels, n independent observations are taken. The researcher is willing to assume that the n observations in any one level are

independent observations from a normal population with variance  $\sigma^t$ . The variance  $\sigma^t$  is assumed to be equal in all J of the levels. (This assumption is often called the assumption of homoscedasticity; the word "homoscedasticity" derives from two Greek words meaning "same" and" scatter" or "dispersion.")

The ANOVA follows five steps:

- 1. A linear model is postulated to explain the data.  $X_{ij} \leftrightarrow \mu + \alpha_j + \epsilon_{ij}$ , where  $\alpha_1 + \ldots + \alpha_J = 0$  and the  $\epsilon_{ij}$  are all independent.
- 2. The null hypothesis  $H_0$  and an alternative hypothesis  $H_1$  are stated.  $H_0: \alpha_1 = \alpha_2 = \ldots = \alpha_J$ ;  $H_1$ : "At least two  $\alpha_J$ 's are different." Or equivalently

$$H_0: \mu_1 = \ldots = \mu_J$$
  $H_1: \sum_{i=1}^{J} (\mu_i - \hat{\mu}_i)^i \neq 0.$ 

- 3. An a-level is selected, i.e., the researcher decides what probability he will assign to the event that he rejects H<sub>0</sub> when it is true (concludes there are differences between the levels of the treatment when there actually are none). This α-level is typically .10, .05, .01, or .001. Values of π much greater than .10 probably allow too great a risk of falsely rejecting H<sub>0</sub>, and values of π much less than .001 are probably overly conservative.
- Computations of sums of squares [parts (a) and (b) below], degrees
  of freedom [parts (c) and (d) below], and mean squares [part (e) below]
  are made.

The data from the experiment may be laid out as follows:

# 

**a.**  $SS_w = \sum_{j=1}^{J} \sum_{i=1}^{n} (X_{ij} - \bar{X}_{ij})^2$ . By algebraic manipulations,  $SS_w$  can be written in a form that is more efficient for computation:

$$SS_{w} = \sum_{i=1}^{J} \sum_{i=1}^{n} \chi_{ij}^{2} - \frac{\sum_{i=1}^{J} \left(\sum_{i=1}^{n} \chi_{ij}\right)^{2}}{n}.$$

Step 1. To calculate  $SS_{\pi}$ , square each individual observation and sum these squared numbers. You then have

$$\sum_{i=1}^{J} \sum_{i=1}^{n} X_{ij}^{2}.$$

Step 2. Find the sum of the original observations for each of the J columns. The sum of the jth column is

$$\sum_{i=1}^{n} X_{ij}.$$

Square each of the J column sums and add these J squared numbers together. Divide this sum by n. You now have the quantity

$$\sum_{j=1}^{J} \left( \sum_{i=1}^{n} X_{ij} \right)^{2}$$

Step 3. Find  $SS_w$  by subtracting the result of Step 2 from the result of Step 1.

b.  $SS_b = \sum_{j=1}^J n(\bar{X}_{.j} - \bar{X}_{.j})^2$ . Again by algebraic manipulations it may be shown that

$$SS_b = \frac{\sum_{i=1}^{J} \left(\sum_{j=1}^{n} X_{ij}\right)^2}{n} - \frac{\left(\sum_{j=1}^{J} \sum_{i=1}^{n} X_{ij}\right)^2}{Jn}.$$

Step 1. Bring forward the value

$$\sum_{j=1}^{J} \left( \sum_{i=1}^{n} X_{ij} \right)^2$$

that was calculated in Step 2 of (a).

- Step 2. Add together all of the Jn original observations and square this sum. Divide the squared sum by Jn. You now have the quantity  $\left(\sum_{i=1}^{J}\sum_{j=1}^{n}X_{ij}^{3}\right)^{j}/Jn$ .
- Step 3. Find SS<sub>b</sub> by subtracting the result of Step 2 from the result of Step 1.
  - c. The degrees of freedom associated with  $SS_w$  are J(n-1).

- d. The degrees of freedom associated with  $SS_b$  are J-1.
- e. Calculate the MS., and the MS. as follows:

$$MS_{w} = \frac{SS_{w}}{J(n-1)}.$$

$$MS_{b} = \frac{SS_{b}}{I-1}.$$

5. The F-ratio  $MS_0/MS_w$  is calculated and compared to the 100(1 —  $\alpha$ ) percentile point in the distribution  $F_{J-1,J(n-1)}$ .

The test of the null hypothesis is made by comparing the ratio  $MS_b/MS_w$ , an F-ratio with J-1 and J(n-1) degrees of freedom, with a value obtained from an F-table that is the value exceeded by a per cent of F-ratios obtained under conditions of a true null hypothesis. If  $MS_b/MS_w$  exceeds that value (denoted by  $\sum_{i=1}^{n} f_{i-1} J_{i-1} J_{i-1}$ ), then the researcher rejects  $H_a$  as a true statement-

To illustrate the above calculations, we shall return now to the example of Sec. 15.1 in which an experimenter sought to determine the relative effectiveness of four different levels of learner activity on learning about the English monetary system. Forty experimental subjects were randomly placed among the four levels (10 to each level) of learner activity: (1) outline the instructional material; (2) write a summary of the material; (3) study an instructional program of the material; (4) study the program on a teaching machine. Ten subjects studied English money under each of these conditions on five successive days; then a 100-item multiple-choice test on the English money system was given. A person's "score" was the number of item answered correctly on the 100-item test. These scores appear in Table 15.1.

TABLE 15.1 SCORES ON A 100-ITEM TEST OVER THE ENGLISH MONETARY SYSTEM FOR 40 SUBJECTS STUDYING UNDER FOUR LEVELS OF LEARNER ACTIVITY

1 Outline	2 Written summary	3 Program	4 Teaching machine
26	51	52	41
34	50	64	49
46	33	39	56
48	28	54	64
42	47	58	72
49	50	53	65
74	48	77	63
61	60	56	87
51	71	63	77
53	42	59	62

The calculations for obtaining  $SS_b$ ,  $SS_w$ ,  $MS_b$  and  $MS_w$  are performed in Table 15.2. In the upper portion of Table 15.2, the data in Table 15.1 are summarized by the necessary quantities.

TABLE 15.2 ILLUSTRATION OF CALCULATIONS FOR A ONE-FACTOR ANOVA WITH EQUAL n's: DATA FROM TABLE 15.1

ANOVA WITH EQUAL 
$$n's$$
; DATA FROM TABLE 15.1

1 2 3 4

 $n = 10$   $n = 10$   $n = 10$   $n = 10$   $n = 10$ 
 $X_{.1} = 48.40$   $X_{.3} = 48.00$   $X_{.4} = 57.50$   $X_{.4} = 63.60$ 

$$\sum_{i=1}^{10} X_{i1} = 484$$
 
$$\sum_{i=1}^{10} X_{i2} = 480$$
 
$$\sum_{i=1}^{10} X_{i3} = 57.5$$
 
$$\sum_{i=1}^{10} X_{i4} = 636$$

$$\sum_{i=1}^{10} X_{i1}^{1} = 25,024$$
 
$$\sum_{i=1}^{10} X_{i1}^{1} = 24,392$$
 
$$\sum_{i=1}^{10} X_{i2}^{1} = 33,925$$
 
$$\sum_{i=1}^{10} X_{i4}^{1} = 42,034$$

$$\sum_{i=1}^{4} \sum_{i=1}^{10} X_{i1} = 2175$$
 
$$\sum_{i=1}^{4} \sum_{i=1}^{10} X_{i2}^{1} = 125,375$$

$$SS_{b} = \frac{(484)^{5} + (480)^{5} + (575)^{2} + (636)^{5}}{40}$$

$$\approx 119,977.70 - 118,265.62 = 1712.08$$

$$SS_{w} = 125,375 - \frac{(484)^{3} + (480)^{3} + (575)^{2} + (636)^{5}}{10}$$

$$= 125,275 - 119,977.70 = 5397.30$$

$$MS_{b} = \frac{SS_{b}}{f-1} = \frac{1712.08}{3} = 570.69$$
  $MS_{w} = \frac{SS_{w}}{f(n-1)} = \frac{5397.30}{36} \approx 149.93$ 

The null hypothesis  $H_0$ , that  $\mu_1 = \mu_2 = \mu_3 = \mu_4$ , is tested by referring  $F = MS_0/MS_2$  to the F-distributions in Table E of Appendix A. The obtained value of F is located relative to various percentile points of the F-distribution with 3 and 36 degrees of freedom to determine whether it could reasonably be regarded as an observation randomly sampled from that distribution—which it must be if  $H_0$  is true.

The value of the F-ratio for the data in Tables 15.1 and 15.2 is F = 570.69/149.93 = 3.81. In Table E we can find the percentile points in  $F_{3.00}$  and  $F_{3.40}$  but not in  $F_{3.30}$  since not all F-distributions are tabulated there. We can, however, interpolate between 30 and 40 to obtain approximate values of the percentile points in  $F_{3.00}$ . The interpolation is performed with the reciprocals of the degrees of freedom instead of the degrees of freedom instead of the degrees, e.g., to find  $_{30}F_{3.30}$  one solves the following equation:

$$\frac{\frac{1}{30} - \frac{1}{36}}{\frac{1}{30} - \frac{1}{40}} = \frac{.85F_{3.30} - .95F_{3.36}}{.85F_{3.30} - .95F_{3.46}}.$$

THE ONE-FACTOR ANALYSIS OF VARIANCE—FIXED EFFECTS

The following percentile points are obtained by interpolation in Table E:

$$_{75}F_{3,36} = 1.43$$
  
 $_{50}F_{3,36} = 2.25$   
 $_{55}F_{3,36} = 2.86$   
 $F = 3.81 -$ 
obtained value of  $F = MS_b/MS_w$ 

The single obtained value of F from the data in Table 15.1 lies between the 95th and the 99th percentiles in the F-distribution with 3 and 36 degrees of freedom. In fact it lies at about the 98th percentile in the  $F_{3,36}$  distribution. Hence, if  $H_0$ :  $\mu_1 = \mu_2 = \mu_3 = \mu_4$  were true, an F-ratio as large as 3.81 or larger would occur with a probability approximately equal to .02. If we were testing  $H_0$  at the  $\alpha = .05$  level in this instance, we would reject  $H_0$  in favor of  $H_1$  in which it is held that not all four  $\mu$ 's are equal. If we adhered rigidly to the  $\alpha = .01$  level of significance we would not reject  $H_0$ . We would be inclined to proceed as though  $H_0$  were false.

## [5.12 THE ONE-WAY ANOVA WITH UNEQUAL n's

Not infrequently, the numbers of scores are unequal in the J cells of a onefactor ANOVA. For example, five persons might be observed under level 1 of the factor, and 10 and 20 persons might be observed under levels 2 and 3 respectively. The technique of analysis of variance that was developed in the preceding sections of this chapter can be easily modified to accommodate these "unequal n's designs." The theory is essentially the same; only slight modifications are made. The computational techniques are quite similar to those in the equal n's ANOVA.

# Notation

362

You are accustomed by now to calling the ith score in the jth group,  $X_{ij}$ . When we said that i runs from 1 to n and j runs from 1 to J, we knew that there were J groups, each containing n measures.

Because in the present situation the groups may contain different numbers of scores, it will be necessary to refer to the number of scores in the first group as  $n_1$ , the number of scores in the second group as  $n_2, \ldots$ , and the number of scores in the Jth group as  $n_J$ . Data gathered in a design that is to be analyzed with a one-factor unequal n's ANOVA could be depicted

Group 1	Group 2	Group 3
X <sub>1,1</sub>	X <sub>1,2</sub>	X <sub>1,3</sub>
$X_{2,1}$	$X_{2,2}$	$X_{2,3}$
•		
	•	
	•	
	$X_{25,2}$	$X_{18,3}$
X20 .		

In the above example, group 1 contains 20 scores, group 2 contains 15 scores, and group 3 contains 18 scores. Consequently,  $n_1 = 20$ ,  $n_2 = 15$ , and  $n_3 = 18$ .

#### The Model

The same assumptions about parent populations are made that were made for the equal n's ANOVA: the J samples are randomly drawn from normal populations with equal variances  $\sigma^2$ , and they are independent. As before, it is assumed that the scores  $X_{tt}$  can be thought of in terms of the following linear model:

$$X_{ij} = \mu + \alpha_j + e_{ij},$$

where Xii is the ith score in the jth group,

 $\mu$  is the average of the J population means,

 $\alpha_j$  is the difference between the mean of the jth population,  $\mu_j$ , and  $\mu_j$  and

 $e_{ij}$  is the difference between  $X_{ij}$  and  $\mu_{ij}$ , the mean of the jth population.

The subscript i runs from 1 to  $n_i$  in the jth group. We shall denote the total number of scores in all groups by  $N_i$  of course,  $N = n_1 + n_2 + ... + n_{x_i}$ .

A restriction placed on the equal n's ANOVA was that  $\alpha_1 + \ldots + \alpha_J = 0$ . In the unequal n's ANOVA this assumption becomes  $n, \alpha_1 + n, \alpha_2 + \ldots + n, \alpha_J = 0$ . This is the only substantial alteration of the theoretical model, and you should not be greatly concerned.

As before, the least-squares estimators of the treatment effects,  $\alpha_j$ , are obtained and become the basis of tests of the hypothesis that in the population all  $\alpha_j = 0$ , i.e., all J population means are equal. The least-squares estimator of  $\alpha_j$  is  $X_j - X_n$ , i.e., the mean of the  $n_j$  scores in the jth group minus the mean of all scores.

$$\hat{\alpha}_i = \bar{X}_j - \bar{X}_n = \frac{\sum\limits_{i=1}^{n_j} X_{ij}}{n_j} - \bar{X}_n.$$

Again, the least-squares estimator of  $\mu$  is simply

$$\widehat{\mu} = \widehat{X}_{..} = \frac{\sum\limits_{i=1}^{J}\sum\limits_{i=1}^{n_i} X_{ij}}{N}.$$

The estimator of the error component, ei, is given by

$$\hat{e}_{ij} = X_{ij} - \bar{X}_{ij}$$

The null hypothesis we wish to test is that in the population all of the  $\alpha$ , are equal, and hence equal zero. In symbols, the null hypothesis is as follows:

$$H_0: \alpha_j = 0$$
 for all  $j$ .

An equivalent statement of  $H_0$  in terms of the J population means is  $H_0$ :  $\mu_1 = \mu_2 = \ldots = \mu_J$ .

#### Sums of Squares

As before, the route to a statistical significance test of  $H_0$  goes by way of the sum of squared estimates of the treatment effects,  $\alpha_j$ , and the sum of squared estimates of the errors,  $e_{ij}$ .

First, a sum of squares within,  $SS_w$ , and a sum of squares between,  $SS_y$ , are found.  $SS_w$  is simply a weighted sum of the J sample variances for each group:

$$SS_{w} = (n_{1} - 1)s_{1}^{2} + (n_{2} - 1)s_{2}^{2} + \dots + (n_{J} - 1)s_{J}^{2}$$

$$= \sum_{i=1}^{n_{1}} (X_{i1} - \bar{X}_{i})^{2} + \sum_{i=1}^{n_{1}} (X_{i2} - \bar{X}_{i})^{2} + \dots + \sum_{i=1}^{n_{J}} (X_{iJ} - \bar{X}_{J})^{2}.$$

A simpler notation for the above weighted sum is:

$$SS_w = \sum_{i=1}^{J} \sum_{i=1}^{n_f} (X_{if} - \bar{X}_{,f})^2$$

Using sigma notation, SS, can be written

$$SS_w = \sum_{i=1}^{J} \sum_{i=1}^{n_i} X_{ii}^2 - \sum_{j=1}^{J} \frac{\left(\sum_{i=1}^{n_j} X_{ij}\right)^2}{n_i}.$$

In words,

- Square each of the N scores and add them up to obtain the first term.
- Sum the scores in the jth group, square this sum, and divide by n<sub>j</sub>.
  Do this for all J groups.
  - 3. Sum the I quantities obtained in step 2.

Therefore

$$E\left(\frac{SS_{\varphi}}{N-J}\right)=E(MS_{\varphi})=\sigma^{2}.$$

Whether Ho is true or false, the long-run average value of MS, (which would be obtained by averaging the MS s from a huge number of instances of running the same experiment) is equal to o2, the common variance of each of the J populations.

The value of E(MSb) is somewhat more difficult to derive, so we state without proof that the following is true:

$$E(MS_b) = \sigma^2 + \frac{\sum\limits_{1}^{J} n_j (\mu_j - \mu)^2}{J - 1}. \label{eq:energy}$$

The formula for E(MS) for the unequal n's ANOVA is quite similar to E(MS,) for the equal n's case. (See Eq. (15.8).) They differ only in that the factor n, by which all squared deviations \(\mu\_1 - \mu \) are weighted in the equal n's case, becomes a differential weight n, for each squared deviation in the unequal n's case.

The essential property of MS, is the same as when n's were equal, and this property is best revealed by inspecting  $E(MS_h)$ . When  $H_0$  is true, one expects MS, and MS, to be the same size, namely o2. When Ho is false, one expects MS, to be larger than MS. These considerations, and others concerning distribution theory quite like that in the equal n's case, provide the techniques for making the F-test of Ha.

# The F-test of Ho: Rationale and Procedure

When the null hypothesis is true the ratio MSy/MS, will follow a central F-distribution with J-1 and N-J degrees of freedom. When the null hypothesis is false, the ratio MS./MS., will follow a noncentral F-distribution, which has a larger mean than the central F-distribution and more of its area above the bulk of the area of  $F_{J-1,N-J}$  (see Fig. 15.2).

The strategy by which Ho is tested is no different from what it was in the equal n's case. One adopts an  $\alpha$ , finds the  $100(1-\alpha)$  percentile point in the table of the F-distribution which has J-1 and N-J degrees of freedom, compares MSb/MS, with 1-xFJ-1,N-J, and decides to accept or reject II. If MSJMS exceeds 1-FJ-1.N-J, then it is considered unlikely that II. is true; for to do so forces one to regard the obtained F-ratio as an unlikely event (in fact, an event with probability a or less of happening). It would make better sense to regard this large F-ratio as one of the typical values from a noncentral F-distribution that would describe the distribution of MSJMS, if H, were false.

# An Example

An experiment was performed in which the effects of perceived similarity to a social group on conforming behavior were observed. A group of 60 experimental subjects was divided at random into three groups: low similarity—subjects in this group were informed that their expressed opinions were generally at variance with those of college students in general; medium similarity—subjects were told that their opinions agreed with those of college students in general only moderately often; high similarity—subjects were told that their opinions were usually exactly like those of students in general. After subjects were thus informed, they were asked to express their opinions about 18 current issues (capital punishment, birth control, etc.). Before expressing their opinion, however, the subjects were told how students in general felt about each issue. The number of times out of 18 a subject's general felt about each issue. The number of times out of 18 a subject's expressed opinion was the same as what was portrayed to be the opinion of students in general was taken to be the number of times the subject "conformed to majority opinion." The actual "conformity scores" for the 60 subjects are presented in Table 15.3.

TABLE 15.3 CONFORMITY SCORES OF SUBJECTS WHO WERE TOLD THAT THEY HELD OPINIONS OF LOW, MEDIUM, OR HIGH SIMILARITY TO STUDENTS IN GENERAL

I STANTS	2 Medium simılarıty	Ηį	gh si	3 mılar	rity
15 13 11 9 14 13 10 9 14 13 10 9 14 13 10 8 13 13 10 8 13 13 10 8	15 12 15 11 14 11 14 10 14 10 13 10		14 14 14	12 12 12 12	

Twelve Ss were observed under the medium similarity condition; 24 were observed under low similarity and 24 under high similarity. The null hypothesis of interest is that in the populations from which these samples can be considered to have been drawn at random—these are hypothetical populations of comparable subjects participating in this same experiment—the means of the three conditions are equal. That is, if we number the groups 1, 2, and 3 from low similarity to high, then  $H_0$ :  $\mu_1 = \mu_2 = \mu_3$ . The calculation of sums of squares is performed in Table 15.4.

The degrees of freedom for  $MS_b$  and  $MS_g$  are J - 1 = 2 and N - J = the degrees of freedom for  $MS_b$  and  $MS_b$  are J - 1 = 2 and N - J = 60 - 3 = 57, respectively. Hence,  $MS_b = 20.416/2 = 10.208$  and  $MS_w = 304.167/57 = 5.336$ . The ANOVA results are tabulated as follows:

Source of variation:	df	MS	F
Between groups	2	10.208	1,91
Within groups	57	5.336	

To test the null hypothesis  $H_0$  that  $\mu_1 = \mu_2 = \mu_3$  (or equivalently that  $\frac{3}{2}z_1^2 = 0$ ), refer an obtained F-ratio of 1.91 to the F-distribution with 2 and 57 degrees of freedom. The 75th and 90th percentiles in the distribution  $F_{1,3}$ , are 1.42 and 2.40, respectively. In replications of the experiment in Table 15.3, F-ratios exceeding 2.40 will be obtained 10% of the time when the null hypothesis is true. The obtained F-ratio of 1.91 will not allow confident rejection of the null hypothesis.

TABLE 15.4 ILLUSTRATION OF CALCULATION OF SUMS OF SQUARES FOR A ONE-FACTOR ANOVA WITH UNEQUAL n's: DATA FROM TABLE 15.3

$$\frac{1}{n_1 = 24} \frac{2}{n_2 = 24}$$

$$\frac{1}{n_1 = 24} \frac{1}{n_2 = 24} \frac{1}{n_2 = 24}$$

$$\frac{1}{2} \frac{1}{1} X_{11} = 273$$

$$\frac{1}{2} \frac{1}{2} X_{12} = 149$$

$$\frac{1}{2} \frac{1}{1} X_{12} = 303$$

$$\frac{2}{1} \frac{1}{2} \frac{1}{1} \frac{1}{1} X_{12} = 715$$

$$\frac{2}{1} \frac{1}{1} \frac{1}$$

IS,I3
CONSEQUENCES OF FAILURE
TO MEET THE ANOYA
ASSUMPTIONS: THE
"ROBUSTNESS" OF ANOYA

The problem of what happens (to levels of significance and power) when the assumptions underlying an analysis of variance model are violated presents considerable difficulty to the mathematical statistician. This is to variances and normality. The actual significance levels are based on samples of 1000 t-ratios. Entire frequency distributions of the t-ratios obtained when t-tests are performed on nonormal populations with unequal variances are reproduced in the 1960 article by Boneau. These graphs are illuminating and ought to be studied. We shall give only a few of the more easily reproduced findings from Boneau's work. These results appear in Table 15.7. Try to integrate Boneau's findings with those of Box and Pearson.

In summary, the fixed effects ANOVA appears to be remarkably insensitive to departures from normality; and when n's are equal, it is equally unaffected by heterogeneous variances. Bot has used the word "robustness" for this insensitivity of a statistical test to violation of its assumptions.

# IS.14 TESTING HOMOGENEITY OF VARIANCES

Two instances in which testing for heterogeneity of population variances is worthwhile come to mind: (1) when one wishes to make inferences about population variances because they are of scientific interest, and (2) when one suspects heterogeneity of variances in an analysis of variance in which not all factors have fixed effects.

The best known test of the homogeneity of a set of variances is due to M. S. Bartlett (1937). Statisticians are sometimes reluctant to run Bartlett's test because the necessary calculations are arduous. Bartlett's test requires the calculation of the J sample variances and their logarithms. In addition to the drawback of laborious calculations, Bartlett's test has been found to be quite sensitive to violation of the necessary assumption that the J samples come from normal populations (Box, 1953). If nonnormal populations are sampled, the probability of a type I error (rejecting a true  $H_0$ ) with Bartlett's test may be far greater than the z chosen by the test user. Bartlett's test is so sensitive to the assumption of normal populations that it may even be a good test of normality (Box, 1953)! Scheffe's test of homogeneity of variances can be every bit as tedious arithmetically as Bartlett's. It is much less sensitive to violation of the normality assumption, however (Scheffé, 1959). A test devised by Hartley (see Biometrika Tables for Statisticians, pp. 60-61) is a useful short-cut test of  $H_{\bullet}$ . It consists of referring the ratio of the largest sample variance to the smallest variance to a special table (Table 31, Biometrika Tables). Hartley's test does not use all of the information the observations provide about heterogeneity of variances, and it might be expected that far more powerful tests could be found. Cochran (1951) proposed a test of  $H_0$  that is simple to run and that uses more of the information in the data than does Hartley's. Winer (1962, p. 96) maintained

that there is evidence that the tests of Bartlett, Hartley, and Cochran are sensitive to the assumption, upon which each rests, that the J samples are from normal distributions. The two latter tests also require special tables for their execution

Levene (1960) proposed a test of  $H_0$  that is both simple to run (it uses ordinary analysis of variance techniques) and insensitive to violation of the normality assumption in most cases. Levene's test is simply a one-way analysis of variance on the absolute values of the differences between each observation and the mean of its group. For example, the data from an experiment have the following lavour:

One wishes to test the hypothesis  $H_0: \sigma_2^2 = \dots = \sigma_J^2$ . No assumption about the shapes of the distributions underlying the J samples is made. A statistic will be appropriate for testing  $H_0$  on the above data provided that (1) one can approximate the 95th and 99th percentiles, say, in the distribution of the statistic when  $H_0$  is true, and (2) the test based on the statistic has good power for rejecting  $H_0$  when it is false. The test statistic for Levene's test is the ratio of  $MS_0$  to  $MS_0$  of transformed scores  $Z_{ij}$  that are related to the  $X_{ij}$ 's by  $Z_{ij} = |X_{ij} - X_{ji}|$ . If this F-ratio exceeds the  $1 - \alpha$  percentile point of the ordinary F-distribution with degrees of freedom J - 1 and J(I - 1), one concludes with approximately  $1 - \alpha$  confidence that the population variances are different.

Levene's paper (1960) reported on his investigation of the properties of the above test. Among other things, he studied the correspondence between the probability of obtaining a significant F-ratio on the  $Z_0$ 's and the nominal probability adopted by the test user. That is, the correspondence between the sampling distribution of the F-ratio on the  $Z_0$ 's and the F-distribution was examined. It was anticipated that this correspondence would be close due to the widely recognized "robustness" of the fixed-effects analysis of variance. Since the fixed-effects analysis of variance is robust for equal numbers of observations per group, Levene restricted attention to this case. The robustness of Levene's test has thus been examined only for equal n's. The mathematics for Levene's investigation proved to be intractable; thus empirical sampling distributions of F-ratios calculated on the  $Z_0$ 's were compared with the F-distribution with J-1 and J(n-1)

degrees of freedom. In general, the agreement of the nominal (based on the F-distribution) and the empirical (based on 1000 observations) probabilities of rejecting a true H. was remarkably close.

Levene also investigated the power of his test for rejecting  $H_0$ . The power was found to be generally quite satisfactory. Assuming normal distributions and J=2, the efficiency of Levene's test relative to the exact F-test was good. These efficiencies ranged between .75 and .96 for various values of  $\alpha$  and various alternative hypotheses  $(\sigma_0^2/\sigma_0^2)$ . For J samples the efficiency of Levene's test relative to the approximate efficiency of Bartlett's test, assuming normal distributions, ranged from .83 to .90.

In summary, Levene's test appears to be a "robust" test (for equal numbers of observations in J samples) of the hypothesis of equal population variances that has satisfactory power to reject alternative hypotheses. Its ease of calculation and use of standard tables of the F-distribution make it an attractive alternative for more laborious and less robust tests that have been in common use. (See Glass, 1966a.)

## 15.15 POWER OF THE F-TEST

The power of a particular F-test depends on four quantities: the degrees of freedom "between" denoted by  $n_1$ , the degrees of freedom "within" denoted by  $n_2$ , a quantity denoted by  $\phi_2$ , which is a measure of the degree of "falseness" of the null hypothesis, and  $\alpha$ , the level of significance of the test. Of course, in the one-factor ANOVA the value of  $n_1$  is J-1, and the value of  $n_2$  is N-J. The power of the F-test is defined for a particular set of values of  $\mu_1, \dots, \mu_{J^*}$ . The quantity  $\phi$  has the following definition:

$$\phi = \sqrt{\frac{\sum\limits_{j=1}^{J} n_j (\mu_j - \bar{\mu})^2}{I\sigma^2}},$$

where  $\bar{\mu} = (\mu_1 + \ldots + \mu_J)/J$ , which also equals  $\mu$ .

The calculation of  $\phi$  involves  $\sigma^2$ , the population variance common to all J populations. Normally  $\sigma^2$  will not be known; this necessitates either gathering preliminary data to estimate it, making a shrewd guess as to its value, or measuring the differences among the  $\mu$ , in  $\sigma$ -units so that  $\sigma^2$  need not be known (e.g., what is the power of the F-test if  $\mu_1 - \mu_2$  equals  $\sigma/27$ ).

Once  $\phi$ ,  $n_1$ ,  $n_2$ , and  $\alpha$  are known, the power of the f-test can be determined from Table N in Appendix A. For example, suppose that an experiment is performed comparing J=3 treatments. Assume that there are n=11 observations per group and that the f-test of the null hypothesis is to be performed at the  $\alpha=0.05$  level. We wish to calculate the power of the F-test

for the following alternative hypothesis:

$$\mu_1 = 68, \quad \mu_2 = 66, \quad \mu_3 = 64.$$

Suppose that previous experiments on similar phenomena lead one to believe that  $\sigma^2$  is reasonably close to 20. The value of  $\phi$  is

$$\phi = \sqrt{\frac{11[(68 - 66)^2 + (66 - 66)^3 + (64 - 66)^3]}{3 \cdot 20}}$$

$$= \sqrt{\frac{11(8)}{200}} = \sqrt{\frac{88}{100}} = \sqrt{1.47} = 1.21.$$

Table N is entered with  $n_1=(J-1)\approx 2$ ,  $n_2=(N-J)\approx 30$ ,  $\alpha=.05$ , and  $\phi=1.21$ . The power of the F-test of  $H_0$  against the alternative hypothesis that  $\mu_1=68$ ,  $\mu_2=66$ , and  $\mu_3=64$  is approximately 0.40. Thus, there are four chances in 10 of rejecting  $H_0$  in favor of  $H_1$  if in fact  $\mu_1=68$ ,  $\mu_2=66$ , and  $\mu_3=64$ .

Verify that for  $n_1 = J - 1 = 1$ ,  $n_2 = N - J = 60$ ,  $\alpha = .01$ , and  $\phi = 5.00$ , the power of the F-test is approximately 0.94.

#### PROBLEMS AND EXERCISES

 Calculate the degrees of freedom for both MS<sub>b</sub> and MS<sub>w</sub> in each of the following instances:

instances.  
a. 
$$J = 2$$
,  $n = 4$ .  
b.  $J = 5$ ,  $n = 2$ .  
c.  $J = 3$ ;  $n_1 = 3$ ,  $n_2 = 6$ ,  $n_3 = 4$ .  
d.  $J = 3$ ;  $n_1 = 4$ ,  $n_2 = 1$ ,  $n_3 = 5$ .

 Determine the critical value of F = MS<sub>b</sub>/MS<sub>w</sub> for testing H<sub>0</sub> in each of the following instances:

a. 
$$J=2; n=6; \alpha=.01$$
. b.  $J=5; n=7; \alpha=.10$ . c.  $J=3; n_1=4, n_2=6, n_3=8; \alpha=.05$ .

3. Which one or more of the following statements are equivalent to the alternative hypothesis H<sub>1</sub> in the one-factor ANOVA?

a. 
$$\mu_1 \neq \mu_{j^*}$$
 for some  $j$  and  $j^*$ . b.  $\mu_1 \neq \mu_2 \neq \ldots \neq \mu_{J-1} \neq \mu_{J}$ .

c. 
$$\sum_{i=1}^{J} (\mu_i - \bar{\mu})^2 \neq 0$$
. d.  $\mu_j \neq \mu_{j*}$  for all pairs of  $j$  and  $j^*$ .

 Given only the following data in an ANOVA table, determine MS<sub>b</sub>, MS<sub>w</sub>, and F.

Source of variation	df	55	MS	_F
Between groups	4	81.25		
Within groups Total	49	378.60		

5. The report of an experiment in which three treatment groups were compared contained only the following data:

$$n_1 = 10$$
  $n_2 = 10$   $n_3 = 10$   $n_3 = 10$   $n_3 = 10$   $n_3 = 10.06$   $n_3 = 102.06$   $n_3 = 105.06$   $n_3 = 105.06$ 

Using these summary statistics, calculate  $MS_0$  and  $MS_0$  and test  $H_0$ :  $\mu_1=\mu_2=\mu_3$  at the .01 level of significance.

 In Prob. 2 at the end of Chapter 14, a t-test of the significance of the difference between two means was called for on the following data:

Group 1
 Group 2

 
$$n_1 = 10$$
 $n_2 = 10$ 
 $X_{.1} = 19.20$ 
 $X_{.2} = 11.30$ 
 $s_1 = 7.93$ 
 $s_2 = 5.79$ 

The value of 1 was 2.55, which was significant at the .05 level. Perform a one-way ANOVA with J=2 on the above data and show that  $F=MS_0MS_0$  equals  $t^2=(2.55)^2$  which is also significant at the .05 level. Thus, verify for one particular case that the one-way ANOVA for two groups is equivalent to a t-test of  $H_0^2$ ,  $H_0=H_2$ .

7. Guthrie (1967) studied the effectiveness of three different modes of training on deciphering cryptograms. Group I was trained by first being presented rules for deciphering cryptograms, then working examples. Group II worked examples first then was told the rules. Group III worked only examples. A control group studied Russian vocabulary during the training period. A group of 72 subjects was randomly split into four groups of 18 each and assigned to the four training condutions. After training, the S's were given a 10-ltem test comprising 10 new cryptograms like those studied during training. The means and standard deviations of the number of cryptograms solved on the criterion test in each group are as follows:

Group I (rule-example)	Group II (example-rule)	Group III (example)	Control
$n = 18$ $\overline{X}_{.1} = 8.06$ $s_1 = 2.37$	$n = 18$ $\vec{X}_{.2} = 6.94$ $s_2 = 2.31$	$n = 18$ $X_{.3} = 7.11$ $s_3 = 2.42$	$n = 18$ $X_{.4} = 4.73$ $s_4 = 2.27$

Perform a one-way ANOVA on Guthrie's data to test the null hypothesis that the means of the populations underlying the four groups are equal. Test  $H_0$  at the .03 level of significance. (Save your calculations because they will be used again in the problems at the end of Chapter 16.)

8. Three methods of teaching foreign language vocabulary were compared in an experiment. To evaluate the instruction, a 50-ttern vocabulary test was administered to the 24 students in the experiment; eight students were in each group. The following data—expressed as number of correct items out of 50—were obtained:

Aural-oral method	Translation method	Combined methods
$X_{11} = 19$	$X_{12} = 21$	$X_{13} = 17$
$X_{21} = 37$	$X_{22} = 18$	$X_{23} = 20$
$X_{31} = 28$	$X_{32} = 15$	$X_{33} = 28$
$X_{41} = 31$	$X_{42} = 23$	$X_{43} = 30$
$X_{51} = 29$	$X_{52} = 20$	$X_{53} = 13$
$X_{61} = 25$	$X_{62} = 22$	$X_{63} = 18$
$X_{71} = 36$	$X_{72} = 26$	$X_{73} = 19$
$X_{81} = 33$	$X_{82} = 14$	$X_{83} = 23$

Perform an F-test of the null hypothesis that  $\mu_1 = \mu_2 = \mu_3$  at the .05 level of significance. (Save your results for the exercises at the end of Chapter 17.)

9. Harrington (1968) experimented with the sequencing of instructional material and "organizers" that structured the material for the learner. A group of 30 persons were randomly split into three groups of 10 each. Group I received organizing material before studying instructional materials on mathematics; group II received the "organizer" after studying the mathematics; group III received no organizing material in connection with studying the mathematics instructional materials. On a 10-tiem test over the mathematics covered in the instructional materials, the following scores were earned:

Group I (pre-organizer)	Group II (post-organizer)	(no organizer
(pre-diguinate)	5	5
4	4	4
5	4	6
3	7	2
6	8	2
6	7	2
3	,	6
3	6	4
4	4	· i
4	4	Ā
2	7	
		thecic that u. =

Perform a one-factor ANOVA; test the null hypothesis that  $\mu_1 = \mu_2 = \mu_3$  at any level of significance you wish.

10. Perform an F-test of the null hypothesis  $H_0: \sum_{i=1}^{k} (\mu_i - \mu_i)^2 = 0$  at the .01 level on the following data which represent weight losses in pounds of subjects under four different diets.

n:	Diet B	Diet C	Diet D
Diet A $ X_{11} = 6 $ $ X_{21} = 8 $ $ X_{31} = 3 $ $ X_{41} = 5 $ $ X_{51} = 6 $	$X_{12} = 11  X_{22} = 13  X_{32} = 15$	$X_{13} = 21$ $X_{23} = 20$ $X_{33} = 17$ $X_{43} = 16$	$X_{14} = 5$ $X_{24} = 9$ $X_{34} = 10$ $X_{44} = 7$ $X_{51} = 7$

- 11. The null hypothesis  $H_0: \mu_1 = \ldots = \mu_0$  is to be tested with an  $\approx$  of .05 and with samples of size n = 5. Suppose that unknown to the experimenter,  $\mu_1 = 10$ ,  $\mu_2 = 10$ ,  $\mu_2 = 12$ ,  $\mu_4 = 14$ , and  $\mu_4 = 14$ ; also, the value of  $\sigma^2$  is 8.00. What is the power of the F-test of  $H_0$  under these circumstances?
- Ten samples of 20 scores each are drawn at random from a single normal
  population with mean κ and variance σ<sup>2</sup>. The sample means of the 10 samples
  have a variance of 2.40, i.e.,

$$s_{x}^{2} = \frac{\sum_{j=1}^{10} (\bar{X}_{j} - \bar{X}_{j})^{2}}{2} = 2.40.$$

Find an estimate of o2, the variance of the original normal population sampled.

- 13. Unknown to the experimenter, the value of σ<sup>2</sup> is 20, and μ₁ = 10, μ₂ = 15, and μ₂ = 20 in the three populations from which he has drawn samples of size n = 10. What is the expected value of MS₂ in this experiment, i.e., what would be the average value of MS₂ over an infinite number of replications of this same experiment?
- 14. Prove that  $SS_w = \sum_{j=1}^{J} \left[ \sum_{k=1}^{n} (X_{ij} \bar{X}_{j})^2 \right]$  is equal to its computational formula

$$SS_{w} = \sum_{j=1}^{J} \sum_{i=1}^{n} X_{ij}^{k} - \sum_{j=1}^{J} \frac{\sum_{i=1}^{n} X_{ij}^{k}}{n}. \quad (Hint: First show that \sum_{i=1}^{n} (X_{ij} - X_{ji})^{k} equals$$

$$\sum_{i=1}^{n} X_{ij}^{k} - \left(\sum_{i=1}^{n} X_{ij}^{k}\right)^{n}, \text{ as was done in Chapter 5.} \quad \text{Then sum the } J \text{ sums of squares}$$
for each erroup across  $L$ )

15. Prove that the value of  $F = MS_b | MS_b$  in a one-way ANOVA with J = 2 equals the square of the value of t for testing the significance of the difference between the two means, i.e., prove that

$$t^2 = \frac{(\bar{X}_{,1} - \bar{X}_{,2})^2}{\{(s_1^2 + s_2^2)/2\}(2/n)}$$

equals

$$F = \frac{n[(X_1 - \overline{X})^2 + (X_2 - \overline{X})^2]}{(s_1^2 + s_2^2)/2}.$$

Start by noting that  $\bar{X}_2 = (X_1 + X_2)/2$ , and then manipulate the numerator of F into the form  $n(\bar{X}_1 - \bar{X}_2)^2/2$ .

#### Reference

Now that you have completed this chapter on the analysis of variance for the one-factor, fixed-effects model, you may wish to read Stanley's (1968a) encyclopedia article to review some points already covered and to introduce certain other concepts that will be discussed in later chapters.

than a simple one-factor ANOVA, both experimenters will end at precisely the same point, i.e., concluding that  $H_0$  is false with a high level of significance (low  $\alpha$ ), although the patterns of the differences between the population means in the two experiments are quite different. This is indeed an unsatisfactory situation.

Most multiple comparison procedures are designed to be used after the null hypothesis of no treatment differences in an ANOVA has been rejected. The purpose of these procedures is the isolation of comparisons between means that are responsible for or contributed to the rejection of  $H_0$ . For example, in the first experiment above, comparing  $\mu_1$  and  $\mu_2$  by forming  $\mu_1 - \mu_1 = 20 - 20 = 0$  reveals that these two means would not have given a high probability of rejecting  $H_0$  if they were the only two treatment groups present. However,  $\mu_4 - \mu_1 = 30 - 20 = 10$  was partly responsible for the significance of the obtained results. The application of multiple comparison procedures to large samples in the first experiment above should have a high probability of leading the experimenter to conclude that  $\mu_1$ ,  $\mu_2$ , and  $\mu_3$  are the same and  $\mu_4$  is different from (and greater than) all three of them.

Your first reaction to this problem is probably the thought that t-tests performed on all possible pairs of means involved in the F-test will reveal where significant differences between means lie. This is an understandable first impression; however, it is quite unacceptable methodology. The t-test was not designed for this use and is invalid when so applied. The t-test is properly applied to 1800 random samples. It is quite a different matter to expect the t-test to be valid for determining the significance of the difference between the smallest and largest sample means in a collection of J means. A t-test applied to the largest and smallest of J means takes no account of how large J is. Isn't it clear that if we take J to be 50 and we draw 50 samples randomly from the same normal population, a t-test will show the largest and smallest to be "significantly different" a large proportion of the time (a far larger proportion than the required α)? In spite of the patent invalidity of t-testing following a significant F-ratio in the analysis of variance, or multiple t-testing in lieu of the analysis of variance, this method has often been and continues to be used.

There are several multiple comparison procedures available, but we shall deal with just two of them in any detail. Such procedures are a relatively recent addition to statistical theory, having been developed during 1957 (see MeHuph and Ellis, 1955; Stanley, 1957a; Sparks, 1963; and Kenyon, 1965).

One of the earliest procedures grew out of Keuls's (1952) extension of some early work by Newman (1939) and is called, quite appropriately, the Newman-Keuls procedure (see Winer, 1962, p. 80 ff.). Another early worker in this field was Duncan (1955), who produced Duncan's New Multiple

SEC. 16.2 THE T-METHOD 383

Range Test. Duncan's procedure has probably been the most popular in the behavioral sciences and education; but until mathematical statisticians resolve their differences or satisfy one another that the derivation of the procedure is valid (see Scheffé, 1959, p. 78, fn. 16), it might be wise for persons in applied fields to observe a moratorium on Duncan's procedure.

If a researcher wishes to compare several treatment groups with a control group to decide which treatments differ from the single control, the procedure due to Dunnett (1955) is available (see Winer, 1962, pp. 90-91). One of the few published reports of educational research in which Dunnett's procedure was used is an article by Scannell and Marshall (1966). You should find this article informative as to the context in which the Dunnett procedure is used. The two most useful multiple comparison procedures are those developed by Tukey and Scheffe: the T-method and the S-method, respectively. Although the T-method and S-method have very general forms, we shall deal with specific cases that encompass almost all of the applications of these methods to problems you are likely to meet.

# 16.2 THE *T*-METHOD

The null hypothesis in a one-way ANOVA has been rejected at the  $\alpha$ -level of significance. J treatments were compared, and the number of observations,  $n_i$  neach group was equal. All assumptions for running the ANOVA were met—at least no evidence existed that any one was violated. Among the J treatments there are J(J-1)/2 pairs (i.e., combinations of J means taken two at a time). Each of these yields a simple comparison of the form  $X_J - X_{J^*}$ , where  $j \neq j^*$ . By observation of these J(J-1)/2 differences between sample means and application of the T-method, the experimenter wishes to decide whether he can regard each  $\mu_I - \mu_I$ , as being different from zero. First we shall see how his decision is reached, then we shall discuss the probability values that apply to his decisions.

- Step 1. All J(J-1)/2 comparisons between sample means of the form  $\bar{X}_1 \bar{X}_2$ , are computed. For example, if three treatments are compared, then  $\bar{X}_1 \bar{X}_2$ ,  $\bar{X}_1 \bar{X}_3$ , and  $\bar{X}_2 \bar{X}_3$  are calculated.
- Step 2. All comparisons  $\overline{X}_J \overline{X}_{J*}$  are divided by  $\sqrt{MS_w} | n_*$  where  $MS_w$  is the mean-square within factor levels from the one-way ANOVA and n is the number of observations in any one group.

Step 3. The  $100(1-\alpha)$  percentile point in the Studentized range distribution with degrees of freedom J and J(n-1) is found from Table F in Appendix A. This percentile point is denoted  $\frac{1}{n} \frac{1}{n} \frac{1}{n} \frac{1}{n} \frac{1}{n-1} \frac{1}{n}$ .

The Studentized range is the difference between the largest and the smallest means of J independent samples each of size n from a normal population, divided by  $\sqrt{MS_n/n}$ . There is a family of distributions of the Studentized range, since a different distribution results for all pairs of values of J and n. The two parameters used to identify a particular Studentized range distribution are J, the number of samples, and J(n-1), the degrees of freedom for  $MS_n$ .

Step 4. All J(J-1)/2 differences  $\overline{X}_{.J} - \overline{X}_{.J}$ , divided by  $\sqrt{MS_w/n}$  are compared with the percentile point. It is concluded that  $\overline{X}_{.J}$  and  $\overline{X}_{.J}$ , are significantly different, i.e., present evidence that  $\mu_J$  and  $\mu_J$ , are different, if  $|\overline{X}_{.J} - \overline{X}_{.J}|$  over  $\sqrt{MS_w/n}$  is greater than  $1_{-x}q_{.J}J_{(J-1)}$ .

An example of the application of the T-method is as follows. An experiment comparing three methods (T=3), each having II observations per group (n=11), yields  $X_1=22.60$ ,  $X_2=23.40$ ,  $X_3=28.50$ , and  $MS_0=4.10$ . An F-ratio significant at the .05 level is obtained.

Step 1. 
$$\overline{X}_{.1} - \overline{X}_{.2} = -0.80$$
.  
 $\overline{X}_{.1} - \overline{X}_{.3} = -5.90$ .  
 $\overline{X}_{.2} - \overline{X}_{.3} = -5.10$ .

Step 2. Dividing the above differences by  $\sqrt{4.10/11} = .610$ , gives -1.311, -9.672, and -8.361.

Step 3. From Table F we see that  $_{,93}q_{3,20} = 3.49$ .

Step 4. Any absolute difference between means divided by √MS<sub>a</sub> in that exceeds 3.49 is significant. Thus, it can be concluded—on the basis of the 7-method—that the population means for groups 1 and 2 do not differ (because 1.311 < 3.49), and the mean of population 3 differs from the means of both population 1 and 2.</p>

# I6.3 CONFIDENCE INTERVALS AROUND CONTRASTS BY THE T-METHOD

Establishing confidence intervals around the differences  $\bar{X}_{.j} - \bar{X}_{.j}$ , should be considered as important, or perhaps more important, than the mere decision of whether the difference is significant. Using the T-method, one can establish a set of simultaneous confidence intervals for the differences between sample means. The confidence interval around  $\bar{X}_{.j} - \bar{X}_{.j}$ , is found from the following formula:

$$(\bar{X}_{,i} - \bar{X}_{,i}) \pm {}_{1-x}q_{J,J(n-1)}\sqrt{MS_w/n}.$$
 (16.1)

In the example used to illustrate the significance testing function of the T-method,  $\bar{X}_{.1}=22.60$ ,  $\bar{X}_{.2}=23.40$ ,  $\bar{X}_{.3}=28.50$ ,  $\sqrt{MS_w}/n=.610$ , and  $.897_{.30}=3.49$ . To establish confidence intervals around the three possible differences between means, one adds and subtracts (3.49)(.610) = 2.13 from each difference. These calculations are performed in Table 16.1.

TABLE 16.1 ESTABLISHING CONFIDENCE INTERVALS AROUND DIFFERENCES
BETWEEN THREE SAMPLE MEANS USING THE I-METHOD

$\overline{X}_{.i} - \overline{X}_{.i}$	$_{1-\alpha}q_{J,J(n-1)}(\sqrt{MS_{n}/n})$	Final calculations
$\vec{X}_{.1} - \vec{X}_{.2} = -0.80$	(3.49)(.610) = 2.13	$-0.8 \pm 2.13 = (-2.93, 1.33)$
$\vec{X}_{.1} - \vec{X}_{.3} = -5.90$	2.13	$-5.9 \pm 2.13 = (-8.03, -3.77)$
$\vec{X}_{.2} - \vec{X}_{.3} = -5.10$	2.13	$-5.1 \pm 2.13 = (-7.23, -2.97)$

Notice that the single confidence interval that includes zero between its bounds corresponds to the nonsignificant difference between the sample means  $X_1$  and  $X_2$ , whereas the other two differences are significant.

Notice that the quantity added to and subtracted from the sample mean difference is the same for any value of j and  $j^*$ . Thus, one determines  $1-\alpha f_{J,I_{n-1}J}\sqrt{MS_n}/n$  and adds it to and subtracts it from each of the J(J-1)/2 mean differences to establish a set of simultaneous confidence intervals by the T-method. Of course, the purpose of any confidence interval of this type is to capture within its limits the value of  $\mu_i - \mu_j$ . The simultaneous confidence intervals are constructed in such a way that the confidence coefficient for any single interval is  $not\ 1-\alpha_s$  however. The meaning of the term simultaneous and the nature of the inference being made with such confidence intervals should become clearer in the following paragraph.

In an experiment comparing five groups, there are 5(4)/2 = 10 comparisons of sample means that can be made. Confidence intervals can be

established around all 10 of these sample mean comparisons by the T-method using Eq. (16.1) and an a of .05, for example. In the set of 10 confidence intervals, some of them will include  $\mu_i - \mu_i$ , within their limits and some will not. The same experiment could be run again and again until thousands of sets of 10 confidence intervals are accumulated. The proportion of these thousands of experiments in which all 10 of the computed confidence intervals around the 10 comparisons  $X_{.j} - X_{.j*}$  include within their limits the value of  $\mu_i - \mu_i$ , is equal to .95, i.e.,  $(1 - \alpha)$ . In other words, in  $100(1 - \alpha)$ percent of the experiments in which J groups of n each are compared, the T-method will yield J(J-1)/2 confidence intervals all of which include  $\mu_i - \mu_{i*}$  within their limits. In some experiments, perhaps only one of the J(J-1)/2 confidence intervals does not capture  $\mu_i - \mu_{i*}$ ; in other experiments, more than one of the set of simultaneous confidence intervals will not capture  $\mu_i - \mu_i$ . In the long run, only 100(z) percent of the experiments will have one or more of the J(J-1)/2 confidence intervals that do not capture  $\mu_i - \mu_{i*}$ .

The idea of a set of simultaneous confidence intervals is a radical departure from the simpler notions of confidence intervals met earlier in this text. Formerly, we spoke of the probability that a confidence interval of the form  $X \pm z \sigma / \sqrt{n}$  included  $\mu$  within its limits. Now we are talking about the probability that for an experiment with J groups every one of a collection of confidence intervals captures the parameters of interest. In probability terms, a collection of intervals was the sample space to which the confidence coefficient,  $1 - \alpha$ , applied. With sets of simultaneous confidence intervals, the sample space is an infinite population of experiments, each experiment yielding J(J-1)/2 sample mean differences.

Suppose that J = 3 groups of n = 10 persons are being compared on some variable, and that  $\mu_1=13$ ,  $\mu_2=10$ , and  $\mu_2=5$ . There are 3(2)/2=3comparisons of means that can be made following the experiment. Imagine that T-method confidence intervals with  $\alpha = .10$  are established around the three comparisons  $\vec{X}_1 - \vec{X}_2$ ,  $\vec{X}_3 - \vec{X}_3$ , and  $\vec{X}_2 - \vec{X}_3$ . Conceivably, the process of drawing three groups of 10 observations and establishing Tmethod intervals around the comparisons could be repeated indefinitely. The graph in Fig. 16.1 shows conceivable results of this process.

The small dots represent sample mean differences. The three dots above I on the horizontal scale in Fig. 16.1 represent the three differences between means obtained the first time the experiment was run. The lines that extend an equal distance above and below each dot represent the intervals established by the T-method. Notice that in the first experiment all three confidence intervals include the true value of  $\mu_j - \mu_j$ . We should expect this to be true with 90% = 100(1-z) of the experiments (J=3, n=10) performed. Notice that in the fifth experiment two of the three intervals do not cover the population mean differences. We would expect that one, two, or all




FIG. 16.1 Sets of simultaneous confidence intervals constructed by the T-method for 10 replications of an experiment comparing three treatments.

three of the intervals would fail to capture the population mean differences in about  $10\% = 100(\alpha)$  of the experiments.

One weakness seems to exist in the adoption of an "experiment-wise" error rate when making multiple comparisons. This weakness relates to the concept of an experiment itself. Generally in research in education and the social sciences, the choice of the number of levels that will constitute a factor in an experiment is arbitrary. Seldom do compelling reasons exist for making this choice. The experimenter would willingly include any number of additional levels in the experiment for comparison if they presented themselves on his doorstep. Some experimenters choose to include a "control themselves on his doorstep. Some experimenters choose to include a "control

group" in any experiment, whereas others do not. The definition of a "factor" (a collection of levels to be compared) in an experiment is quite arbitrary. However, a contrast between method A and method B always means the same thing whether it stands alone or is imbedded in a factor with a dozen levels. (See Wilson, 1962 and Ryan, 1959 and 1962.)

Inspection of the Tukey method shows that the probability of detecting a significant contrast between two means (the power of the contrast) depends upon J, the number of groups compared in the experiment. The width of confidence intervals around individual contrasts depends upon J (as well as n and  $MS_{ob}$ ) in the Tukey method. It seems undesirable to allow either of these related conditions (power and widths of intervals) to be affected by the arbitrary choice of how many levels will constitute an "experiment." After all, int it a contrast that is primary? And shouldn't the probability of cring in concluding that  $\mu_1 - \mu_2$ , is or is not different from zero be unaffected by how many and which other groups one happens to include in the "experiment?"

Although some case can be made for the pre-eminence of a contrast or comparison instead of the experiment as the choice for controlling the rate of errors of the first kind, the statistical techniques most commonly regarded as achieving this purpose-most notably the Newman-Keuls procedure (see Winer, 1962, pp. 80–81)—are not well understood. Suffice it to say that between means separately for all pairs of means with a contrast-wise error rate of  $\alpha$  as is widely believed.

#### 16.4 THE S-METHOD

The Scheffé method of multiple comparisons can be applied whenever the T-method is applicable, as well as in some situations where the T-method is not. The most comprehensive discussion of the S-method appears in Scheffé

The notion of a comparison  $(\mu_1 - \mu_2, say)$  that was met in the previous section is actually a special case of the more general concept of a contrast.

Definition: A contrast among the J population means  $\mu_1, \dots, \mu_J$  is a linear combination

$$\label{eq:power_power} \psi = c_1\mu_1 + c_2\mu_2 + \ldots + c_J\mu_J,$$
 such that  $c_1 + c_2 + \ldots + c_J = 0.$ 

The constants  $c_1, \dots, c_r$  are simply positive and negative real numbers that sum to zero. Thus,  $\psi = \mu_1 + \mu_2 - 2\mu_2$  is a contrast in the three means

 $\mu_1$ ,  $\mu_2$ , and  $\mu_3$ , where  $c_1 = 1$ ,  $c_2 = 1$ , and  $c_3 = -2$ . The difference between any two means is a contrast by the above definition:  $\psi=\mu_3-\mu_6$  corresponds to  $c_3 = 1$ ,  $c_6 = -1$ , and all other c's equal to 0.

Suppose there are five population means  $\mu_1, \ldots, \mu_5$ . Table 16.2 lists

TABLE 16.2

6.2		i	Values of	-	
Contrast	$c_1$	c,	C <sub>3</sub>	C.	
1. $\mu_1 - \mu_1$ 2. $\mu_2 - [(\mu_1 + \mu_4)/2]$ 3. $[(\mu_1 + \mu_2 + \mu_4)/3] - [(\mu_2 + \mu_4)/2]$ 4. $\mu_3 - \mu_1$	1 -1 1 -1	-1 1 -½ 0	0 0 <del>1</del> 1	0 -1 1 0	0 -1 0

contrasts among the five means. Any contrast  $\psi$  can be estimated by replacing  $\mu$ 's with sample means, i.e., an estimate  $\psi$  of  $\psi$  is given by

while means, i.e., an earning 
$$\hat{\psi} = c_1 \vec{X}_{.1} + c_2 \vec{X}_{.2} + \dots + c_J \vec{X}_{.J}.$$
 (16.2)

For example, if  $\psi=\mu_2-\mu_3$ , then an estimate of  $\psi$  is  $\bar{X}_2-\bar{X}_3$ .

Before the significance of any estimated contrast can be judged, it is necessary to estimate the variance of  $\psi$ . If a one-way ANOVA has been performed and  $MS_{\omega}$  found, then an estimate of the variance of  $\psi$  is given by (16.3)

$$\hat{\sigma}_{\psi}^{2} = MS_{w} \left( \frac{c_{1}^{2}}{n_{1}} + \frac{c_{2}^{2}}{n_{2}} + \dots + \frac{c_{J}^{2}}{n_{J}} \right), \tag{16.3}$$

where MSw is the "mean-square within,"

c, is the constant that multiplies the jth mean, and

 $n_i$  is the number of observations in the jth group.

The degrees of freedom (df) associated with this estimate of variance are the df for  $MS_{\omega}$ , N-J, where N, the total number of observations in the entire one-factor ANOVA that precedes the S-method, is equal to  $n_1 +$ 

For example, if J=4, all n's equal 10, and  $\psi=\vec{X}_{.1}-\vec{X}_{.4}$ , an estimate  $n_0 + \ldots + n_J$ of the variance of  $\hat{\psi}$  is

$$\hat{\sigma}_{\psi}^{2} = MS_{w} \left[ \frac{1^{2}}{10} + \frac{0^{2}}{10} + \frac{0^{2}}{10} + \frac{(-1)^{2}}{10} \right] = MS_{w} \left( \frac{2}{10} \right).$$

The degrees of freedom for this estimate of variance are N-J=40-4=

Usually, ψ will be obtained to test the significance of its difference from 36. zero. This test answers questions like the following:

- 1. Is there any difference between overt and covert responding to a teaching program?  $\mu_1 \mu_2$
- 2. Is the average of overt responding with and without confirmation of response different from the same average for covert responding?  $[(\mu_1 + \mu_2)/2] [(\mu_3 + \mu_4)/2]$
- Is the average of the three levels of positive reinforcement in an
  experiment different from the average of the two levels of negative
  reinforcement? [(μ<sub>4</sub> + μ<sub>2</sub> + μ<sub>3</sub>)/3] [(μ<sub>4</sub> + μ<sub>4</sub>)/2]

There are five steps to follow in testing the significance of a contrast among means:

- Step 1. Specify and estimate ψ. The coefficients c<sub>1</sub>,..., c<sub>J</sub> that determine the contrast of interest must be specified. Once this is done, ψ is obtained by substituting sample means for population means, as in Eq. (16.2).
- Step 2. Find an estimate of the variance of ψ. The one-factor ANOVA that precedes application of the S-method contains a MS<sub>w</sub>. This value, the c's, and the sample sizes are substituted into Eq. (16.3) to produce ô<sup>2</sup><sub>c</sub>.
- Step 3. Find  $\hat{\sigma}_{\hat{v}}$ . Take the positive square root of  $\hat{\sigma}_{\hat{v}}^2$ , which was found in step 2.
- Step 4. Form the ratio of  $\psi$  to  $\hat{\sigma}_{\psi}$ . The value of  $\psi$  obtained in step 1 is divided by the value of  $\hat{\sigma}_{\psi}$  found in step 3.
- Step 5. Compare the absolute value of the ratio with the test statistic. The absolute value of the ratio found in step 4 is compared with the square root of (J-1) times the  $100(1-\alpha)$  percentile in the F-distribution with degrees of freedom J-1 and N-J. That is, the absolute value of the ratio is compared with  $\sqrt{(J-1)_{1-4}F_{J-1}}$ . The hypothesis that y=0 is rejected if the absolute value of the ratio exceeds the square root of (J-1) times the percentile point, i.e.,

reject 
$$H_0$$
:  $\psi = 0$  if  $\frac{|\psi|}{\hat{\sigma}_{\psi}} > \sqrt{(J-1)_{1-k}F_{J-1,N-J}}$ .

hangine that four ways of teaching the oxidation-reduction method of balancing chemical equations were compared. The data (scores on a 15-item achievement test) obtained in the experiment appear in Table 16.3.

#### TABLE 16.3

Group	Sample mean	n	
1	$X_1 = 10.32$	$n_1 = 20$	
2	$X_{.2} = 10.54$	$n_2 = 25$	$MS_w = 8.35$
3	$\bar{X}_{s} = 12.86$	$n_2 = 20$	
4	$X_{.4} = 7.17$	$n_4 = 15$	

The F-ratio in the analysis of variance of these data is significant at the .05 level. It is now the purpose of the S-method to uncover the groups contributing to this significant result.

Very likely, all possible pairs of differences between the sample means will be of interest. Thus, the significance of the following differences between sample means must be determined:

$$\begin{split} & \bar{X}_{.1} - \bar{X}_{.2} & \bar{X}_{.2} - \bar{X}_{.3} \\ & \bar{X}_{.1} - \bar{X}_{.3} & \bar{X}_{.2} - \bar{X}_{.4} \\ & \bar{X}_{.1} - \bar{X}_{.4} & \bar{X}_{.2} - \bar{X}_{.4} \end{split}$$

Suppose in addition that a slightly more complicated contrast is of interest to the experimenter. It was the case that groups 1 and 2 were taught by two methods differing only in that group 1 actually observed the pre-

TABLE 16.4

4	Contrast	Estimate of contrast	Estimate of variance of contrast
	A. μ <sub>1</sub> - μ <sub>2</sub>	$\vec{X}_{,1} - \vec{X}_{,2}$	$MS_w\left(\frac{1}{n_1} + \frac{1}{n_2}\right)$
	B. $\mu_1 - \mu_2$	$X_{,1} - X_{,3}$	$MS_{w}\left(\frac{1}{n_1}+\frac{1}{n_2}\right)$
	C. $\mu_1 \sim \mu_4$	$\overline{X}_{-1} - \overline{X}_{-4}$	$MS_w\left(\frac{1}{n_1} + \frac{1}{n_4}\right)$
	$D. \mu_3 \sim \mu_3$	$\overline{X}_{,2} - \overline{X}_{,3}$	$MS_u\left(\frac{1}{n_z} + \frac{1}{n_z}\right)$
	$E. \mu_2 \sim \mu_4$	$\vec{X}_{.5} - \vec{X}_{.4}$	$MS_w\left(\frac{1}{n_k} + \frac{1}{n_k}\right)$
	F. $\mu_3 \sim \mu_4$	$\vec{X}_{.3} - \vec{X}_{.4}$	$MS_{\omega}\left(\frac{1}{n_4}+\frac{1}{n_4}\right)$
	$G. \frac{\mu_1 + \mu_2}{2} - \mu_2$	$\frac{X_{.1} + \bar{X}_{.3}}{2} - \bar{X}_{.3}$	$MS_w\left(\frac{\frac{1}{n_1}+\frac{1}{n_2}+\frac{1}{n_3}\right)$

cipitation of silver chloride while group 2 did not. The data suggest that this experience for group 1 was irrelevant; in fact, groups 1 and 2 may not be different. The experimenter wishes to know whether the average of groups 1 and 2 differs significantly from group 3, which was taught by a substantially different method. This question suggests the contrast  $\{(\mu_1 + \mu_2)(2) - \mu_2\}$ .

The contrasts in which the experimenter is interested appear in Table 16.4. These specifications complete steps 1 and 2 of the five steps to be followed. If we substitute the data into the formulas in Table 16.4 the estimates (see Table 16.5) of the contrasts and the variances of the contrasts result.

TABLE 14.5

Contrast	Ý	σ̂.
A. $\mu_1 - \mu_2$	~0.22	0.752
$B$ , $\mu_1 - \mu_2$	2.54	0.83
$C. \mu_1 - \mu_4$	3.15	0,969
D. μ <sub>2</sub> μ <sub>3</sub>	-2.32	0.75
$E. \mu_3 - \mu_4$	3.37	0.88
$F. \mu_1 - \mu_4$	5.69	0.969
G. $[(\mu_1 + \mu_2)/2] - \mu_2$	-2.43	0,60

As one example of how the calculations in Table 16.5 were made, consider contrast G. The contrast is  $\varphi = [(\mu_1 + \mu_2)/2] - \mu_2$ ; the estimate  $\psi$  of the contrast (step I) is

$$\hat{y} = \frac{\bar{X}_1 + \bar{X}_2}{2} - \bar{X}_3 = \frac{10.32 + 10.54}{2} - 12.86 = -2.43.$$

The estimate of the variance of this contrast (step 2) is

$$\hat{\sigma}_{\psi}^{2} = MS_{w} \left( \frac{1}{4n_{1}} + \frac{1}{4n_{2}} + \frac{1}{n_{3}} \right) = 8.35 (\hat{s}_{0}^{1} + \hat{1}_{00}^{1} + \frac{1}{20}) = 0.61.$$

Step 3. The positive square roots of the estimates of the variance of all six contrasts are now found:

Contrast	- ó.	á;
A	0.752	0.867
В	0.835	0.914
Ç	0.969	0.985
D	0.752	0 867
E	0.885	0.941
F	0.967	0.985
<u> </u>	0,605	0.778

Step 4. The ratio of  $\hat{\psi}$  to  $\hat{\sigma}_{\hat{\varphi}}$  is found for each of the six contrasts.

Contrast	$\psi/\hat{\sigma}_{\psi}^{\star}$
A	-0.22/0.867 = -0.25
В	-2.54/0.914 = -2.76
С	3.15/0.985 = 3.21
D	-2.32/0.867 = -2.67
E	3.37/0.941 = 3.59
F	5.69/0.985 = 5.81
G	-2.43/0.778 = -3.12

Step 5. The absolute values of the ratios found in step 4 are compared with  $\sqrt{(J-1)_{1-2}}F_{J-1}N_{-J}I$ .

$$N = 20 + 25 + 20 + 15 = 80$$
 and  $J = 4$ .  

$$\sqrt{(J-1)_{.85}F_{J-1.N-J}} = \sqrt{3(2.72)} = 2.86.$$

Therefore, if any ratio found in step 4 is greater than 2.86 in absolute value, i.e., if the ratio is above 2.86 or below -2.86, the corresponding contrast is significant. By this criterion, contrasts C, E, F, and G are significantly different from zero by the S-method with  $\alpha = .05$ . Thus the experimenter concludes that

$$\mu_1 - \mu_4$$
,  $\mu_2 - \mu_4$ ,  $\mu_3 - \mu_4$ , and  $\frac{\mu_1 + \mu_2}{2} - \mu_3$ 

are not zero. It cannot be concluded that  $\mu_1$  and  $\mu_2$  differ, or that  $\mu_1$  and  $\mu_2$  differ, or that  $\mu_2$  and  $\mu_3$  differ. For a published application of the S-method to educational psychological research, see Travers et al. (1964).

#### I6.5 CONFIDENCE INTERVALS AROUND CONTRASTS BY THE S-METHOD

The confidence interval around the estimate of a contrast is constructed as follows with the S-method:

$$\hat{\psi} \pm \hat{\sigma}_{\hat{\nu}} \sqrt{(J-1)_{1-\alpha} F_{J-1,N-J}}$$
 (16.4)

Of course, the purpose of this confidence interval is to capture between its extremes the true value of the contrast in the population means,  $\psi$ .

If  $\psi=c_1\mu_1+c_2\mu_2+\ldots+c_J\mu_J$ , then the confidence interval for  $\psi$  around  $\psi$  is

around 
$$\psi$$
 is 
$$(c_1 \overline{X}_{.1} + c_2 \overline{X}_{.2} + \dots + c_J \overline{X}_{.J})$$

$$\pm \sqrt{MS_n \left(\frac{c_1^2}{n_1} + \frac{c_2^2}{n_2} + \dots + \frac{c_J^2}{n_J}\right)} \sqrt{(J-1)_{1-n}F_{J-1,N-J}}. \quad (16.5)$$

Suppose, for example, that  $\psi = \mu_1 - \mu_2$ . The confidence interval around  $\hat{\psi}$  would be

$$(\bar{X}_{.1} - \bar{X}_{.2}) \pm \sqrt{MS_w(\frac{1}{n_1} + \frac{1}{n_2})} \sqrt{(J-1)_{1-a}F_{J-1,N-J}}.$$

If the significance of the difference of the estimated contrasts from zero has been tested first, use of Eq. (16.4) for obtaining confidence intervals is quite convenient. The construction of the S-method confidence intervals will now be illustrated on the sample used in Sec. 16.4.

A sample contrast is significantly different from zero if the confidence interval around it does *not* span zero. Hence, contrasts C, E, F, and G in Table 16.6 correspond to contrasts that differ significantly from zero. In

TABLE 16.6 ILLUSTRATION OF THE CONSTRUCTION OF S-METHOD CONFIDENCE INTERVALS BY EQ. (16.4)

#	σ̂ŷ	$\hat{\psi} \pm \hat{\sigma}_{\psi}^{\wedge} \sqrt{(J-1)_{1-\alpha} F_{J-1,N-J}} = \hat{\psi} \pm \hat{\sigma}_{\psi}^{\wedge}(2.86)$
A0.22 B2.54 C. 3.15 D2.32 E. 3.37 F. 5.69 G2.43	0.867 0.914 0.985 0.867 0.941 0.985 0.778	(-2.71, 2.27) (-5.17, 0.09) (0.35, 5.95) (-4.80, 0.16) (0.68, 6.06) (2.89, 8.49) (-4.66, -0.20)

the case of contrast C, one can state with the confidence afforded by the S-method that the difference between  $\mu_1$  and  $\mu_4$  is some number greater than zero; indeed it probably lies between  $\mu_3$  and 5.95. Similarly, the confidence interval on contrast G tells us that the difference between  $\mu_3$  and the average of the population means  $\mu_1$  and  $\mu_2$  is very probably not zero. On the other hand, the S-method gives us no evidence to conclude with confidence that I-me confidence I-me thand I-me confidence I-me thand I-me confidence I-me than I-me confidence I-me confidence I-me than I-me confidence I-me confidence I-me than I-me confidence I-me confiden

The confidence intervals established by the S-method must be regarded as a set of simultaneous confidence intervals. As was the case with intervals established by the T-method, no probability statement is made about each individual interval. On the contrary, the intervals are constructed in such a

395

way that the entire set of those which can be constructed in any one experiment has a probability of  $1-\alpha$  of capturing the true value of the contrasts estimated within its bounds. In other words, if all possible contrasts among a set of means are formed by the S-method with a confidence coefficient of  $1-\alpha$ , the statement that "all of these S-method confidence intervals contain the true value of the contrasts they estimate" will be true  $100(1-\alpha)$  percent of the times this experiment is run.

# 16.6 THE T- AND S-METHODS COMPARED

The Scheffé method of multiple comparisons is generally regarded by mathematicians as superior to the T-method because of its generality (equal n's are not necessary) and greater sensitivity when complex combinations of the sample means are being estimated. However, you will probably find greater use for the T-method.

An obvious criterion for the choice between use of the T-method or the S-method is the distribution of N among the J groups. A requirement of the T-method, which is not made with the S-method, is that the sample sizes are equal. Hence the T-method cannot be used when not all n's are equal: the S-method must be used with unequal n's. This dictum must be tempered with good sense. We shall see presently that if one is interested in only the differences among pairs of J sample means, the T-method is far superior-in terms of power to detect significant differences and shorter confidence intervals-to the S-method. If the n's are only slightly unequal, e.g.,  $n_1 = 21$ ,  $n_2 = 22$ , and  $n_3 = 20$ , it is wiser to discard at random one person from group 1 and two persons from group 2 and apply the T-method than to apply the S-method. The random exclusion of three persons from the analysis will have a negligible effect on the results, and it allows you to use the more powerful T-method for comparing sample means. On the other hand, this strategy would not work well if, for example,  $n_1 = 5$ .  $n_2 = 30$ , and  $n_3 = 40$  because too much data would have to be discarded to achieve equal n's of 5. In this instance it would be better to use the Smethod even if you were interested only in the three possible differences between sample means.

Suppose that after finding a significant F-ratio in an analysis of variance with equal n's, the experimenter was interested in probing among the J(J-1)/2 differences between sample means. Should he use the T-method the S-method? Either can be applied, but which one is preferred? The T-method is preferred because it produces a greater number of significant differences between means. Equivalently, the T-method will give shorter confidence intervals around differences between means than the S-method.




FIG. 162 Schema for making multiple comparisons among J group means. (Based on Hopkins and Chadbourn, 1967.)

As an example of the relative widths of the confidence intervals around  $X_3 - X_3$ , given by the T-method and the S-method, take the contrast between  $X_4$  and  $X_3$  in Table 16.1. The .95 confidence interval around  $X_3 - X_3$  was found to be (-7.23, -2.97). Now let's establish a .95 confidence interval around the same difference using the S-method.

$$\ddot{S}_{1} - \ddot{X}_{2} = -5.1$$
,  $MS_{e} = 4.1$ ,  $n_{1} = n_{2} = 11$ .  
 $\dot{\sigma}_{1}^{1} = 4.1(\dot{\gamma}_{1}^{1} + \dot{\gamma}_{1}^{1}) = 0.745$ ,  $\sqrt{(J-1)_{3i}F_{J-1,X-J}} = \sqrt{2(3.2)} = 2.58$ .  
The S-method confidence interval around  $\ddot{X}_{1} - \ddot{X}_{2}$  is  $-5.1 \pm (2.58)(.86) = (-7.32, -2.84)$ .

This confidence interval is wider, but not by much, than the confidence interval around the same difference between means yielded by the T-method (-7.23, -2.97). In some instances, the S-method will produce a confidence interval that includes zero—in which case the difference between the means is judged nonsignificant—while the T-method will produce a confidence interval around the same difference that does not include zero, thus leading to judgment of a significant difference between the two means.

When one is interested in contrasts that are more complex than a simple difference between means, e.g.,  $(\mu_1 + \mu_3)/2 = \mu_3$ , then the S-method has more power (and thus yields shorter confidence intervals around an estimate of the contrast) than an extension of the T-method appropriate for such contrasts. We have not discussed the extension of the T-method to contrasts other than  $\mu_1 - \mu_j$ , because it would only be a needless complication in view of the fact that we have developed the S-method, which is preferable with the more complex contrasts. You can find a readable account of the more general T-method in Guenther (1964, pp. 54-57) and a mathematical discussion of the method in Scheffé (1959, pp. 73-77).

The application of the T-method (or the S-method, when n's differ appreciably) to the differences between sample means accounts for almost all of the applications of multiple comparison procedures in educational research and the behavioral sciences. Interest in more complex contrasts—for which the S-method would be more appropriate—is not common. Undoubtedly, however, many opportunities for exploring interesting and worthwhile contrasts with the S-method are overlooked.

A recent textbook on simultaneous statistical inference by Miller (1966) is an excellent reference on these and related topics.

#### PROBLEMS AND EXERCISES

- 1. From Table F in Appendix A, determine the following percentile points:
  - a. 6592.30 b. 9098.120 c. 90910.10 d. 90912.8 e. 9592.80
- 2. Suppose that five samples each of size 7 are to be randomly drawn from a single normal distribution with mean μ and variance σ². The range between the largest, X̄<sub>L</sub>, and smallest, X̄<sub>S</sub>, sample mean will be found and divided by ¬MS<sub>ω</sub>T, where MS<sub>ω</sub> is the average of the five sample variances.
  - a. What is the probability that  $(\bar{X}_L \bar{X}_S)/\sqrt{MS_w/7}$  will be larger than 4.102?
  - b. If this sampling procedure were carried out thousands of times, what percent of the values of  $(X_R X_R)/\sqrt{MS_w/T}$  would exceed 5.048?
- 3. If (X<sub>j</sub> − X<sub>j\*</sub>) ± <sup>38</sup>/<sub>2</sub>Q<sub>j\*</sub>Q<sub>j\*</sub>Q<sub>j\*</sub> does not contain zero, then the difference between X<sub>j\*</sub> and X<sub>j\*</sub> is judged to be significant at the .05 level by the Tukey method. This procedure is equivalent to concluding that X<sub>j</sub> and X<sub>j\*</sub> differ

398

significantly at the .05 level if

$$\frac{|\overline{X}_j - \overline{X}_{j*}|}{\sqrt{MS_n/n}} > .95q_{J,J(n-1)}.$$

Suppose that in a particular experiment, J=6, n=11, and  $MS_w=44,00$ . Which of the following means differ significantly from each other at the z=.05 level by the Tmethol:  $\vec{X}_1=61.25$ ,  $\vec{X}_2=64.72$ ,  $\vec{X}_3=70.57$ ,  $\vec{X}_4=73.42$ ,  $\vec{X}_3=81.66$ , and  $\vec{X}_{.4}=82.17$ ?

- 4. In Prob. 7 at the end of Chapter 15, you performed an ANOVA on data gathered by Guthrie (1967) in an experiment on "discovery learning." Using the 7-method, perform multiple comparisons on all pairs of means in Guthrie's experiment using an a of .05.
- 5. Jeffrey and Samuels (1967) compared phonic (letter) and look-say (word) methods of reading instruction on a measure of transfer of training. Twenty kindergarten pupils learned a list of eight words by the phonic method and 20 pupils learned the same words by the look-say method. A control group of 20 pupils did not learn the original list of words. All 60 pupils were subsequently taught a new list of eight words; the dependent variable in the experiment was the number of trials required to learn the second list. The following data were obtained.

Number of trials to learn second list

	Phonic method	Look-say method	Control
$\overline{X}_{\cdot}$	20 13.50	20 27.20	20 29.25
52	9.94	10.37	9.99

The value of  $MS_w$  was 102.05. Perform multiple comparisons at the .01 level using the Tukey method. Determine which pairs of means differ significantly.

6. Four methods of teaching percentage (case method, formula method, equation method, unitary analysis method) to sixth graders were compared (Sparks, 1963). Thenty-eight sixth-grade classes were randomly assigned to the four methods; seven classes studied under each method. At the conclusion of the teaching unit a 45-item test on computing percentages was administered to each class and the class average recorded. The following data were obtained.

Case	Average test	st score for each class		
method (1)	Formula method (2)	Equation method (3)	Unitary analysis method (4)	
14.59 23.44 25.43 18.15 20.82 14.06 14.26	20.27 26.84 14.71 22.34 19.49 24.92 20.20	27.82 24.92 28.68 23.32 32.85 33.90 23.42	33.16 26.93 30.43 36.43 37.04 29.76 33.88	

- a. Run a one-way analysis of variance on these data. Test the null hypothesis
  of no differences among the four teaching methods at the .05 level of significance.
- b. After completing the F-test, use both the T-method and the S-method to determine which of the six pairs of means can be considered to be significantly different. Which method produces a greater number of significant differences?
- different. Which method produces a greater number of significant differences? c. Set up the 95% confidence interval for  $[(\mu_2 + \mu_3)/2] - [(\mu_1 + \mu_4)/2]$  using the 5-method.
- 7. Klausmeier (1963) studied the effects of accelerating bright older pupuls from the second grade to the fourth grade, thus skipping the third grade. The following five groups of n = 20 pupils each were tested on the Metropolitan Achievement Test during their year in the fifth grade:
  - Ace: a group of 20 bright pupils who were promoted to the fourth grade from
    the second grade after a five-week special summer session. The Ace group is
    now in the fifth grade. Average age of this group is 9 years 6 months.
  - SY: a group of 20 pupils who did not skip the third grade and who are of superior (S) ability and are young (Y), i.e., below median age of fifth graders.
     The SY group is now in the fifth grade and has an average age of 10 years 0 months.
  - SO: 20 nonaccelerated pupils of superior (S) ability and who are older (O)
    than the median fifth grader. The SO group, now in the fifth grade, has an
    average age of 10 years 5 months.
  - 4. AY: 20 nonaccelerated pupils of average ability who are younger than the median fifth grader. The AY group, now in the fifth grade, has an average age of 10 years, 0 months.
  - AO: 20 nonaccelerated fifth graders of average (A) ability who are above the age of the median fifth grader (average age, 10 years, 6 months).

The following table presents data for the Total Language scores on the Metropolitan Achievement Test for the five groups:

	Group				
	Acc	SY	50	AY	AO
n	20	20	20	20	20
$\bar{X}$	55.15	56.40	63.55	47.15	53.45
5-	6.86	4.59	6.86	7.75	12.39

- a. Perform an F-test at the .05 level of  $H_0$ :  $\mu_1 = \ldots = \mu_5$  on the above data.
- b. Follow the F-test in (a) with T-method multiple comparisons to determine which pairs of means differ significantly.
- c. Use the S-method with  $\alpha=.05$  to place a confidence interval on  $\psi=\mu_1-[(\mu_2+\ldots+\mu_3)/4]$ . In effect, then, test the hypothesis that  $\mu_1$ , the mean for accelerated pupils, differs from the average of the remaining four means.

# 17

# THE TWO-FACTOR ANALYSIS OF VARIANCE— FIXED EFFECTS

### I7,I THE LAYOUT AND SYMBOLIZATION OF DATA

The data gathered in a two-factor analysis of variance design are classified with reference to two factors. Such two-factor classifications of things are common in everyone's experience. For example, people can be classified both with respect to sex and race; automobiles can be classified with respect to "year" and "make."

Consider as an example of a two-factor design an experiment in which three methods of teaching beginning reading are compared. The pupils involved in this experiment can be classified in two ways (among others): with respect to the teaching method under which they study (factor A) and factor B, sex, has two levels. Since both boys and girls study under each method in the experiment, six unique combinations of the levels of the two attoins of the levels. Suppose the variable that will be observed in the evaluation of the outcome of this experiment is reading comprehension, symbolized

Factor A, teaching	Factor	Β, sex
method	Male (1)	Female (2)
	X111	X <sub>121</sub>
. [	X <sub>II2</sub>	X <sub>122</sub>
1 {	X113	X <sub>123</sub>
	X <sub>114</sub>	X <sub>124</sub>
	X <sub>211</sub>	X <sub>221</sub>
	X <sub>212</sub>	X <sub>222</sub>
2	X <sub>213</sub>	X223
{	X <sub>214</sub>	X <sub>224</sub>
	X <sub>311</sub>	X <sub>321</sub>
1	X312	X322
3	X <sub>313</sub>	X <sub>323</sub>
	X314	X <sub>324</sub>

FIG. 17.1 Layout of data in a 3  $\times$  2 two-factor ANOVA design with four observations per cell ( $X_{IB}$  notation, where i=1,2,3 for method, j=1,2 for sex, and k=1,2,3,4 for pupil within method-sex group).

by X. Observations on X are taken by administering a standardized test of reading comprehension. If four boys and four girls were taught to read by method 1, four boys and four girls by method 2, etc., the data could be tabulated and symbolized as shown in Fig. 17.1.

A total of 24 pupils participated in this experiment. X<sub>111</sub> represents the reading comprehension test score of the "first"—arbitrarily designated—boy who studied under method 1. X<sub>243</sub> stands for the test score of the "fourth" girl who studied under method 3. The "3" stands for the method, the "2" for the sex (male—1, female—2), and the "4" designates the arbitrarily labeled fourth pupil in the group of four girls under method 3.

In general, an observation in a two-factor ANOVA design is denoted by  $X_{ik}$ , where i is a subscript for factor A and takes on the values  $1, 2, \dots, I_j$  jis the subscript for factor B and takes on the values  $1, 2, \dots, I_j$ ; and k is the subscript that identifies the observations within a cell (combination of levels of factors A and B) of the design and takes on the values  $1, 2, \dots, n$ . If we want to denote the score of the third (3) girl (2) studying under the first (1) method, we let i = 1, j = 2, and k = 3 to obtain  $X_{12}$ . Often with statistical notation one in effect reads subscripts or summation signs from

right to left. To summarize, in  $X_{(0)}$ ,

$$i = 1, 2, ..., I$$
 for factor  $A$ ,  
 $j = 1, 2, ..., J$  for factor  $B$ , and  
 $k = 1, 2, ..., n$  for "within cells,"

This notation is a slight departure from the conventions established in Chapters 15 and 16. In particular, the right-most subscript now indexes observations within a cell, where previously the left index ranged over withincell observations. Master this switch now so that it will not cause confusion later. Note also that now k = 1, 2, ..., n, whereas for the one-factor design  $i = 1, 2, \ldots, n$ .

# 17.2 A MODEL FOR THE DATA

402

Our interest in gathering the data was to determine how the size of the scores varies with the levels of the two factors, i.e., whether boys score higher than girls, whether method 2 gives higher scores than method 1, etc. Toward this end, we shall now devise a fairly abstract model—a generalization of that used with the one-factor ANOVA—to explain, in a general way, how the data

Perhaps the simplest model for data in a two-factor ANOVA would be one that involved only four types of terms or "effects": a term  $\mu$ , the same for each datum, which describes the general size of the scores; I terms  $\alpha_t$ , one for each level of factor A, which describe the excess of the data above or below  $\mu$  for those scores at the ith level of factor A;J terms  $\beta_j$ , one for each level of factor B, which describe the excess above or below  $\mu$  for the scores at the jth level of factor B; and a term  $e_{ijk}$ , for each score that would essentially make up the difference between the score and the sum of the other terms in the model. Under this rudimentary model, a score  $X_{ijk}$  would be

$$X_{ijk} = \mu + \alpha_i + \beta_j + e_{ijk}.$$
(17.1)

For example, if adult males in general tend to be  $5'7' = \mu$ , if feeding them yoghurt tends to add  $1' = x_1$  to their height, if sleeping on a hard mattress tends to add  $1'=\beta_1$  to their height, and if Joe's unique individual history (environmental influences, heredity, errors of measurement, etc.) has taken 3° off the height he would have attained under standard conditions,

$$X_{11(1)ee} = \mu + z_1 + \beta_1 + \epsilon_{11(1)ee} = 5'7'' + 1'' + \frac{1}{2}'' - 3'' = 5'5\frac{1}{2}''$$
.

Unfortunately, the most useful and the state of the st

Unfortunately, the most useful and widely applicable model for the data in a two-factor ANOVA is slightly more complex than the model in Eq.

(17.1). The more useful and descriptive model differs from Eq. (17.1) by a term that denotes the effect of the unique result of combining level i of factor A with level j of factor B. "Unique" in this context means that the outcome one would obtain from combining level i of A with level i of B may not be the simple sum of  $\alpha_i$  and  $\beta_j$ . If not, perhaps a new  $\alpha \beta_{ij}$ , which is not the product of  $\alpha_i$  and  $\beta_i$ , is needed to describe the scores in the ijth cell. Such a term is called an interaction term. [The concept of the interaction of two independent variables is due to the famed English statistician-geneticist. Ronald Fisher. It was Fisher's concepts of experimental control through randomization and the study of the effects of several factors and their interactions simultaneously that successfully overthrew the "one variable at a time" orthodoxy of experimental agriculture in the early 1900's. See Fisher (1925) and Stanley (1966b). If an interaction term is needed in the model to describe the scores, then knowing the general effect of level i of A and of level i of B is not enough to predict the outcome in cell ii. An interaction is tenuously analogous to two polarized lenses: light passes through each lens individually, but when one is laid on top of the other (when they are combined) no light passes through the pair. The separate effects of the lenses differ from their combined effect.

The expanded model, the model upon which the analyses in this chapter are based, is

$$X_{ijk} = \mu + \alpha_i + \beta_j + \alpha \beta_{ij} + e_{ijk}. \qquad (17.2)$$

Without any loss of generality the  $\alpha$ ,  $\beta$ , and  $\alpha\beta$  terms in the above model are assumed to sum to zero over both i and j, i.e.,  $\sum_{i} \alpha_{i} = \sum_{j} \beta_{j} = \sum_{i} \alpha\beta_{ij} = \sum_{i} \alpha\beta_{ij} = 0$ .

In Sec. 17.4 we shall take a closer look at the nature of an ANOVA interaction and examine some illustrations.

#### 17.3 LEAST-SQUARES ESTIMATION OF THE MODEL

With data in hand and having adopted a particular abstract model to explain them, we still have the task of relating the data to the model. What features of the data influence the value of  $\mu$  in Eq. (17.2)? How can the data be manipulated to disclose information about the values of the  $\alpha_i^*$ s, the  $\beta_i^*$ s and the  $\alpha \beta_{ij}$ 's?

This is the same general problem we faced when we sought to predict Y from X by means of a straight line or when we sought to explain the differences between factor levels in a one-factor ANOVA. The problem is basically the same, and our approach to its solution is no different. We fit the model in Eq. (17.2) to the data so that a criterion of least squares is satisfied. In this instance as in the others, the criterion of least squares is

right to left. To summarize, in X12.

$$i = 1, 2, ..., I$$
 for factor  $A$ ,  
 $j = 1, 2, ..., J$  for factor  $B$ , and  
 $k = 1, 2, ..., n$  for "within cells,"

This notation is a slight departure from the conventions established in Chapters 15 and 16. In particular, the right-most subscript now indexes observations within a cell, where previously the left index ranged over withincell observations. Master this switch now so that it will not cause confusion later. Note also that now k = 1, 2, ..., n, whereas for the one-factor design  $i = 1, 2, \ldots, n$ .

#### 17.2

# A MODEL FOR THE DATA

Our interest in gathering the data was to determine how the size of the scores varies with the levels of the two factors, i.e., whether boys score higher than girls, whether method 2 gives higher scores than method 1, etc. Toward this end, we shall now devise a fairly abstract model—a generalization of that used with the one-factor ANOVA-to explain, in a general way, how the data

Perhaps the simplest model for data in a two-factor ANOVA would be one that involved only four types of terms or "effects": a term  $\mu$ , the same for each datum, which describes the general size of the scores; I terms  $\alpha_i$ , one for each level of factor A, which describe the excess of the data above or below  $\mu$  for those scores at the  $\hbar h$  level of factor A; J terms  $\beta_I$ , one for each level of factor B, which describe the excess above or below  $\mu$  for the scores at the jth level of factor B; and a term  $e_{ijk}$ , for each score that would essentially make up the difference between the score and the sum of the other terms in the model. Under this rudimentary model, a score  $X_{ijk}$  would be

$$X_{ijk} = \mu + \alpha_i + \beta_j + e_{ijk}. \tag{17.1}$$

For example, if adult males in general tend to be 5'7' =  $\mu$ , if feeding them yoghurt tends to add  $1' = \alpha_1$  to their height, if sleeping on a hard mattress tends to add  $\frac{1}{4}'=\beta_1$  to their height, and if Joc's unique individual history (environmental influences, heredity, errors of measurement, etc.) has taken 3° off the height he would have attained under standard conditions,

$$X_{\text{IIIJoe}} = \mu + \alpha_1 + \beta_1 + e_{\text{IIIIJoe}} = 5'7'' + 1'' + \frac{1}{2}'' - 3'' = 5'5\frac{1}{2}''$$
.

Unfortunately, the most useful and with the formula of the state of the sta

Unfortunately, the most useful and widely applicable model for the data in a two-factor ANOVA is slightly more complex than the model in Eq.

(17.1). The more useful and descriptive model differs from Eq. (17.1) by a term that denotes the effect of the unique result of combining level i of factor A with level j of factor B. "Unique" in this context means that the outcome one would obtain from combining level i of A with level i of B may not be the simple sum of  $\alpha_i$  and  $\beta_i$ . If not, perhaps a new  $\alpha\beta_{ij}$ , which is not the product of  $\alpha$ , and  $\beta_i$ , is needed to describe the scores in the iith cell. Such a term is called an interaction term. [The concept of the interaction of two independent variables is due to the famed English statistician-geneticist, Ronald Fisher. It was Fisher's concepts of experimental control through randomization and the study of the effects of several factors and their interactions simultaneously that successfully overthrew the "one variable at a time" orthodoxy of experimental agriculture in the early 1900's. See Fisher (1925) and Stanley (1966b). I If an interaction term is needed in the model to describe the scores, then knowing the general effect of level i of A and of level j of B is not enough to predict the outcome in cell ij. An interaction is tenuously analogous to two polarized lenses: light passes through each lens individually, but when one is laid on top of the other (when they are combined) no light passes through the pair. The separate effects of the lenses differ from their combined effect.

The expanded model, the model upon which the analyses in this chapter are based, is

$$X_{ijk} = \mu + \alpha_i + \beta_i + \alpha \beta_{ij} + e_{ijk}. \tag{17.2}$$


Without any loss of generality the  $\alpha$ ,  $\beta$ , and  $\alpha\beta$  terms in the above model are assumed to sum to zero over both i and j, i.e.,  $\sum_{i} \alpha_{i} = \sum_{j} \beta_{ij} = \sum_{i} \alpha\beta_{ij} = \sum_{i} \alpha\beta_{ij} = 0$ .

In Sec. 17.4 we shall take a closer look at the nature of an ANOVA interaction and examine some illustrations.

#### I7.3 LEAST-SQUARES ESTIMATION OF THE MODEL

With data in hand and having adopted a particular abstract model to explain them, we still have the task of relating the data to the model. What features of the data influence the value of  $\mu$  in Eq. (17.2)? How can the data be manipulated to disclose information about the values of the  $\alpha_i$ 's, the  $\beta_i$ 's and the  $\alpha\beta_i$ 's?

This is the same general problem we faced when we sought to predict Y from X by means of a straight line or when we sought to explain the differences between factor levels in a one-factor ANOVA. The problem is basically the same, and our approach to its solution is no different. We fit the model in Eq. (17.2) to the data so that a criterion of least squares is satisfied. In this instance as in the others, the criterion of least squares is a



follows: (1) values are substituted into Eq. (17.2) for  $\mu$ ,  $\alpha_1, \ldots, \alpha_{I}$ ,  $\beta_1, \ldots, \beta_J$ ,  $\alpha_{I_{11}}, \ldots, \alpha_{I_{J_{12}}}$  (2) these values along with the data  $X_{G2}$  determine, by subtraction, values for the Un errors  $e_{G3}$ ; (3) when the sum of the squared errors so determined is as small as it is possible to make it, the least-squares estimates of  $\mu$ , the  $\alpha_i$ 's, the  $\beta_i$ 's and the  $\alpha_i\beta_i$ 's have been found.

For example, eight scores are gathered in a simple  $2 \times 2$  design, with n also 2, as shown in Fig. 17.2. We postulate the following model for the data:

$$X_{ijk} = \mu + \alpha_i + \beta_j + \alpha \beta_{ij} + \epsilon_{ijk}, \qquad i = 1, 2$$

$$j = 1, 2$$

The hypothetical main and interaction effects underlie the data in Fig. 17.2 in the manner depicted in Fig. 17.3. Hence it is assumed, for example, that

$$X_{111} = \mu + \alpha_1 + \beta_1 + \alpha \beta_{11} + e_{112}.$$

By subtraction of appropriate terms, we see that

$$e_{112} = X_{112} - (\mu + \alpha_1 + \beta_1 + \alpha \beta_{11}).$$
 (17.3)

k = 1.2

Suppose we let  $\mu=5$ ,  $\alpha_1=2$ ,  $\alpha_2=1$ ,  $\beta_1=0$ ,  $\beta_2=1$ , and all the  $\alpha\beta_{ij}$ 's equal zero. Will this give us the smallest value for the sum of the eight squared errors? We can calculate  $\epsilon_{112}$  by Eq. (17.3):

$$e_{112} = 4 - (5 + 2 + 0 + 0) = -3.$$

Calculating the eight remaining  $e_{in}$ 's similarly, squaring each, and summing the squares produce a value of 132. Is this the smallest sum of squared errors possible? No, it so happens that the smallest sum of squared errors is produced by the following least-squares estimates of the parameters

of the model:

$$\begin{array}{ll} \mu = 4 & \alpha \beta_{11} = -.5 \\ \alpha_1 = -1.5 & \alpha \beta_{12} = .5 \\ \alpha_2 = 1.5 & \alpha \beta_{21} = .5 \\ \beta_1 = 1 & \alpha \beta_{22} = -.5 \\ \beta_2 = -1 & \end{array}$$

The sum of the squared estimated errors, which were obtained by substituting the above least-squares estimates into Eq. (17.2) along with the data  $X_{tit}$ , is equal to 8.00, the smallest possible value for any choices of  $\mu$ , the  $\alpha_i$ , the  $\beta_i$ , and the  $\alpha\beta_{ii}$ .

What manipulations of the data will produce the least-squares estimates? As was true in the one-factor ANOVA the least-squares estimates of the terms in the model for the data are obtained by simple averaging of the data in various ways. For example, the least-squares estimate of  $\mu$  is just the mean of all eight scores in the 2  $\times$  2 table. The least-squares estimate of  $\alpha_1$ is just the mean of the four scores in row 1 of the table minus the mean of all eight scores. The least-squares estimate of  $\beta_2$  is the mean of the four scores in column 2 minus the mean of all eight scores. The least-squares estimate of  $\alpha\beta_{11}$  is the mean of the two scores in the cell at the intersection of row 1 and column 1 minus the mean of row 1 minus the mean of column 1 plus the mean of all eight scores.

Let  $\mu_{ij}$  be the mean of the population of scores from which those scores in the ith row and the jth column of the data layout were sampled. Let  $\bar{\mu}_{i}$ , be the average of the  $J \mu_{ij}$ 's in the *i*th row; let  $\bar{\mu}_{ij}$  be the average of the  $I \mu_{ij}$ 's in the jth column; and let  $\mu$  be the average of all the  $\mu_{ij}$ 's. For example, see Fig. 17.4.

The least-squares estimates of the terms in the model for the  $2 \times 2$ ANOVA design can be characterized in terms of their definition in the sample and their long-range average (or expected) value that they attain in the population. This has been done in Table 17.1.




FIG. 17.4

TABLE 17.1 LEAST-SQUARES ESTIMATION IN THE TWO-FACTOR FIXED-EFFECTS ANOVA

	ANOVA Term in the model	Least-squares estimate	Population value
_	μ	$\hat{\mu} \approx \bar{x}_{}$ $\hat{\alpha}_1 = \bar{x}_{} - \bar{x}_{}$	$\mu \atop \mu_{1} \sim \mu$
	α <sub>1</sub> α <sub>1</sub>	$\hat{a}_1 = \hat{x}_1 - \hat{x}_1$	$\hat{\mu}_{t}$ . — $\mu$
		$\hat{\beta}_1 = \bar{x}_{.1}, \cdots \bar{x}_{}$	$\mu_{.1} - \mu_{.1}$
	$\beta_1$ $\beta_2$	$\hat{\beta}_2 = \bar{x}_{,2} - \bar{x}_{,}$	$\tilde{\mu}_{,1} \sim \mu$
	$x_{\beta_{22}}$	$\alpha \tilde{\beta}_{11} = \tilde{x}_{11} - \hat{x}_{1} - \tilde{x}_{.1.} + \hat{x}_{}$	$\mu_{11} \sim \hat{\mu}_{11} - \hat{\mu}_{A} + \mu$
	2311	$\widehat{x}\widehat{\beta}_{12} = \widehat{x}_{12} - \widehat{x}_{1.} - \widehat{x}_{3.} + \widehat{x}_{}$	$\mu_{11} \sim \hat{\mu}_{1} = \hat{\mu}_{-1} + \mu$
	$a\beta_{st}$	$\widehat{\alpha\beta}_{22} = \bar{x}_{21}, -\bar{x}_{2.}, -\bar{x}_{i,1} + \bar{x}_{i,1}$	$\mu_{21} \sim \tilde{\mu}_{1.} - \tilde{\mu}_{.1} + \mu$
	$\alpha \beta_{11}$	$\alpha \beta_{23} = \hat{x}_{23} - \hat{x}_{2.} - \hat{x}_{3.} + \hat{x}_{}$	$\mu_{24} \sim \hat{\mu}_{2*} - \hat{\mu}_{*2} + \mu$

Before proceeding to the problem of testing hypotheses about the terms of the two-factor ANOVA model, it would be wise to dwell a little longer on the meaning of what has been called the interaction of factors A and B.

#### 17.4 THE NATURE OF INTERACTION

In addition to being interested solely in the effect one variable (independent) has on another variable (dependent), investigators frequently ask whether this effect is the same for all levels of a second, independent variable. If this effect is not the same, an interaction between the two independent variables is said to exist. Suppose three different methods of teaching are being compared experimentally. It is found that one method is relatively best with high ability students whereas another method is relatively best with low ability students. We say that an interaction between teaching method and student ability exists.

In a study on test-wiseness, multiple-choice and free-response items were given to both American and Indonesian students. Interest was not focused on which type of item was harder or which group of students had the superior scores, but on the interaction between nationality and item type. Did one maitonality do telatively better on one type of item and the other nationality do telatively better on the other type (interaction); or was the degree of superiority of one nationality over the other the same for both types of item (no interaction)?

An expanded the same for both

An example of hypothetical data displaying an interaction effect between sex and type of reading material on reading speed is shown in Fig. 12.5. Notice that the boys read relatively faster with material of one type; girls, with material of another type. The difference between boys' and girls' reading speed is different for different reading matter. Notice that when




FIG. 17.5 Graphic representation of the mean reading speed scores for boys and girls on three types of reading material.

boys and girls are combined, i.e., the lines are averaged, the three word groups are equally difficult. We say that there is no main effect for "type of reading material." Further, when the three reading materials are combined, the boys and girls read equally fast.

Consider the data in Fig. 17.6. Here again the boys read relatively fastest with material on "sports." Even though the girls are clearly faster than the boys, their superiority is not the same for all reading materials. Thus, there is an interaction between sex and reading materials (as well as a difference between the sexes). Notice that the requirement for interaction is merely that the differences between the sexes on the different reading




FIG. 17.6 Graphic representation of the mean reading speed scores for boys and girls on three types of reading material.

materials be different—they do not have to be reversed, as was the case in Fig. 17.5. This is why the words "relatively the same" and "relatively fastest" were used in the introductory paragraphs.

Differences across reading material in the difference between the sexes are reflected geometrically by nonparallel slopes. The fact that the solid and dotted lines are nonparallel reflects the presence of interaction.

A. Demonstration that Absence of Interaction Between Two Factors Implies Parallel Lines in the Graph of Cell Means

Suppose we have a  $2 \times 2$  analysis of variance design as shown in Fig. 17.7. In this diagram,  $\mu$ 's represent population means for rows, columns, cells, and total. If no interaction exists between the two independent variables, then the cell mean equals the sum of the grand mean, the row main effect, and the column main effect, e.g.,

$$\mu_{11} = \mu + (\bar{\mu}_{1.} - \mu) + (\bar{\mu}_{.1} - \mu).$$

If interaction does exist, then this equality does not hold.





Suppose we graph the four cell means in Fig. 17.7 in such a way that  $(\mu_1, \mu_1) = (\mu_1, \mu_2) = (\mu_1, \mu_2)$ , which results in the parallel lines shown in Fig. 17.8 for the lines in Fig. 17.8 to be parallel, the vertical distance from one line to the other must be equal at all points. In particular, the distance from  $\mu_1$  to  $\mu_1$ , must equal the distance from  $\mu_1$  to  $\mu_2$ , \*

<sup>&</sup>quot;The lines are not essential, of course, but just a convenient device to aid the reader A can be takeled with \$\tilde{\ell}\_{i,k}\$ and \$\tilde{\ell}\_{i,k}\$. At and \$A\_k\$, arbitrarily ordered and spaced, would ANOVA for factor \$A\$ in mind. See Stanley (1992)




FIG. 17.9 Graphic representation of the mean reading speed scores for boys and girls on three types of reading material.

Assuming no interaction in the analysis of variance sense, we have

and 
$$\mu_{11} = \mu + (\bar{\mu}_1 - \mu) + (\bar{\mu}_{.1} - \mu)$$
 
$$\mu_{12} = \mu + (\bar{\mu}_1 - \mu) + (\bar{\mu}_{.3} - \mu).$$
 Also, 
$$\mu_{21} = \mu + (\bar{\mu}_2 - \mu) + (\bar{\mu}_{.1} - \mu)$$
 and 
$$\mu_{22} = \mu + (\bar{\mu}_2 - \mu) + (\bar{\mu}_2 - \mu).$$

The distance from  $\mu_{21}$  to  $\mu_{12}$  is  $\mu_{11} - \mu_{12}$ , which you can easily show to be equal to  $\bar{\mu}_1 - \bar{\mu}_{22}$ . The distance from  $\mu_{21}$  to  $\mu_{22}$  is  $\mu_{21} - \mu_{22}$ , which also equals  $\bar{\mu}_1 - \bar{\mu}_{22}$ . The two distances between the pairs of means are equal. Thus, the lines are parallel.

Notice that the slopes of the lines for the boys (and, in turn, for the girls) is the same in Fig. 17.5 and in Fig. 17.6. Consequently, the degree or magnitude of the nonparallelism, and hence the degree of interaction, is identical in Fig. 17.5 and Fig. 17.6.

Figure 17.9 illustrates a situation in which no interaction is present. Notice that the lines for the boys and girls are parallel and thus the difference in reading speed between the sexes is the same for each type of reading material.

#### B. A Caution

One should not overlook artificial reasons for the existence of interaction. For example, consider Fig. 17.10, which illustrates the interaction




FIG. 17.10 Illustration of a possibly artificial interaction between intelligence and learning condition.

between intelligence and learning condition. In a sense, the definition of an interaction effect in terms of a simple sum of population means and the consequence that "no interaction" implies parallel lines when the means are graphed is an imperfect interpretation of our notion of a "unique" and "unpredictable" outcome resulting from combining the levels of two factors. Certainly, many graphs of means from a two-factor design evidence nonparallel lines, yet we feel uncomfortable talking about "unique and unpredictable" effects of the combining of two levels when the nonparallel lines result merely from a "ceiling" or "cellar" effect on a particular paper-andpencil test. (For an example of such interaction, see Stanley, 1969.) This illustrates the fact that perhaps allowing our least-squares estimation procedure to determine that interaction effects would be measured by  $\mu_B$  - $\tilde{\mu}_{i} - \tilde{\mu}_{i} + \mu$  did not capture the full richness of our intuitive notion of  $\alpha\beta_{tt}$ , the unique and unpredictable (from main effects, at least) result of combining factor levels. Such failure to see our intuitive notions reflected perfectly in our mathematical models is a hazard (or reality) of an attempt to represent the real world mathematically.

## C. Two Types of Interaction

In the statistical literature, a useful distinction is made (e.g., by Lubin, 1962a) between two types of interaction: ordinal and distordinal. In the ordinal case, the rank order of the categories of one variable on the basis of their dependent variable scores is the same within each category of the second independent variable. In Fig. 17.6 we see an example of ordinal interaction. It is ordinal because the girls read faster than the boys on each type of reading material. Figure 17.10 also illustrates ordinal interaction. However, in Fig. 17.5 it is not the case that girls read faster than boys on each type of

reading material. Figure 17.5 presents an example of disordinal interaction. When the lines do not cross, interaction is said to be "ordinal"; when the lines cross, interaction is said to be "disordinal."

The importance of this distinction for interpretation is this. When interaction is ordinal, it makes sense to assume, for example, that when girls score higher than boys on the dependent variable the superiority exists for all types of reading material. That is, even though an interaction exists, it is still meaningful to make a single statement about boys and girls, over-all, without qualification or reference to some other variable. However, when there is a disordinal interaction, asserting that girls score higher (i.e., read faster) than boys cannot be understood to mean the superiority is maintained on all types of reading material. Disordinal interaction indicates that only in some situations are girls superior to boys on the dependent variable.

Let's consider an example in which teaching methods 1 and 2 are being tried out in either urban or rural settings. If there is a disordinal interaction between "teaching method" and "setting" and a significant difference between methods in favor of method 1, it would not be possible to claim that method 1 is better than 2 "across the board." Before we could recommend that a school adopt one method or the other, we would first want to know whether the school was rural or urban.

The advice usually given is that whenever there is a significant interaction, one should plot the means for the various combinations. Unfortunately, research reports usually do not present sufficient data to allow the reader to follow this advice.

It should be emphasized that "ordinality" and "disordinality" are properties of graphs. A choice exists between placing factor A or factor B on the abscissa when graphing an interaction. The same cell means can give an ordinal interaction when factor A is on the abscissa and a disordinal interaction when factor B is on the abscissa.

A technical article well worth reading when one has acquired an introduction to analysis of variance is Lubin (1962a).

#### 17.5 STATEMENT OF NULL HYPOTHESES

If in Fig. 17.1 factor B, "sex," was disregarded, the data would be identical to those gathered in a one-factor experiment comparing three teaching methods. Eight observations would have been gathered under each level of factor A, and the one-factor ANOVA model would be appropriate for a statistical inferential test of the hypothesis that the three population means underlying the teaching methods were equal. This null hypothesis, namely that the population means for the three teaching methods are equal,

is identical to one null hypothesis of interest in the two-factor ANOVA. Specifically, we are interested in whether the data gathered in a two-factor ANOVA support or run counter to a decision to accept as true the statement  $H_1$  that  $\tilde{\mu}_1 = \tilde{\mu}_2 = \tilde{\mu}_2$ .

In any two-factor ANOVA, the null hypothesis for factor A can be stated as follows:

$$H_0: \bar{\mu}_1 = \bar{\mu}_2 = \ldots = \bar{\mu}_L$$

(Null hypothesis for factor A)

Notice that the equality of the I population means underlying the I levels of factor A has implications that can be used to state H<sub>0</sub> in several equivalent forms.

Because, when  $k=1,2,\ldots,n$  for every lith factor-level combination,  $\mu$  is the average of the IJ population means (one for each cell), equality of the  $\hat{\mu}_{c}$ 's implies that  $each \, \hat{\mu}_{i_{c}} = quals \, \mu$ . If each  $\hat{\mu}_{i_{c}} = \mu$ , then  $\hat{\mu}_{i_{c}} - \mu = 0$  for the I levels of factor A. Since  $a_{i_{c}}$  the main effect for level i of factor A, is equal to  $\hat{\mu}_{i_{c}} - \mu$ , all  $a_{i_{c}}$ 's equal zero if the null hypothesis is true. The following statements are equivalent ways of stating the null hypothesis for factor A:

1. 
$$H_0$$
:  $\bar{\mu}_{1.} = \ldots = \bar{\mu}_{L}$ ,  
2.  $H_0$ :  $\sum_{i=1}^{I} (\bar{\mu}_{i.} - \mu)^2 = 0$ ,  
3.  $H_0$ :  $\sum_{i=1}^{I} \alpha_i^2 = 0$ ,

4. 
$$H_0$$
: all  $(\bar{\mu}_i - \mu) = 0$ , and

5.  $H_0$ : all  $x_i = 0$ ,

In the above discussion, it was immaterial that we chose to talk about for A instead of factor B. The development of the statement of the null hypothesis for factor B is perfectly analogous to the development of  $H_0$  for A. Specifically, as regards factor B, we are interested in rejecting or accepting the null hypothesis that the J population means underlying the levels of factor B are all equal. The following statements are equivalent ways of stating the null hypothesis for factor B:

1. 
$$H_{\bullet}$$
:  $\tilde{\mu}_{.1} = \tilde{\mu}_{.2} = \ldots = \tilde{\mu}_{.J}$ ,

2. 
$$H_{\mathbf{0}}: \sum_{i=1}^{J} (\hat{\mu}_{.i} - \mu)^{\mathbf{1}} = 0$$
,

3. 
$$H_0$$
:  $\sum_{i=1}^{J} \beta_i^2 \approx 0$ ,

4. 
$$H_0$$
: all  $(\bar{\mu}_1 - \mu) = 0$ , and

5.  $H_{\bullet}$ : all  $\beta_i = 0$ 

There are many ways in which the null hypothesis about a main effect, i.e., about a single factor, can be false. For factor A,  $\hat{\mu}_1$ , could equal 20,5 and the remaining  $\hat{\mu}_i$  could equal 21,1, for example. Or  $\hat{\mu}_1 = \hat{\mu}_2 = 16.65$  and  $\hat{\mu}_3 = \hat{\mu}_4 = 17.80$ . In both instances,  $H_0$  is false. All that it takes for  $H_0$  to be false is for at least two population means to be unequal. The decision faced in an ANOVA is whether one should opt for the truth of  $H_0$  or the truth of  $H_1$ , the logical converse of  $H_0$ , which is true when  $H_0$  is false.  $H_1$ , the alternative hypothesis, can be stated in the following equivalent ways for factor A:

- I.  $H_1: \tilde{\mu}_i \neq \tilde{\mu}_i^*$ , where i and i\* are distinct,
- 2.  $H_1: \sum_{i=1}^{I} (\bar{\mu}_{i.} \mu)^2 \neq 0$ ,
- 3.  $H_1: \sum_{i=1}^{I} \alpha_i^2 \neq 0$ ,
- 4.  $H_1$ :  $\vec{\mu}_i$ ,  $-\mu \neq 0$  for at least one i, and
- 5.  $H_1$ :  $\alpha_i \neq 0$  for at least one i.

Each of the above equivalent statements will be true if and only if the null hypothesis for factor A is false. Hence, if we reject  $H_0$ , we automatically accept  $H_2$ .

The form of the alternative hypothesis for factor B is perfectly analogous to the form of  $H_1$  for factor A. The reader may wish to state  $H_1$  for factor B in at least five equivalent ways.

There remains one hypothesis of interest, and it concerns the collection of interaction terms,  $a\beta_{ij}$ . In Sec. 17.4 our attention was directed toward two sets of conditions: (1) the graph of the population means produced parallel lines; (2) the graph of the population means produced nonparallel lines. We saw in Sec. 17.4 that if no interaction exists between A and B, i.e., if the graph of the population means shows parallel lines, then  $\mu_{ij}$  will equal  $\mu + (\bar{\mu}_i - \mu) + (\bar{\mu}_j - \mu) = \mu + \alpha_i + \beta_j$ . An equivalent condition is that  $\mu_{ij} = \bar{\mu}_i + \bar{\mu}_j - \mu$ . Now if this condition is satisfied, then

$$\mu_{ij} - \tilde{\mu}_{ij} - \tilde{\mu}_{ij} + \mu = 0$$
 for all the  $\mu_{ij}$ 's.

Or, in terms of the model in Eq. (17.2),

$$\alpha \beta_{ij} = 0$$
 for all  $i$  and  $j$ .

If the lines in the graph of the cell means are not parallel at any single point on A, then at least one  $\alpha \beta_{ij}$  is not equal to zero. Hence, parallel lines in the graph of the II population means correspond to all of the  $\alpha \beta_{ij}$ 's not equaling zero; nonparallel lines correspond to at least one of the  $\alpha \beta_{ij}$ 's not equaling zero. These two conditions represent the null hypothesis and the alternative hypothesis, respectively, about the interaction of factors A and B. There are several equivalent ways of stating the null hypothesis  $H_0$  and

Equivalent statements of H1 for the

Equivalent statements of H. for the

the alternative hypothesis  $H_1$  about the interaction effects. Some of these follow:

interaction of A and B	interaction of A and B		
1. $H_0$ : all $(\mu_{ii} - \hat{\mu}_{i_*} - \hat{\mu}_{.i} + \mu) = 0$ ,	1. $H_1$ : $\mu_{ij} - \tilde{\mu}_{i_*} - \tilde{\mu}_{,j} + \mu \neq 0$ for at least one $\mu_{ij}$ ,		
2. $H_0$ : all $\alpha \beta_{ii} = 0$ ,	2. $H_1$ : $\alpha \beta_{ij} \neq 0$ for at least one $\alpha \beta_{ij}$ ,		
3. $H_1: \sum_{i=1}^{I} \sum_{i=1}^{J} (\mu_{ii} - \tilde{\mu}_{i_i} - \tilde{\mu}_{J} + \mu)^2 = 0$	3. $H_1$ : $\sum_{i}^{I} \sum_{j}^{J} (\mu_{ij} - \hat{\mu}_{i,} - \hat{\mu}_{,j} + \mu)^{1} \neq 0,$		

In summary, there are three pairs of hypotheses that are typically of interest in the two-factor ANOVA-(1)  $H_0$  and  $H_1$  for factor A, (2)  $H_0$  and  $H_1$  for the interaction of A and B. In the remaining sections of this chapter we shall see how the data gathered in a two-factor experiment are brought to bear on the decision to accept either  $H_0$  or  $H_1$  for  $A_1$  for  $B_2$  and for the interaction of A and B.

#### 17.6 SUMS OF SQUARES IN THE TWO-FACTOR ANOVA

As in the one-factor ANOVA, the route to statistical inferential tests of the three null hypotheses in the two-factor ANOVA is marked by sums of squares, degrees of freedom, mean squares, expected mean squares, and Fratios. As you travel this route, you may begin to feel that the markers do not point toward your destination in any obvious way. Patience is required because the meaning of the preliminary steps will not become clear until you are on the last leg of the journey.

There are four sources of variation, as they are called, in the two-factor NOVA: (1) factor A, (2) factor B, (3) the interaction of A and B, and (4) "within" cells or combinations of levels of A and B. We shall define sum of squares for each source of variation in turn.

# Sum of Squares for Factor A

The sum of squares for factor A, denoted  $SS_A$ , is simply nJ times the sum of the squared least-squares estimates of the  $x_i$ 's.

$$SS_A = nJ \sum_{i=1}^{J} \hat{x}_i^2 = nJ \sum_{i=1}^{J} (X_{i,.} - X_{..})^2.$$
 (17.4)

415

Recall that  $\alpha_t = \bar{\mu}_t$ ,  $-\mu$  is estimated by  $\bar{X}_{t...}$ , the mean of the nJ scores in level i of factor A, minus  $\bar{X}_{...}$ , the mean of all nIJ scores in the layout of data. The sum over all I levels of A of these squared estimates,  $(\bar{X}_{t...} - \bar{X}_{...})^2$ , is called the sum of squares for factor A.

Equation (17.4) for  $SS_A$  is definitional and does not make for easy computing. We shall return to the matter of actually calculating  $SS_A$  in Sec. 17.9.

#### Sum of Squares for Factor B

The sum of squares for factor B is nI times the sum of the squared least-squares estimates of the  $B_i$ 's:

$$SS_B = nI \sum_{i=1}^{J} \hat{\beta}_j^2 = nI \sum_{i=1}^{J} (\vec{X}_{.j.} - \vec{X}_{...})^2.$$
 (17.5)

Notice that nI is the number of scores averaged to obtain  $X_{J,\cdot}$ . In  $SS_A$ , nI was the number of scores averaged to obtain  $X_{I,\cdot}$ . Again, the above formula for  $SS_B$  is not convenient for computation. Computational formulas will be presented in Sec. 17.9.

# Sum of Squares for the Interaction of A and B

$$SS_{AB} = n \sum_{i=1}^{I} \sum_{j=1}^{J} \widehat{\alpha} \widehat{\beta}_{ij}^{2} = n \sum_{i=1}^{I} \sum_{j=1}^{J} (\vec{X}_{ij} - \vec{X}_{in} - \vec{X}_{j} + \vec{X}_{n})^{2}. \quad (17.6)$$

Notice as before that n, the factor multiplying the sum, is the number of scores averaged to obtain  $X_{ij}$ .

# Sum of Squares "Within" Cells

There remains one sum of squares, SSw, the sum of squares within cells.

$$SS_{w} = \sum_{i=1}^{J} \sum_{j=1}^{J} \sum_{k=1}^{n} (X_{ijk} - \bar{X}_{ij})^{2}.$$
 (17.7)

The meaning of these four sums of squares will begin to emerge when the corresponding mean squares and their expected values are considered in the following sections.

Incidentally, though it is of no great significance in and of itself, the sum of the squared deviations of each of the nII scores in a two-factor design around  $\bar{X}_{m}$  is exactly equal to  $SS_{A} + SS_{B} + SS_{AB} + SS_{m}$ , i.e.,

$$\sum_{i=1}^{J} \sum_{j=1}^{n} (X_{ijk} - \bar{X}_{...})^2 = SS_A + SS_B + SS_{AB} + SS_{\omega}.$$
 (17.8)

The term on the left-hand side of Eq. (17.8) is called the total sum of squares and denoted SStotal for obvious reasons. The total rariance of the collection of nIJ scores is analyzed into four additive components; hence, we use the expression analysis of variance, abbreviated ANOVA or sometimes anova.

#### 17.7 DEGREES OF FREEDOM

Each of the four sums of squares in the two-factor ANOVA is converted into a mean square by dividing it by its degrees of freedom. The degrees of freedom of a sum of squares equals the number of least-squares estimates of effects that comprise the sum of squares less the number of independent linear restrictions placed on these estimates. Admittedly, this is a difficult and abstract notion, and we shall discuss it in some detail here merely for the sake of comprehensiveness of coverage. If you are reading this section for the first time, the day on which you come to "understand" (rather than "work with") the concept of degrees of freedom probably lies far in the future.

 $SS_A$  is calculated from the I least-squares estimates  $\hat{x}_1, \ldots, \hat{x}_I$ . It was natural and unrestrictive to specify in the model in Eq. (17.2) that  $\alpha_1+\ldots$  $+\alpha_I=0$ . Furthermore, although we did not see it happen, it was necessary to assume that  $\hat{a}_1 + \ldots + \hat{a}_I = 0$  before the solution to the mathematical criterion of least-squares estimation could be found. Indeed, as they must, the least-squares estimates of the  $\alpha_i$  satisfy this restriction, i.e.,  $\hat{\alpha}_1 + \ldots + \hat{\alpha}_I = 0$ . This is easy to demonstrate:

$$\sum_{i=1}^{I} \hat{x}_i = \sum_{i=1}^{I} (\vec{X}_{i..} - \vec{X}_{...}) = 0,$$

because  $\widetilde{X}_{...}$  is the mean of the I column means  $\overline{X}_{t...}$ 

There are I least-squares estimates in the calculation of  $SS_A$  and they must conform to the single linear restriction that their sum be zero. Hence, the degrees of freedom for  $SS_A$  are I-1. An exactly analogous line of reasoning would lead to the correct conclusion that  $SS_B$  has degrees of

The calculation of  $SS_{AB}$  involves the II least-squares estimates of the  $\alpha\beta_{ij}$  terms. The restrictions it was necessary to impose on these estimates to solve the least-squares problem were that summing the estimates across rows for any given column yields a sum of zero and summing the estimates across the columns for any given row yields a sum of zero, i.e.,

$$\sum_{i=1}^{J} \widehat{x} \widehat{\beta}_{ij} = 0 \text{ for each } i$$
(17.9)
$$\sum_{i=1}^{J} \widehat{x} \widehat{\beta}_{ij} = 0 \text{ for each } j.$$
(17.10)

$$\sum_{i=1}^{n} \widehat{z} \widehat{\beta}_{ij} = 0 \quad \text{for each } j. \tag{17.10}$$

The conditions in Eq. (17.9) are I in number; there are J restrictions represented in Eq. (17.10). Not all I+J of these restrictions are independent, however. Namely, given the restrictions in Eq. (17.9) and knowing that  $\sum_{i=1}^{J} \widehat{\alpha} \widehat{\beta}_{ij}$  equals zero for  $j=1,\ldots,J-1$ , it must necessarily follow that  $\sum_{i=1}^{J} \widehat{\alpha} \widehat{\beta}_{ij} = 0$ . Hence only I+J-1 of the linear restrictions on the IJ values of  $\widehat{\alpha} \widehat{\beta}_{ij}$  are independent. Therefore, the degrees of freedom for  $SS_{AB}$  are

$$IJ - (I + J - 1) = IJ - I - J + 1.$$

Notice that this expression "factors" into (I-1)(J-1).

The sum of squares within cells,  $SS_{\omega_1}$  is actually the sum of the squares of the nU least-squares estimates of the  $e_{ik}$  terms in the model in Eq. (17.2). Any single  $e_{ik}$  is estimated by  $\hat{e}_{ik} = X_{ik} - \bar{X}_{ik}$ .

Since  $\hat{e}_{ijk}$  is the deviation of a score from its cell mean, the sum of the n values of  $\hat{e}_{ijk}$  within each cell is zero. Thus, there are IJ independent linear restrictions on the nIJ values of  $\hat{e}_{ijk}$ . Consequently, the degrees of freedom associated with  $SS_n$  are IJn - IJ = II(n - 1).

The above results can be summarized as follows:

Sum of squares	Degrees of freedo		
SSA	I-1		
$SS_B$	J-1		
$SS_{AB}$	(I-1)(J-1)		
SS,	IJ(n-1)		

Now look at a very simple example of linear restriction. How many independent restraints on the data of the  $2 \times 3$  table in Fig. 17.11 are there?

Row Column number			Row	
number	1	2	3	sums
1	1			7
2		4		9
Column sums	4	6		

FIG. 17.11

The conditions in Eq. (17.9) are I in number; there are J restrictions represented in Eq. (17.10). Not all I+J of these restrictions are independent, however. Namely, given the restrictions in Eq. (17.9) and knowing that  $\sum_{i=1}^{J} \widehat{\alpha} \widehat{\beta}_{ij} \text{ equals zero for } j=1,\ldots,J-1, \text{ it must necessarily follow that } \sum_{i=1}^{J} \widehat{\alpha} \widehat{\beta}_{iJ} = 0.$  Hence only I+J-1 of the linear restrictions on the JJ values of  $\widehat{\alpha} \widehat{\beta}_{iJ}$  are independent. Therefore, the degrees of freedom for  $SS_{AB}$  are

$$IJ - (I + J - 1) \approx IJ - I - J + 1.$$

Notice that this expression "factors" into (I-1)(J-1).

The sum of squares within cells,  $SS_{er}$ , is actually the sum of the squares of the nU least-squares estimates of the  $e_{th}$  terms in the model in Eq. (17.2). Any single  $e_{th}$  is estimated by  $\hat{e}_{th} = X_{th} - X_{tt}$ .

Since  $\ell_{iR}$  is the deviation of a score from its cell mean, the sum of the n values of  $\ell_{iR}$  within each cell is zero. Thus, there are II independent linear restrictions on the nII values of  $\ell_{iR}$ . Consequently, the degrees of freedom associated with  $SS_n$  are III - II = II(n-1).

The above results can be summarized as follows:

Sum of squares	Degrees of freedo.		
SSA	<i>I</i> – 1		
$SS_n$	J-1		
$SS_{AB}$	(I-1)(J-1)		
SS	IJ(n-1)		

Now look at a very simple example of linear restriction. How many independent restraints on the data of the  $2 \times 3$  table in Fig. 17.11 are there?

Row	Column number		Row	
number	,	2	3	sums
1	1			7
2		4		9
Column	4	6		

FIG. 17.11

(Try filling in the four missing cell entries, the missing column sum, and the missing grand sum.)

If the number of degrees of freedom is the number of cells (here,  $2 \times 3 = 6$ ) minus the number of independent linear restrictions on the data, how many degrees of freedom are there for a table of this type? That is, how many cell entries are free to vary? Is this analogous to the degrees of freedom for the interaction of factor A with factor B in a  $2 \times 3$  factorial design? [Hint: For that ANOVA, the cell entries are interaction residuals, the six  $(X_{1i}, -X_{1i}, -X_{2i}, +X_{ii})$ 's, and every row sum and column sum is zero.]

#### 17.8 MEAN SQUARES

For each sum of squares there is a mean square (abbreviated MS) defined by dividing the sum of squares by its degrees of freedom:

$$MS_A = \frac{SS_A}{l-1}$$

$$MS_B = \frac{SS_B}{J-1},$$

$$MS_{AB} = \frac{SS_{AB}}{(l-1)(J-1)},$$

$$MS_{\varphi} = \frac{SS_{\varphi}}{Ll(p-1)}.$$

As in the one-factor ANOVA, the mean squares are the final stage in calculations leading toward significance tests of the null hypotheses.

## 17.9 COMPUTATIONAL PROCEDURES

In this section we shall see how computational formulas are applied in finding the sums of squares and mean squares. For a two-factor ANOVA design in which factor A has J levels, factor B has J levels, and each of the IJ cells contains n observations, the four sums of squares can be obtained most conveniently from the formulas in Table 17.2.

From a population of 2500 tenth-grade geometry classes 48 were selected at random for use in an experiment. The researcher wished to determine the efficiency of two different methods and media for geometry instruction and their interactions. Classes were randomly assigned in equal numbers to the four combinations of media and method. After one semester

# **TABLE 17.2**

## COMPUTATIONAL FORMULAS FOR SUMS OF SQUARES IN THE TWO-FACTOR ANOVA WITH EQUAL n's

$$SS_{a} = \sum_{i=1}^{J} \frac{\left(\sum_{i=1}^{J} \sum_{k=1}^{n} X_{i0}\right)^{k}}{nJ} - \frac{\left(\sum_{i=1}^{J} \sum_{k=1}^{n} X_{i0k}\right)^{k}}{nIJ}$$

$$SS_{B} = \sum_{j=1}^{J} \frac{\left(\sum_{i=1}^{n} \sum_{k=1}^{n} X_{i0}\right)^{k}}{nI} - \frac{\left(\sum_{i=1}^{J} \sum_{j=1}^{n} X_{i0k}\right)^{k}}{nIJ}$$

$$SS_{AB} = \sum_{j=1}^{J} \frac{\left(\sum_{i=1}^{n} \sum_{k=1}^{n} X_{i0}\right)^{k}}{nIJ} - SS_{A} - SS_{B} - \frac{\left(\sum_{i=1}^{J} \sum_{j=1}^{n} X_{i0k}\right)^{k}}{nIJ}$$

$$SS_{AB} = \sum_{i=1}^{J} \sum_{j=1}^{n} \sum_{k=1}^{n} X_{i0k}^{i} - \sum_{i=1}^{J} \sum_{j=1}^{J} \frac{\left(\sum_{k=1}^{n} X_{i0k}\right)^{k}}{nIJ}$$

$$SS_{B} = \sum_{i=1}^{J} \sum_{j=1}^{J} \sum_{k=1}^{n} X_{ijk}^{i} - \sum_{i=1}^{J} \sum_{j=1}^{J} \frac{\left(\sum_{k=1}^{n} X_{i0k}\right)^{k}}{n}$$

of instruction the researcher administered the same geometry achievement test to each class. Since the sampling unit was "classroom," the unit of analysis is taken to be the class mean (to the nearest whole number) on the criterion achievement test. The results are tabulated in Fig. 17.12.\*

In the above example, i = 1, 2; j = 1, 2; and  $k = 1, \ldots, 12$ . Certain basic calculations are indicated in Fig. 17.12. In addition, one more quantity is required: the sum of all 48 scores after each has been squared. This quantity is:

$$\sum_{1}^{2} \sum_{k=1}^{2} \sum_{k=1}^{12} X_{ijk}^{2} = 2^{2} + 5^{2} + \dots + 35^{2} = 14,969.$$

Only those quantities in Fig. 17.12 are required to find the four sums of squares by the formulas in Table 17.2.

The sum of squares for factor A,  $SS_A$ , is calculated as follows:

$$SS_A = \frac{(229)^2 + (506)^2}{2 \cdot 12} - \frac{(735)^2}{2 \cdot 2 \cdot 12} = 1598.52.$$

The remaining sums of squares are found as indicated hereafter.

$$SS_B = \frac{(235)^2 + (500)^2}{2 \cdot 12} - \frac{(735)^2}{2 \cdot 2 \cdot 12} = 1463.02.$$

\* Twenty-four score points have been subtracted from each class mean to make the calculations less arduous on a desk calculator. Recall that  $s_{(x^*,c)}^* = s_s^*$ , so the sums of squares and the mean squares of an analysis of variance are not affected by adding a constant to each observation. However, the mean of (X + c) is X + c, so every mean in the ANOVA has been reduced 24 points.

Factor B, media

	Classroom lecture (1)	Programmed instruction (2)	Row sums
Traditional (1)	2, 5, 6, 7, 4, 6 7, 8, 4, 6, 7, 10	9, 12, 14, 15, 10, 13 14,16, 10, 13, 14, 17	$\sum_{j=1}^{2} \sum_{k=1}^{12} X_{1jk} = 229$
A, methods	$\sum_{k=1}^{12} X_{11k} = 72$	∑ X <sub>12k</sub> = 157	•
actor	10,13,14,16,10,13	21, 25, 31, 33, 22, 26 32, 34, 22, 30, 32, 35	$\sum_{i=1}^{2} \sum_{k=1}^{12} X_{2/k} = 506$
Modern (2)	$\sum_{k=1}^{12} X_{21k} = 163$	$\sum_{k=1}^{12} X_{22k} = 343$	j=1 k=1
Column sums	$\sum_{i=1}^{2} \sum_{k=1}^{12} X_{i1k} = 235$	$\sum_{i=1}^{2} \sum_{k=1}^{12} X_{i2k} = 500$	$\sum_{i=1}^{2} \sum_{j=1}^{2} \sum_{k=1}^{12} X_{ijk} = 735$

FIG. 17.12 Illustration of basic calculations in a two-way analysis of

$$SS_{AB} = \frac{(72)^2 + (157)^2 + (163)^2 + (343)^2}{12} - SS_A - SS_B - \frac{(735)^2}{2 \cdot 2 \cdot 12}$$

$$= \frac{(72)^2 + (157)^2 + (163)^2 + (343)^2}{12} - 1598.52 - 1463.02 - \frac{(735)^2}{2 \cdot 2 \cdot 12}$$

$$= 188.02.$$

$$SS_{\bullet} = 14,969 - \frac{(72)^2 + (157)^2 + (163)^2 + (343)^2}{12} = 464,75.$$

The sums of squares, degrees of freedom, and mean squares are reported in Table 17.3.

TABLE IT.]

SUHS OF SQUARES, DEGREES OF FREEDOM, AND MEAN SQUARES FOR THE

Source of praintane

Source of praintane

Factor A (method)	df	55	MS
Factor B (media) Interaction of A and B Within cells	I-1 = 1  J-1 = 1  (I-1)(J-1) = 1  IJ(n-1) = 44	1598.52 1463.02 188.02 464.75	1598.52 1463.02 188.02 10.56

#### 17.10 EXPECTED VALUES OF MEAN SQUARES

The computational aspects of the two-factor ANOVA are complete. Now it is time to turn our attention once again to the purpose of the computations. We are seeking a statistical inferential test for deciding whether the data support the null hypothesis  $H_0$  or the alternative hypothesis  $H_1$  about the main effects of factors A and B and their interaction effects. As was true in the one-factor ANOVA, the expected values of the mean squares reveal how they bear on the truth or falsity of the three null hypotheses.

The expected value (or "long-run average value") of  $MS_w$  is the mean of all the  $MS_w$ 's that would be obtained if the same two-factor ANOVA design were performed an infinite number of times with independent observations. Another way to look at  $E(MS_w)$  is that it is the variance of the population from which the observations in any one cell of the two-factor ANOVA design have been sampled. We shall assume that the variance of the population from which the n observations in any cell have been sampled is equal to  $\sigma_v^*$ . In other words, the variances of each of the populations underlying each of the IJ cells are equal to the same value,  $\sigma_v^*$ . This is an extension to the two-factor ANOVA of the assumption of homogeneous variances that we saw in the one-factor ANOVA.

If the *n* observations in the *ij*th cell are assumed to have been drawn from a population with variance  $\sigma_s^2$ , then  $E(s_{is}^2) = \sigma_s^2$ .  $MS_w$  has the following form:

$$MS_{w} = \frac{SS_{w}}{IJ(n-1)} = \frac{\sum_{j}^{I} \sum_{k}^{J} \sum_{k}^{n} (X_{ijk} - \vec{X}_{ij})^{2}}{IJ(n-1)}.$$

We recognize  $MS_w$  to be the average of the IJ within cell sample variances, i.e.,

$$MS_w = \frac{\sum_{i=1}^{J} \sum_{j=1}^{J} s_{ij}^2}{\prod_{i=1}^{J} s_{ij}^2}.$$

We see, then, that

$$E(MS_w) = E\left[\frac{\sum_{i=1}^{J} \sum_{j=1}^{J} s_{ij}^2}{JI}\right] = \sum_{i=1}^{J} \sum_{j=1}^{J} \frac{E(s_{ij}^2)}{IJ} = \sum_{i=1}^{J} \sum_{j=1}^{J} \frac{\sigma_s^2}{IJ} = \frac{IJ\sigma_s^2}{IJ} = \sigma_s^2.$$

The expected value of  $MS_A$  is the average of an infinite number of  $MS_A$ 's, each one obtained from an independent replication of the same two-factor ANOVA design—I levels of factor A, J levels of factor B, and n cases in each of the IJ cells. In the  $2 \times 2$  experiment in the preceding section, the value of  $MS_A$  was 198.52. This is just one observation from a hypothetically infinite

population of  $MS_A$ 's that could be generated by replicating the same methods  $\times$  media experiment with a new set of 48 classrooms randomly drawn from the population. We don't know whether 1598.52 is above or below the population average value of  $MS_A$ , i.e., the expected value of  $MS_A$ ,  $E(MS_A)$ .

From one replication of the experiment, we can calculate one value of  $MS_A$ . However, we cannot calculate the numerical value of  $E(MS_A)$ ; if we could there would be no need for inferential statistics. The algebraic formula for  $E(MS_A)$  in terms of the parameters of the model in Eq. (17.2) can be found. We shall not bother with the details of the derivation here; we shall simply state that  $E(MS_A)$  has the following form:

$$E(MS_A) = \sigma_e^2 + \frac{nJ \sum_{i}^{J} \alpha_i^2}{I - 1},$$
 (17.11)

where  $\sigma_e^2$  is the variance of the error term in Eq. (17.2) and is estimated by  $MS_e$ , and

 $\alpha_i$  is the main effect of the *i*th level of factor A, i.e.,  $\alpha_i = \bar{\mu}_i - \mu$ .

Suppose that, unknown to the researcher, the true value of  $\sigma_{\star}^{z}$  (the true "within cell" variance) is 15.0, and that  $\bar{\mu}_{1}$ , = 12 and  $\bar{\mu}_{2}$ , = 22. Since  $\mu = \frac{1}{2}(12+22) = 17$ ,  $\alpha_{1}$  is 12 - 17 = -5 and  $\alpha_{2}$  is 22 - 17 = 5. Substituting these values and J = 2 and n = 12 into Eq. (17.11) yields

$$E(MS_A) = 15.0 + 12 \cdot 2 \frac{[(-5)^2 + 5^2]}{2 - 1} = 1215.$$

The above calculations were performed to illustrate the nature of the terms in Eq. (17.11). It must be emphasized that one never actually calculates a value for  $E(MS_A)$ . What is important is to note the relationship between the expression for  $E(MS_A)$  and the truth or falsity of the null hypothesis about factor A. Notice that the third of several equivalent statements of  $H_0$  for factor A in Sec. [1.2, 5 is  $H_0$ :  $\sum_{i=1}^{L} z_i^2 = 0$ . The quantity hypothesized to be zero in  $H_0$  for factor A is the same quantity,  $\sum_{i=1}^{L} z_i^2$ , that appears in the numerator of the second term for  $E(MS_A)$ . Thus, if  $H_0$  is true—which means that  $\sum_{i=1}^{L} c_i - t_{i+1}$ 

$$E(MS_A) = \sigma_e^2 + \frac{nJ(0)}{I-1} = \sigma_e^2.$$

On the other hand, if  $H_0$  is false—which means that  $\sum \alpha_i^2$  is positive—then

$$E(MS_A) = \sigma_s^2 + \frac{nJ \sum \alpha_t^2}{1 - 1} > \sigma_s^2.$$

This is an important relationship to understand: If He is true, we expect

 $MS_A$  to be the same size as the true within-cell variance,  $\sigma_e^2$ ; if  $H_0$  is false, we expect  $MS_A$  to be larger than  $\sigma_e^2$ .

The expected value of MSn has the form

$$E(MS_B) = \sigma_e^2 + \frac{nI \sum_{j}^{J} \beta_j^2}{J-1}.$$
 (17.12)

The null hypothesis  $H_0$  for the main effect of factor B is  $H_0$ :  $\int_1^J \beta_i^2 = 0$ . If  $H_0$  for factor B is true, then

$$E(MS_B) = \sigma_e^2 + \frac{nI(0)}{I-1} = \sigma_e^2.$$

If  $H_0$  for factor B is true, we expect  $MS_B$  to be equal to  $E(MS_m) = \sigma_s^2$ ,  $MS_B$  can be expected to be larger than  $\sigma_s^2$  when  $H_0$  for factor B is false.

The expected value of MSAR is

$$E(MS_{AB}) = \sigma_{\epsilon}^{2} + \frac{n \sum_{i}^{J} \sum_{j}^{J} \alpha \beta_{ij}^{2}}{(I - 1)(J - 1)}$$
 (17.13)

The null hypothesis for the interaction of factors A and B can be stated as  $H_0$ :  $\sum_{i}^{J} \sum_{j}^{J} \alpha \beta_{ij}^2 = 0$ . Thus, we see that if  $H_0$  for the interaction of A and B is true, then  $E(MS_{AB})$  equals  $\sigma_e^2$ : if  $H_0$  is false, then  $E(MS_{AB}) > \sigma_e^2$ . All of these important relationships are summarized in Table 17.4.

TABLE 17.4 RELATIONSHIPS BETWEEN NULL HYPOTHESES AND EXPECTED VALUES OF

MEAN SQUARES		
Mean square	Expected mean square when H <sub>0</sub> is true for source of variation in question	Expected mean square when H <sub>0</sub> is false
Factor A, MSA	o <sub>s</sub>	$\sigma_s^{z} + \frac{nf\sum \alpha_i^{z}}{I-1}$
Factor B, MS <sub>B</sub>	σ;	$\sigma_{\bullet}^3 + \frac{nI \sum \beta_{f}^3}{J-1}$
Interaction of A and B, MSAR	σ.*	$\sigma_s^2 + \frac{n \sum \sum \alpha \beta_{ii}^2}{(I-1)(I-1)}$ $\sigma_s^2$
Within cells, MS.	$\sigma_s^s$	o. 1)(0 1)

As we shall see in greater detail later, the comparison of  $MS_A$  with  $MS_\omega$  reflects on the truth of  $H_0: \sum x_i^2 = 0$ . If  $H_0$  is true, then  $MS_A$  and  $MS_\omega$  have the same expected value; if  $H_0$  is false, then  $MS_A$  has an expected value larger

than  $\sigma_s^2$  but the expected value of  $MS_\omega$  is still  $\sigma_s^2$ . Naturally, if  $MS_A$  proves to be much larger than  $MS_\omega$  in a particular run of the experiment, we are inclined to think that  $H_s$  is false; if  $MS_A$  and  $MS_\omega$  are much the same size in a particular replication of the experiment, we are inclined to think that they are both estimating the same quantity,  $\sigma_s^2$ , which is the case when  $H_s$  is true. Comparisons of either  $MS_B$  or  $MS_A$ , with  $MS_\omega$  bear on the truth or falsity of the null hypotheses about the main effects of B and the interaction effects of A and B, respectively, in the same manner that comparing  $MS_A$  with  $MS_\omega$  tells us something about the plausibility of  $\sum \sigma_s^2 = 0$  being true.

The problem of deciding when  $MS_A$   $(MS_B$  or  $MS_{AB})$  is sufficiently larger than  $MS_a$  so that we should consider the truth of  $H_0$  implausible is a problem of the variability of the values of mean squares from one replication of the two-factor experiment to the next. This problem will occupy our attention in the next section.

# 17.11 THE DISTRIBUTIONS OF THE MEAN SQUARES

Before proceeding to the question of the statistical distributions of the four mean squares in a two-factor ANOVA design, it is advisable to clarify the nature of the statistical inference one makes in this situation. In the twofactor design in Sec. 17.9, 12 observations were taken in each of the  $2 \times 2 = 4$ cells of the design. These 48 observations can be considered to be randomly drawn from four hypothetical populations (one for each cell) that contain the scores an infinite number of classrooms would obtain under the same experimental conditions. The 48 observations in Fig. 17.12 will be called a complete replication of the experiment. This complete replication produced the following mean squares:  $MS_A = 1598.52$ ,  $MS_B = 1463.02$ ,  $MS_{AB} = 1663.02$ 188.02, and  $MS_w = 10.56$ . A second complete replication of the experiment could be obtained by performing the same experiment with a different set of 48 classrooms (12 in each cell); this second replication would yield different values for each of the four mean squares. Conceptually at least, third, fourth, fifth, etc. replications of the experiment could be run, and each replication would produce its own set of four mean squares. Now the question is, what will be the distribution of values of  $MS_A$  obtained from an infinite number of replications of the experiment? We ask the same question individually about MSB, MSAB, and MSw.

Before these questions can be answered, it is necessary to add an assumption to our model in Eq. (17.2). You will recall that the model in Eq. (17.2) itself an assumption and that in Sec. 17.10 it was necessary to assume that the variances of the hypothetical populations underlying the II cells of the experiment all have the same variance,  $\sigma^T$ . To these assumptions we now add the assumption that these populations are normally distributed.

#### The Distribution of MS.

With the addition of the normality assumption, we know that the n observations in any cell-the ijth cell-constitute a random sample from a normal distribution with mean  $\mu_{ij}$  and variance  $\sigma_{ij}^2$ . Each cell variance  $s_{ij}^2$  is an unbiased estimator of  $\sigma_s^2$ ; furthermore, we can conclude that

$$\frac{s_{ij}^2}{\sigma_e^2} \sim \frac{\chi_{n-1}^2}{n-1}$$

that is,  $s_{ij}^2/\sigma_s^2$  has a distribution equal to the chi-square distribution (df =n-1) divided by n-1. This statement is true for the II independent cell variances  $s_{ij}^2$ . From the additive property of chi-square variables, we know that

$$\sum_{i=1}^{J} \sum_{j=1}^{S_{i,j}^2} \sim \frac{\chi_{n-1}^2}{n-1} + \ldots + \frac{\chi_{n-1}^2}{n-1} \sim \frac{\chi_{IJ(n-1)}^2}{n-1}.$$

Dividing the above quantities by IJ yields:

$$\frac{\sum\limits_{i=1}^{I}\sum\limits_{j=1}^{J}s_{ij}^2}{IJ\sigma_{\epsilon}^2} = \frac{MS_w}{\sigma_{\epsilon}^2} \sim \frac{\chi_{IJ(n-1)}^2}{IJ(n-1)}.$$

 $MS_w/\sigma_s^2$  has a chi-square distribution, df = IJ(n-1), divided by IJ(n-1). For example, in the problem in Sec. 17.9, I=J=2 and n=12. Suppose the value of  $\sigma_e^2$  is 15. Then

$$\frac{MS_w}{15} \sim \frac{\chi_{44}^2}{44}$$

and the single observed value of  $MS_w = 10.56$  over 15 is one observation from a chi-square distribution (df = 44) that has been rescaled by division by 44.

#### The Distribution of MSA

We must consider two cases in this instance: the distribution of  $MS_A$  when  $H_0$ :  $\sum \alpha_i^2 = 0$  is true and when  $H_0$  is false. If  $H_0$  is true, then

$$f_0: \sum \alpha_i^2 = 0$$
 is true and when  $H_0$  is true.
$$\frac{MS_A}{\sigma^2} \sim \frac{\chi_{I-1}^2}{I-1},$$

$$\frac{MS_A}{\sigma_a^2} \sim \frac{\chi_{I-1}^2}{I-1}$$

that is,  $MS_A/\sigma_e^2$  has a distribution over complete replications of the twofactor design that is the chi-square distribution (df = I - 1) divided by

If  $H_0$  is false, then  $MS_A/\sigma_s^2$  has what is called a noncentral chi-square distribution (df = I - 1) divided by I - 1. The noncentral chi-square distribution with df = I - 1 is a mathematical curve that has a higher mean




FIG. 17.13 Distribution of MS. Io.

and is generally to the right of the chi-square distribution (df = I - 1). The relationship of the chi-square distribution to the noncentral chi-square distribution is illustrated in Fig. 17.13.

When  $H_0$  is false, the values of  $MS_A$  tend to be larger than those of  $MS_A$  when  $H_0$  is true. This is reflected in the displacement to the right of the non-central chi-square distribution. The larger the value of  $\sum \alpha_i^2$ , the further to the right of  $\chi_{I-1}^2/(I-1)$  the values of  $MS_A/\sigma_i^2$  will be displaced. Thus, the noncentral chi-square distribution in Fig. 17.13 is for one particular value of  $\sum \alpha_i^2$  only; there exists a separate noncentral  $\chi^2$  distribution for each value of  $\sum \alpha_i^2$ .

#### The Distribution of MS.

The distributional statements that can be made about  $MS_B$  are quite analogous to those for  $MS_A$ , as you might expect.

If  $H_0: \sum \beta_1^2 = 0$  is true, then

$$\frac{MS_B}{\sigma_*^2} \sim \frac{\chi_{J-1}^2}{J-1}.$$

If  $H_0$  is false, then  $MS_B/\sigma_r^2$  has a noncentral chi-square distribution (df = J - 1) divided by J - 1.

The Distribution of MSAB

If  $H_0$ :  $\sum \sum \alpha \beta_{ij}^a = 0$ , i.e., if there is no interaction between factors A and B, then

$$\frac{MS_{AB}}{\sigma_4^2} \sim \frac{\chi_{(I-1)(J-1)}^2}{(I-1)(J-1)},$$

that is,  $MS_{AB}/\sigma_s^2$  has a chi-square distribution, df = (I-1)(J-1), divided by (I-1)(J-1).

Again, if the null hypothesis about the interaction of A and B is false,  $MS_{AB}/\sigma_1^2$  has a noncentral chi-square distribution, df = (I - 1)(J - 1), divided by (I - 1)(J - 1).

We are now prepared to combine the above facts into the major results of this section. Recall that the ratio of two independent chi-square variables, each divided by its own degrees of freedom, has an F-distribution.

Suppose that  $H_0$ :  $\sum \alpha_t^2 = 0$  is true; then  $MS_A/\sigma_t^2 \sim \chi_{t-1}^2/(t-1)$ . Regardless of whether  $H_0$  is true or false,  $MS_w/\sigma_t^2$  has a chi-square distribution divided by IJ(n-1). Now

$$\frac{MS_A/\sigma_o^2}{MS_n/\sigma_o^2} \sim F_{I-1,IJ(n-1)}.$$

But notice that

$$\frac{MS_A/\sigma_e^2}{MS_-/\sigma_e^2} = \frac{MS_A}{MS_m} \sim F_{I-1,IJ(n-1)}.$$

(Luckily we have eliminated σ<sup>2</sup> without having to know its actual value.)

What we have shown is that the ratio of  $MS_A$  to  $MS_a$  has an F-distribution with I-1 and II(n-1) degrees of freedom when  $H_0$  is true. Thus if the value of  $MS_A/MS_a$  for a replication of a two-factor experiment looks like a "typical" observation from the distribution  $F_{I-1,II(n-1)}$ —by "typical" we mean that  $MS_A/MS_a$  does not exceed the 90th or 95th or 99th percentile, say, of that distribution—we are inclined to think that  $H_0$  is true. On the other hand, if  $H_0$  is false, then we expect the value of  $MS_A$  to be larger than the value of  $MS_a$ . Hence, if we obtain a very large value of  $MS_A/MS_a$ —a value that seems not to have been drawn from  $F_{I-1,II(n-1)}$  since it exceeds the 99th percentile, say, of that distribution—then we think that  $H_0$  is probably false.

When  $H_0$ :  $\sum \beta_i^2 = 0$  is true,

$$\frac{MS_B/\sigma_e^2}{MS_m/\sigma_e^2} = \frac{MS_B}{MS_m} \sim F_{J-1,IJ(n-1)}.$$

When  $H_0: \sum \sum \alpha \beta_{ij}^2 \approx 0$  is true, then

$$\frac{MS_{AB}/\sigma_e^2}{MS_{\omega}/\sigma_e^2} = \frac{MS_{AB}}{MS_{\omega}} \sim F_{(I-1)(J-1),IJ(n-1)}.$$

The effect of a false null hypothesis about either the main effects of factor B or the interaction effects of A and B is to increase  $MS_B$  or  $MS_{AB}$  without systematically increasing  $MS_{o}$ , thus producing a distribution, over complete replications of the design, of ratios of mean squares that is displaced to the right of the F-distribution. These relationships are illustrated for factor B in Fig. 17.14.




FIG. 17.14 Distribution of  $MS_s/MS_s$  over complete replications of the two-factor design when  $H_s$  is true and when it is false. (Shaded portion contains the 5% of the ratios that exceed the 95th percentile in the F-distribution.)

#### 17.12 HYPOTHESIS TESTS OF THE NULL HYPOTHESES

The discussion to this point has led us to three ratios of mean squares that will be called F-ratios:

$$F_A = \frac{MS_A}{MS_w}, \qquad F_B = \frac{MS_B}{MS_w}, \qquad F_{AB} = \frac{MS_{AB}}{MS_w}.$$

For the data in Table 17.3 these F-ratios have the following values:

$$F_A = \frac{1598.52}{10.56} = 151.38, F_B = \frac{1463.02}{10.56} = 138.54,$$

$$F_{AB} = \frac{188.02}{10.56} = 17.80.$$

We shall now point out how statistical tests of the three null hypotheses can be made using the F-ratios. These F-tests are similar to the F-test in the one-factor fixed-effects ANOVA. The F-test will be illustrated with the main effects of factor A.

Let's go back to the example in Fig. 17.12 to illustrate the hypothesis tests. There we found that  $F_4=151.38$ . If  $H_6:\sum \alpha_i^2=0$  were true, the distribution of  $F_A$  over repeated complete replications of the  $2\times 2$  design would describe an F-distribution with I-1=1 and II(n-1)=4 degrees of freedom. We would not wish to conclude erroneously that  $H_0$  is false when in fact it is true. Indeed, we wish to adopt a decision rule for choosing between  $H_0$  and  $H_1:\sum \alpha_i^2\neq0$  that will lead us to decide erroneously to reject  $H_0$  in favor of  $H_1$  only one time in 100, say. Hence, we want to adopt a risk of  $\alpha=0$ 1 of committing a type I error, rejecting  $H_0$  when it is true. Since the only evidence in favor of  $H_1$  is a large value of  $F_A$ , we shall place the entire critical region of the test in the upper tail of the distribution  $F_{1,44}$ : hence, the critical value for the test becomes  ${}_{10}F_{1,44}$ . From Table E in Appendix A, we can determine that the 99th percentile in the F-distribution with 1 and 44 degrees of freedom is approximately 7.25.

Any F-ratio  $F_A$  exceeding 7.25 will be taken to be evidence that the hypothesis  $H_0$ :  $\sum a_i^2 = 0$  is false. This statement constitutes the decision rule of the hypothesis test. If in reality  $H_0$  is true, this decision rule will have a probability of  $\alpha = .01$  of falsely rejecting  $H_0$ . Such is the magnitude of the risk one takes in agreeing to reject  $H_0$  if  $F_A$  is greater than 7.25.

For the data in Fig. 17.12 the value of  $F_A$  is 151.38. Since this F-ratio exceeds the critical value of 7.25, the hypothesis that  $\sum \alpha_i^2 = 0$  is rejected. One concludes that the two sample means  $\vec{X}_{1...}$  and  $\vec{X}_{2...}$  are significantly different at the .01 level.

The F-test of the hypothesis  $H_0\colon \sum \beta_i^2=0$  proceeds along similar lines. Suppose it was decided to test  $H_0$  with an  $\alpha$  of .05. From Table E we see that the 95th percentile in the F-distribution with J-1=1 and J(I-1)=4 degrees of freedom is 4.06. Hence, if  $H_0$  is true, the chances of obtaining a value of  $F_D$  exceeding 4.06 are 1 in 20. The chances of obtaining an  $F_D$  of 138.54 or greater when  $\sum \beta_i^2=0$  are infinitesimally small. Thus, rather than regard  $F_D$  as a one-in-a-million occurrence that just so happened to occur even though  $H_0$  was true, we take the more logical course and regard it as evidence that  $H_0$  is false. Our conclusion is stated formally as " $H_0\colon \sum \beta_i^2=0$  is rejected with an  $\alpha$  of .05." Of course, this implies that we are deciding in favor of the conclusion that  $\beta_1\neq\beta_3$ , i.e., that the means for the two media (classroom lecture vs. programmed instruction) of the populations sampled are different.

The *E*-test of the null hypothesis about the interaction of factors A and B is carried out in a similar manner. If  $H_0\colon \sum \alpha \beta_B^n = 0$  is true, then  $F_{AB}$  has an F-distribution with (I-1)(I-1) = 1 and II(I-1) = 44 degrees of freedom. Suppose that we wish to run a test of  $H_0$  with a risk of .01 of rejecting  $H_0$  when it is actually true. The critical value that  $F_{AB}$  must exceed to be counted as evidence of a false  $H_0$  is 7.25, the 99th percentile in the F-distribution with 1 and 44 degrees of freedom.

The value of  $F_{AB}$  is 188.02/10.56 = 17.80. The position of this obtained




FIG. 17.15 Position of an F-ratio of 17.80 relative to the F-distribution with one and 44 degrees of freedom.

F-ratio relative to the F-distribution with 1 and 44 df—the distribution  $F_{AB}$  would follow if  $H_0$  were true—is illustrated in Fig. 17.15.

By inspecting Fig. 17.15, it would appear to be foolish to regard 17.80 as having been drawn at random from the distribution  $F_{1.41}$ , but to argue that the data support the null hypothesis that  $\sum \sum a\beta_{ij}^2 = 0$  would be equivalent to doing so. Thus, the conclusion is to reject  $H_0$ , and the probability that this conclusion is to recovery is a far less than .01.

Since a significant interaction of A and B has been found, it will be illuminating to graph it. The four cell means for the data in Fig. 17.12 are 6.00 for traditional method and classroom lecture, 13.08 for traditional method and programmed instruction, 13.58 for modern method and classroom lecture, and 28.58 for modern method and programmed instruction. The four cell means are depicted in Fig. 17.16.

The departure from parallelism of the two lines in Fig. 17.16 reflects the




FIG. 17.16 Graph of the interaction of factors A and B for the data in Fig. 17.12.

interaction of method with media. The departure of the two lines from parallelism was judged by the F-test to be overwhelmingly statistically significant, i.e., there is hardly the slightest chance that graphing the unknown population means would reveal parallel lines. The interpretation of this interaction is that the modern teaching method emphasized the superiority of programmed instruction over the classroom lecture over what that superiority was for the traditional method. Programmed instruction is about seven points better than classroom lecture under the traditional method; but it is 15 points better with the modern method. Perhaps the modern method lent itself to programming better than did the traditional method.

# 17.13 REVIEW OF THE TWO-FACTOR ANOVA WITH EQUAL NUMBERS OF OBSERVATIONS PER CELL

In outline form, the two-factor fixed-effects model ANOVA with n observations per cell is performed as follows:

- The following model is postulated for the data: X<sub>t/k</sub> = μ + α<sub>i</sub> + β<sub>j</sub> + αβ<sub>ij</sub> + ε<sub>ik</sub>.
   It is assumed that the n observations in any one of the U cells come
- It is assumed that the n observations in any one of the IJ cells come from a normal distribution with variance σ<sub>e</sub><sup>2</sup>, which is constant for each cell, and that the IJ samples are independent of one another.
- A level of significance α is adopted for each of the three F-tests:
   the tests of the null hypothesis for the main effects of factor A and
   factor B and of the interaction effect of A and B.
- The three critical values for the F-tests are determined by reference to Table E in Appendix A. These critical values are as follows:

For 
$$F_A$$
:  $_{1-a}F_{I-1,IJ(n-1)}$ ;  
For  $F_B$ :  $_{1-a}F_{J-1,IJ(n-1)}$ ;  
For  $F_{AB}$ :  $_{1-a}F_{(I-1)(J-1),IJ(n-1)}$ .

- The four mean squares, MS<sub>A</sub>, MS<sub>B</sub>, MS<sub>AB</sub>, and MS<sub>w</sub>, are calculated from the computational formulas for sums of squares in Table 17.2 and the degrees of freedom.
- 6. The three F-ratios are calculated:

$$F_A = \frac{MS_A}{MS_w} \qquad F_B = \frac{MS_B}{MS_w} \qquad F_{AB} = \frac{MS_{AB}}{MS_w} \,.$$

 The F-ratios in step 6 are compared with the corresponding critical values found in step 4. If an F-ratio exceeds the critical value, the corresponding null hypothesis is rejected at the x-level of significance. If an F-ratio does not exceed the critical value, the corresponding null hypothesis is not rejected.

8. The results are tabulated as illustrated in Table 17.4.

TABLE 17.4 TABULATION OF RESULTS OF A TWO-FACTOR FIXED-EFFECTS MODEL ANOVA

Source of cariation	df	MS	F	P
Factor A (methods) Factor B (media) A × B Within ceffs	1 1 1	1598.52 1463.02 188.02 10.56	151.38* 138.54* 17.80*	p < .001 p < .001 p < .001

<sup>•</sup> F-ratio significant at the x = .01 level.

Probability values appear in the last column of Table 17.4. It is becoming customary to report with each F-ratio the proportion, p, of the area in the F-distribution that lies to the right of the obtained F-ratio. For example, if  $F_{AB}$  had been 3.47, one would have indicated that p was between 1.0 and .05, i.e., that less than 10% but more than 5% of the area in the distribution  $F_{AB}$  these above 3.47. One would know immediately that the null hypothesis about interaction could be rejected with an  $\alpha$  of .05.

## 17.14 TWO-FACTOR FIXED-EFFECTS ANOVA WITH UNEQUAL n's

You will probably be surprised to learn that the generalization of the twofactor ANOVA from the equal n's case to that of unequal n's is not simple and straightforward as in the case of the one-factor ANOVA. Of the totality of two-factor layouts of data with unequal n's a subclass of them presents no real difficulties in either theory or application. Unfortunately, the other class does present problems. The remainder of this section is divided into two parts corresponding to the two classes of conditions on the unequal n's.

#### Proportional Cell Frequencies

Now that we are considering the possibility of different cells having different numbers of observations we must adopt a new system of denoting such numbers. To this point, the common number of observations in the II cells of a two-factor design has been denoted by n. Henceforth, the number of

Factor A	Facto		
(1)	1	2	Row sums
1	n <sub>11</sub> = 2	n <sub>12</sub> = 4	n <sub>1.</sub> = 6
2	n <sub>21</sub> ≈ 3	n <sub>22</sub> =6	n <sub>2</sub> = 9
3	n31= 2	n <sub>32</sub> = 4	n <sub>3</sub> = 6
Column sums	n <sub>.1</sub> =7	n,z=14	n = 21

FIG. 17.17 Notation for factorial designs where the  $n_{ij}$ 's are not equal.

observations in the ifth cell of such a design will be denoted by nij. Consider the 3  $\times$  2 design (I = 3, J = 2) for which the n's are shown in Fig. 17.17.

The number of observations in the cell at the intersection of the first column (i = 1) and first row (j = 1) is  $n_{11} = 2$ . There are  $n_{32} = 4$  observations in the cell at the intersection of column three (i = 3) and row two  $(i \approx 2)$ 

It will be necessary eventually to make use of the numbers of observations in any one row or column of a two-factor design. Notice in Fig. 17.17 that there are 2 + 3 + 2 = 7 observations in the first column. This number is found by adding up  $n_{tt}$ 's as follows:

$$n_{.1} = n_{11} + n_{21} + n_{31} = \sum_{i=1}^{3} n_{i1}$$

The notation n, for the total frequencies in the first column conforms to our usage of the subscripted dot to denote having summed over an index. In general, the total frequencies in the jth column is denoted by  $n_{ij}$ . The total number of observations in row 3 is

$$n_{3.} = n_{31} + n_{32} = \sum_{i=1}^{2} n_{3i} = 2 + 4 = 6.$$

In general, there are n, observations in the ith row.

The tetm  $n_{ij}$  denotes  $\sum_{i=1}^{3} \sum_{j=1}^{2} n_{ij}$ , which is the total number of observations in the entire layout of data. We shall also use the familiar notation of N for the total number of observations, i.e., by definition n = N. What is the meaning of n,2 and n1.?

Relatively simple computational procedures may be applied to a twofactor fixed-effects ANOVA with unequal n's provided the cell frequencies are proportional. What does it mean to have proportional cell frequencies? The cell frequencies are proportional when

$$n_{ij} = \frac{n_{i,n,j}}{n_{..}} \tag{17.14}$$

for all II values of  $n_{ij}$ .

In words, the condition of proportionality is satisfied when for each cell the number of observations in that cell equals the product of the numbers of observations in the entire row and entire column in which the cell lies, divided by the total number of observations.\* Notice that this condition is satisfied by each of the six values of  $n_{ij}$  in Fig. 17.17:

$$n_{11} = \frac{6 \cdot 7}{21} = 2$$
,  $n_{21} = \frac{9 \cdot 7}{21} = 3$ ,  $n_{31} = \frac{7 \cdot 6}{21} = 2$ ,  $n_{32} = \frac{6 \cdot 14}{21} = 4$ ,  $n_{32} = \frac{9 \cdot 14}{21} = 6$ ,  $n_{32} = \frac{6 \cdot 14}{21} = 4$ .

Thus the six cell frequencies in Fig. 17.17 are proportional. "Proportional" here means that 2:3:2::4:6:44, and 2:4::3:6::2:4. Both of these are true because the first reduces to "2 is to 3 is to 2, as 2 is to 3 is to 2, and the second reduces to "1 is to 2, as 1 is to 2, as 1 is to 2." Thus the two sets of column frequencies are proportional to each other (or to the column of row sums), and the three sets of row frequencies are proportional to each other (or to the row of column sums).†

Having determined that the cell frequencies are proportional, one may proceed with the calculations as outlined below. The calculations will be illustrated on the data (running long-jump distance measured in feet for boys and girls of three races) in Fig. 17.18.

Notice in the summation signs in Fig. 17.18 that  $n_{ij}$  replaces n in the symbol for the sum of all of the observations in the ijth cell, i.e., the sum

Where did we get Eq. (17.14)? Recall from Chapter 10 the formula for the probability of the currence of both of two independent events,  $(P_n/P_n)$ , where  $P_n$  is the probability of occurrence of event A, and  $P_n$  is the probability of occurrence of event in row t and  $P_n$  is the proportion of the  $n_n$  cases that lie in column f, then it will equal  $(n_n/n_n/n_n)$ . Solve the equation for  $n_n$  taset and you have Eq. (17.14).

Nexed another way, but equivalently, when the  $n_1$ 's are proportional one can consider each as being composed of a proportional prostant for the row,  $r_1$ , and a proportional prostant for the row,  $r_2$ , and a the three row constant or  $r_2 = r_2 = r_3 = r_4$ . For the data of Fig. 17.17  $r_2 = r_3 = r_4 = r_4$ . Using these, one finds, for example, that  $n_2 = r_2 < r_3 = r_4$  and  $r_3 = r_4 < r_5 = r_5 = r_5$ . And  $r_4 = r_5 = r_5 = r_5 = r_5$ . The two column constants are which is true

Factor A	Factor	8 (sex)	Row	Row means	
(roce)	Girls (1)	Boys (2)	sums		
Negro (1)	11.5, 15.0	182,11 3 142,15 9	$\sum_{j=1}^{2} \sum_{k=1}^{n_{1j}} X_{1jk} = 86.1$	14.4	
White (2)	13.1, 10 4	14 3, 15 3 11.8, 11 0 10 9, 10.5	$\sum_{j=1}^{2} \sum_{k=1}^{n_{2j}} X_{2jk}$ = 109.2	12.1	
Oriental (3)	101, 69	12 8, 8 4 10.6, 10 0	$\sum_{j=1}^{2} \sum_{k=1}^{n_{3j}} X_{3jk} = 58.8$	9.8	
Column sums	$ \begin{array}{c c}     & n_{i1} \\     & \sum_{i=1}^{3} \sum_{k=1}^{N} X_{i1k} \\     & = 78.9 \end{array} $	$\sum_{i=1}^{3} \sum_{k=1}^{n_{i2}} X_{i2k}$ =  75.2	$ \sum_{j=1}^{3} \sum_{j=1}^{2} \sum_{k=1}^{n_{ij}} X_{ijk} \\ = 254 \text{ f} $		
Column	113	12.5		Grand mear	

FIG. 17.18 Illustrative data for the proportional-cell-frequency design of Fig. 17.17.

of the  $n_{ij}$  observations in the ijth cell is denoted by

$$\sum_{i=1}^{n_{ij}} X_{ijk}.$$

The six cell sums for the data in Fig. 17.18 are shown in Fig. 17.19.

The only remaining quantity required for the calculation of the mean squares in the two-factor ANOVA is the sum of the squares of the 21 observations in Fig. 17.18:

$$\sum_{i=1}^{3} \sum_{j=1}^{2} \sum_{k=1}^{n(i)} X_{ijk}^{2} = 11.5^{2} + 15.0^{2} + \dots + 10.0^{2} = 3216.27.$$

The computational formulas for the four sums of squares bear a close resemblance to those for the equal n's case. The four computational formulas appear in Table 17.5.

TABLE 17.5 COMPUTATIONAL FORMULAS FOR SUMS OF SQUARES IN THE TWO-FACTOR ANOVAWITH PROPORTIONAL CELL FREQUENCIES

$$SS_{a} = \sum_{i=1}^{J} \frac{\int_{i=1}^{T} \sum_{k=1}^{n_{i}} X_{in}^{k}}{n_{i}} - \frac{(\sum \sum X_{in})^{2}}{N}$$

$$SS_{a} = \sum_{i=1}^{J} \frac{(\sum_{k=1}^{T} \sum_{k=1}^{n_{i}} X_{in})}{n_{i}} - \frac{(\sum \sum X_{in})^{2}}{N}$$

$$SS_{a,b} = \sum_{i=1}^{J} \sum_{k=1}^{T} \frac{\sum_{i=1}^{n_{i}} X_{in}^{k}}{n_{ij}} - SS_{a} - SS_{a} - \frac{(\sum \sum X_{in})^{2}}{N}$$

$$SS_{a} = \sum_{i=1}^{J} \sum_{k=1}^{T} X_{in}^{n} - \sum_{k=1}^{J} \sum_{k=1}^{T} \frac{\sum_{k=1}^{T} X_{in}^{k}}{n_{ij}}$$

The use of the formulas in Table 17.5 will now be illustrated with the data in Figure 17.18.

$$SS_A = \frac{(86.1)^3}{6} + \frac{(109.2)^2}{9} + \frac{(58.8)^3}{6} - \frac{(254.1)^3}{21}$$

$$= 3136.735 - 3074.610 = 62.125.$$

$$SS_B = \frac{(78.9)^3}{7} + \frac{(175.2)^2}{14} - \frac{(254.1)^3}{12} = 3081.819 - 3074.610 = 7.209.$$

$$SS_{AB} = \frac{(26.5)^3}{2} + \frac{(59.6)^3}{4} + \frac{(35.4)^3}{3} + \frac{(73.8)^3}{6} + \frac{(17.0)^2}{2}$$

$$+\frac{(41.8)^2}{4} - 62.125 - 7.209 - \frac{(254.1)^2}{21}$$

$$= 3145.935 - 62.125 - 7.209 - 3074.610 = 1.991.$$

$$SS_w = 3216.270 - \left[\frac{(26.5)^2}{2} + \dots + \frac{(41.8)^4}{4}\right]$$

$$= 3216.270 - 3145.935 = 70.335.$$

The degrees of freedom and expected values of the four mean squares for the two-factor ANOVA with unequal n's are given in Table 17.6. The

TABLE 17.6 DEGREES OF FREEDOM AND EXPECTED VALUES OF MEAN SQUARES IN THE TWO-FACTOR ANOVA WITH UNEQUAL n's

Source of variation	df	E(MS)
Factor A	<i>I</i> – 1	$\sigma_s^2 + \frac{\sum_{i=1}^I n_i, \alpha_i^2}{I-1}$
Factor B	J – 1	$\sigma_s^2 + \frac{\sum_{j=1}^J n_{,j} \beta_j^2}{J-1}$
$A \times B$	(I-1)(J-1)	$\sigma_*^2 + \frac{\sum_{i=1}^{J} \sum_{j=1}^{J} n_{ij} \alpha \beta_{ij}^2}{(I-1)(J-1)}$
Within cells	N-IJ	σ**

obtained values of the four mean squares are as follows:

$$MS_A = \frac{62.125}{2} = 31.062,$$
  $MS_B = \frac{7.209}{1} = 7.209,$   
 $MS_{AB} = \frac{1.991}{2} = 0.996,$   $MS_w = \frac{70.335}{15} = 4.689.$ 

The entries in Table 17.6 possess only minor differences from their counterparts in the equal n's ANOVA. With the unequal n's ANOVA, we write the degrees of freedom within cells as N-II instead of II(n-1); notice, however, that II(n-1)=IIn-II=N-II.

The E(MS)'s differ from those in the equal n's case only in that each squared effect—main or interaction—is differentially weighted by the number of observations available to estimate it in the layout of data.  $n_{ij}$  is

the general case, of which

$$n_1 = n_2 = \ldots = n_I$$
,  $n_A = n_A = \ldots = n_{AJ}$ ,  $n_{11} = n_{12} = \ldots = n_{IJ}$ 

(i.e., equal column frequencies, equal row frequencies, or equal frequencies for each of the H factor-level combinations) are special cases that permit the "weights"  $(n_i, n_j, o \ n_{ij})$  to be moved to the left of one or both of the summation signs. Note, for example, that in Table 17.6 if the  $n_i$ 's are all equal, say, to c, the E(MS) for factor A becomes

$$\sigma_i^2 + \frac{c\sum_{i=1}^I \alpha_i^2}{I-1}.$$

Inspection of the E(MS)'s in Table 17.6 indicates appropriate F-tests of the three null hypotheses:

$$H_0: \sum_{i}^{I} \alpha_i^2 = 0, \quad H_0: \sum_{j}^{J} \beta_j^2 = 0, \quad H_0: \sum_{i}^{I} \sum_{j}^{J} \alpha_i \beta_{ij}^2 = 0.$$

(Notice that if  $\sum_{i=1}^{J} \alpha_{i}^{2} = 0$ , then  $\sum_{i=1}^{J} n_{i} \alpha_{i}^{2}$  is also equal to zero since if the first condition holds then all  $I \alpha_{i}$ 's are zero. The same is true for the  $\beta_{i}$ 's and  $\alpha \beta_{i} \beta_{i}$ 's.)

If  $H_a$ :  $\sum_{i=1}^{L} \alpha_i^a = 0$  is true, then  $MS_A = SS_A|(I-1)$  will be an unbiased estimator of  $\sigma_i^a$ . Regardless of whether  $H_a$  is true or false,  $MS_a = SS_A|(N-I)$  is an unbiased estimator of  $\sigma_i^a$  and is independent of  $MS_A$ . Thus  $F_A = MS_A|MS_a$  will follow the F-distribution with I - 1 and N - II diffusion over complete replications of the two-factor design that lies generally above the F-distribution with I - 1 and N - II dies above  $\frac{1}{1-1} \sum_{i=1}^{L} H_i$  can be rejected at the  $\alpha$ -level of significance; otherwise,  $H_a$  cannot be rejected. For example, for the data in Fig. 17.18  $F_A = 10.05(14.689 = 6.62$ . If  $\alpha$  had been set at .01, the critical value against which  $F_A$  would be compared is  $\alpha_i F_{i,1,3} = 6.36$ . Thus the data constitute evidence for the rejection of  $H_i$ :  $\nabla_a e^2 = 0$ .

Leaving  $\alpha$  at .01, the following conclusions will be drawn concerning the main effect of factor B and the interaction of A and B:

The critical value for  $F_B$  is  $_{BF_B 13} = 8.68$ . The value of  $F_B$  is 7.209/4.689 = 1.54. Thus the data do not support rejection of  $H_{a^*} \sum \beta_{ij}^2 = 0$ . There is insufficient evidence to support the conclusion that boys and girls of this age differ in long-jumping ability.

The critical value for  $F_{AB}$  is  $_{3}F_{2.15} = 6.36$ . The value of  $F_{AB}$  is 0.996/of no interaction of race and sex.

The results may be tabulated as in Table 17.7.

TABLE 17.7 RESULTS OF THE TWO-FACTOR ANOVA OF THE DATA IN FIGURE 17.12

Source of variation	df	MS	F	P
Race (A)	2	31.062	6.62	p < 0 01
Sex (B)	1	7.209	1.54	p > 0.25
Race $\times$ sex $(A \times B)$	2	0.996	0.21	p > 0.50
Within race-sex groups	15	4.689		•

#### Disproportional Cell Frequencies

The numbers of observations in the cells of the two-factor layout shown in Fig. 17.20 are disproportional. Notice, for example, that  $n_{11}$  is not equal to  $n_1, n_{,1}/n_{,..}$ ;

$$5 \neq \frac{12 \cdot 25}{68} = 4.4.$$

Though it is by no means readily apparent, proportional cell frequencies can be achieved in the layout of Fig. 17.20 by the simple expedient of randomly discarding one of the five scores in the cell in row 1 and column 1 and one of the five scores in the cell in row 2 and column 3. After so doing, the cell frequencies are proportional (see Fig. 17.21). (You may check the cell frequencies of Fig. 17.21 against Eq. (17.14) to satisfy yourself that they are actually proportional.)

When proportional cell frequencies can be achieved from disproportional frequencies by randomly discarding only a few observations from the total layout, then by all means one should do so. Avoiding the computational

Foctor B (f)

1 2 3

1 
$$n_1 = 5$$
  $n_1 = 5$   $n_1 = 2$   $n_1 = 1$ 

2  $n_2 = 8$   $n_2 = 10$   $n_2 = 5$   $n_3 = 3$ 

3  $n_3 = 12$   $n_3 = 15$   $n_3 = 6$   $n_3 = 3$ 

3  $n_4 = 25$   $n_3 = 30$   $n_3 = 13$   $n_4 = 6$ 

FIG. 17.20

Factor B(i) 2 3  $n_{12} = 5$  $n_{22} = 10$  $n_{31} = 12$ 

FIG. 17.21

labor necessary for the analysis of disproportional designs is worth the trivial reduction in power resulting from discarding 5%, say, of the data. When proportional cell frequencies cannot be easily or inexpensively achieved (in terms of power of the F-test) by randomly discarding data from various cells, the analysis that will now be presented is recommended.

The unweighted means analysis is probably the simplest and one of the most justifiable techniques for analyzing disproportional designs. First we shall outline briefly the rationale for the unweighted means analysis; then this technique will be illustrated.

Suppose that data are gathered in a two-factor design with disproportional cell frequencies. Next, consider replacing the  $n_{tt}$  observations in any one cell by a single observation  $X_{tt}$ , the mean of those  $n_{tt}$  scores. Now we have a new layout of data: II cells in a two-factor design with one observation per cell, the mean of the original  $n_{tt}$  scores in the cell. Surely this new layout of data has proportional cell frequencies; indeed, the n's are all equal to 1. It is possible to calculate  $MS'_{tt}$ ,  $MS'_{tt}$ , and  $MS'_{tt}$ , on this new layout of data. (The primes indicate that these are not the MS's based on the original individual observations.) However, there exist no degrees of freedom within cells when n = 1; hence, we cannot calculate a  $MS'_{tt}$  from the two-factor array of cell means

We can find a useful analogue of  $MS'_m$ . Any  $MS'_m$  must estimate the variance of the normal populations from which the observation (or observations) in any one of the U cells is sampled. Since we assume that the  $n_U$  observations in the ijth cell come from a normal population with variance  $\sigma^2_m$ , the variance of  $X_{tt}$ , is  $\sigma^2_m n_t$ , by a well-known theorem. We can see from  $\sigma^2_t / n_{tt}$  that the population variances of the U cell means are heterogeneous (since  $n_U$  differs from cell to cell). However, when  $n^2$  are equal (as they are in the case of the layout of U cell means—n = 1), it has been shown that the T-tests operate as they would if all of the variances were equal and equalled the average of the T-test operate as the denominator of any T-tailor, T-tailor,

average variance of 
$$\bar{X}_{ij} = \frac{\sum\limits_{i=j}^{I} \sum\limits_{n_{ij}} \frac{\sigma_e^2}{n_{ij}}}{\sum\limits_{n_{ij}} \sum\limits_{n_{ij}} \sigma_{ij}}$$

which equals

$$\sigma_{\bullet}^{2} = \sum_{i=1}^{I} \sum_{j=1}^{J} \frac{1}{n_{ij}}.$$

We can estimate  $\sigma_n^2$  the variance of the populations of the original  $n_H$  observations. It is estimated by  $MS_m$ , the mean square within for the original

layout of N individual observations. Hence, we take

$$MS'_{w} = MS_{w} \frac{\sum_{i=1}^{J} \frac{J}{n_{ij}}}{IJ}.$$
 (17.15)

Denote the multiplier of  $MS_{\varphi}$  in Eq. (17.15) by c; c is the average of the reciprocals of the numbers of observations in the II cells.  $MS'_{\varphi}$  is appropriate as a denominator for testing  $MS'_{A}$ ,  $MS'_{B}$ , and  $MS'_{AB}$ .

The unweighted means analysis is illustrated on data in Fig. 17.22. These hypothetical data can be thought of as having been gathered in an experiment to test the effects on intelligence of a chemical that enhances the ability of brain tissue to absorb oxygen; interest centers both on the overall effectiveness of the drug and its possible interaction with diagnosis of the child (normal vs. brain damaged). Intelligence is measured with the Stanford-Binet Intelligence Test.

The data in the top half of Fig. 17.22 are transformed into the layout of cell means in the bottom half of the figure. [Note that (92 + 114 + 107)/3 = 104.33.] The layout of data in the bottom half of Fig. 17.22 is now regarded

### 1. Original Data Evidencing Disproportional Cell Frequencies.

• • •	O.19		
	Foo	tor B - Glut	amic acıd dosage
	1 No Dosage	2 Half Dosage	3 Full Dosoge
Normal 1	92,114, 107 n <sub>II</sub> =3	101,118, 98, 96, 105 n <sub>12</sub> =5	89, 120, 110, 115 n <sub>13</sub> =4
Factor A-Diognosis State A-Social Brain S	91, 74, 65, 90 n <sub>21</sub> =4	101, 68, 59 n <sub>22</sub> =3	79, 88, 55, 67, 93 n <sub>23</sub> = 5

2. Layout of Cell Means.

1	Factor B 2	3	Row totals	Row means
104 33	103.60	108.50	316.43	105.48
80.00	76.00	76.40	232,40	77.47
184.33 92.17	179.60 89.80	184.90 92.45	548.83	
	80.00 184.33	1 2 104 33 103.60 80.00 76.00 184.33 179.60	1 2 3 104 33 103.60 108.50 80.00 76.00 76.40 184.33 179.60 184.90 92.17 89.80 92.45	1 2 3 Row totals 104 33 103.60 108.50 316.43 80.00 76.00 76.40 232.40 184.33 179.60 184.90 548.83

FIG. 17.22 Layout of data for the unweighted means analysis.

as a two-factor ANOVA with n = 1 observation per cell. The sums of squares are calculated as follows:

$$SS'_{A} = \sum_{j=1}^{2} \frac{\left(\sum_{i=1}^{3} X_{ij}\right)^{2}}{3} - \frac{\left(\sum_{i=1}^{3} \sum_{i=1}^{2} X_{ij}\right)^{2}}{6} = \frac{(316.43)^{2} + (232.40)^{2}}{3} - \frac{(548.83)^{2}}{6}$$
$$= 51,379.235 - 50,202.395 = 1176.840.$$

$$SS'_B = \sum_{t=1}^{3} \frac{\left(\sum_{t=1}^{2} X_t\right)^2}{2} - \frac{\left(\sum_{t=1}^{3} \sum_{t=1}^{3} X_{tt}\right)^2}{6}$$

$$= \frac{(184.33)^2 + (179.60)^2 + (184.90)^2}{2} - \frac{(548.83)^2}{6}$$

$$= 50,210.860 - 50,202.395 = 8.465.$$

$$SS'_{AB} = \sum_{i=1}^{3} \sum_{j=1}^{3} X_{ij}^{3} - SS'_{A} - SS'_{B} - \frac{\left(\sum_{i=1}^{3} \sum_{j=1}^{3} X_{ij}\right)^{3}}{6}$$

$$= (104.33)^{2} + (103.60)^{2} + \dots + (76.40)^{2}$$

$$- 1176.840 - 8.465 - \frac{(548.83)^{2}}{6}$$

$$= 51,402.919 - 1176.840 - 8.465 - 50,202.395 = 15.219.$$

MS' is obtained by calculating MS from the 24 scores in the top half of Fig. 17.22 and multiplying by c. SS, is given by

$$SS_{w} = \sum_{i}^{I} \sum_{j}^{S_{si}} x_{IR}^{2} - \sum_{i}^{I} \sum_{j}^{\frac{S_{si}}{2}} \frac{\left(\sum_{i}^{S_{si}} X_{IR}\right)^{2}}{n_{ti}}$$

$$= 92^{2} + 114^{2} + 107^{2} + \dots + 93^{2} - \left[\frac{313^{2}}{3} + \dots + \frac{382^{2}}{5}\right]$$

$$= 209,061 - 205,522,933 = 3538.067.$$
The value of

The value of c is given by

$$c = \frac{\sum_{i=1}^{J} \frac{1}{II}}{II} = \frac{\frac{1}{3} + \frac{1}{4} + \frac{1}{3} + \frac{1}{4} + \frac{1}{3}}{6} = 0.2611.$$

Now we are prepared to calculate mean squares and F-ratios:

$$MS'_A = \frac{SS'_A}{1} = \frac{1176.840}{1} = 1176.840.$$

$$MS'_B = \frac{SS'_B}{2} = \frac{8.465}{2} = 4.233.$$

$$MS'_{AB} = \frac{SS'_{AB}}{2} = \frac{15.219}{2} = 7.610.$$

$$MS'_{\Psi} = c\frac{SS_{\Psi}}{2} = 0.2611 \frac{3538.067}{18} = 0.2611(196.559) = 51.322$$

The F-ratios for testing the three null hypotheses are:

$$F_A = \frac{MS'_A}{MS'_w} = \frac{1176.840}{51.322} = 22.93.$$
  $F_B = \frac{MS'_B}{MS'_w} = \frac{4.233}{51.322} = 0.08.$   $F_{AB} = \frac{MS'_AB}{MS'} = \frac{7.610}{51.322} = 0.15.$ 

Quite obviously there exists no evidence to conclude a significant main effect for factor B—glutamic acid dosage or interaction effect of A and B. The F-ratio for factor A is 22.93, which exceeds  $_{.00}F_{1.18} = 8.28$ . The probability p of obtaining an  $F_A$  as large as 22.93 or larger if the null hypothesis for factor A is true is less than .01. Hence,  $H_0: \sum_{i=1}^{2} \alpha_i^2 = 0$  is rejected.

The results of the unweighted means analysis can be presented as is usual for a two-factor ANOVA:

Source of variation	df	MS	F	р	
Diagnosis (A) Glutamic acid	1 2	1176.840 4.233	22.93 0.08	p < .01 p > .50	
dosage (B) $A \times B$ Within cells	2 18	7.610 51.322	0.15	p > .50	

#### 17.15 MULTIPLE COMPARISONS IN THE TWO-FACTOR ANOVA

The rejection of a null hypothesis about the main effect of factor A implies only that at least two of the I population means differ. All I population means may differ, only two may differ, or any number in between may

differ; the F-test does not distinguish among these possibilities. As was true with the one-factor ANOVA, multiple-comparisons procedures are required to determine which of the pairs of I sample means show differences large enough to permit the conclusion that the underlying population means differ. The above remarks apply to factor B as well. The Tukey and Scheffe methods of multiple comparisons were presented in Chapter 16. In this section, we shall indicate how each method would typically be employed in the two-factor case.

When the number of observations upon which each of the I sample means for factor A is based is equal for all I means and when one is interested only in the I(I-1)/2 differences between I population means, the Tukey method should be used. The set of I(I-1)/2 simultaneous confidence intervals on the  $\bar{\mu}_L - \bar{\mu}_L$ ; with confidence coefficient  $1 - \alpha$  is constructed as follows:


$$(\bar{X}_{l.} - \bar{X}_{l.}) \pm {}_{l-e}q_{I,N-IJ} \sqrt{\frac{MS_w}{N|I}},$$
 (17.16)

where  $_{1-\alpha}q_{I,N-IJ}$  is the  $100(1-\alpha)$  percentile point in the Studentized range distribution with I and N-II degrees of freedom (see Table F in Appendix A).

MS. is the mean square within from the two-factor ANOVA, and N/I is the common number of observations on which each of the I sample means is based. (Recall that this number of observations must be equal for each group to use the Tukey method.) When the I is in the II cells are equal, N/I = n/I.

Construction of a set of simultaneous confidence intervals by the Tukey method will be illustrated on the data in Fig. 17.23 for which the ratio  $F_B$  proved to be significant at the  $\alpha = .01$  level.

Notice that N = 27, I = 2, and J = 3; furthermore, the number of observations upon which each column mean is based equals N|J = 27/3 = 9.



The value of  $MS_B$  is 232.78; hence,  $F_B = 232.78/24.62 = 9.45$ , which is significant at the 01 level with 2 and 21 degrees of freedom. The questions remain whether group 1 differs significantly from both groups 2 and 3 or whether groups 2 and 3 are significantly different.

A set of simultaneous confidence intervals may be constructed around the three differences between pairs of means by the method outlined in Table 17.7.

TABLE 17.7 CONSTRUCTION OF A SET OF SIMULTANEOUS CONFIDENCE INTERVALS AROUND THE DIFFERENCES BETWEEN PAIRS OF MEANS BY THE TUKEY METHOD (J = 3,  $\alpha = .01$ )

Differences between means	$. \mathfrak{pq}_{2.11} \sqrt{\frac{MS_{w}}{N/J}}$	Confidence interval [Eq. (17.16)]
$X_{.1.} - X_{.1.} = 9.44$	$(4.61) \sqrt{\frac{24.62}{9}} = 7.62$	(1.82, 17.06)
$X_{1} - X_{2} = 8.00$	7.62	(0.38, 15.62)
$X_{.1.} - X_{.1.} = 8.00$ $X_{.1.} - X_{.1.} = -1.44$	7.62	(-9.06, 6.18)

The confidence intervals on  $\bar{\mu}_1$  —  $\bar{\mu}_2$  and  $\bar{\mu}_1$  —  $\bar{\mu}_3$  do not include zero; here, we conclude a significant difference between  $\bar{X}_1$ , and  $\bar{X}_2$ , and between  $\bar{X}_1$ , and  $\bar{X}_3$ . The confidence interval on  $\bar{\mu}_2$  and  $\bar{\mu}_3$  includes zero; there exists no evidence that  $\bar{\mu}_2$  and  $\bar{\mu}_3$  differ. The confidence afforded by the Tukey method is 99% that all three of these conclusions are simultaneously correct.

The above techniques apply to factor A as well, of course, the only alterations being in the use of  $_{1-q}q_{I,N-IJ}$  and N/I as a divisor of  $MS_w$ .

When one desires to perform multiple comparisons on the means of one factor in a two-factor design in which these row or column means are based on differing numbers of observations, the Scheffe method should be employed. The Scheffe method was discussed at length in Chapter 16. The only alterations to Eq. (16.5) occasioned by the fact that a two-factor ANOVA is being analyzed are that  $MS_w$  now stands for "within cells mean square" and  $n_1, \ldots, n_T$  now become  $n_1, \ldots, n_T$ , i.e., the n's are the total numbers of observations upon which the column (or row) means are based. The formula for constructing a set of simultaneous confidence intervals around contrasts of the row means by the Scheffe method is as follows:

$$(c_1 \vec{X}_{1..} + ... + c_I \vec{X}_{L.}) \pm \sqrt{MS_{\sigma} \left(\frac{c_1^2}{n_1} + ... + \frac{c_I^2}{n_I}\right)} \sqrt{(I-1)_{I-\sigma} F_{I-1..N-IJ}}.$$
(17.17)

#### PROBLEMS AND EXERCISES

1. Complete the ANOVA table and make F-tests of the null hypotheses for factors A and B and the interaction of A and B at the  $\alpha = .01$  level of significance.

Source of variation	df	SS	MS
Factor A	4	64.26	
Factor B	5	46.85	
$A \times B$		40.05	
Within	120	1136.53	
Total	149	2411.69	

2. The figure below represents cell, row, and column population means in a two-factor ANOVA design.

a. Is 
$$H_0$$
:  $\sum_{i=1}^{I} \alpha_i^2 = 0$  true or false?

b. Is 
$$H_0$$
:  $\sum_{j=1}^{J} \beta_j^2 = 0$  true or false?

c. Is 
$$H_0$$
:  $\sum_{i=1}^{J} \sum_{j=1}^{J} \alpha \beta_{ij}^{\alpha} = 0$  true or false?

- 3. You have been told that research shows no interaction between "sex" and "method of foreign language instruction." The main effect of the aural-oral method of instruction is +6 points on a French language mastery test; the main effect for females is +2 points on the same mastery test. In what direction and by how many points should the mean of females studying French by the aural-oral method deviate from the grand mean (the mean for both sexes
  - 4. Graph the interaction of factor B, traditional orthography vs. Initial Teaching Alphabet, and factor A, sex, from the following cell means on the dependent variable, reading performance measured in grade-placement units:

Boys 
$$\vec{X}_{1l} = 4.61$$
  $\vec{X}_{12} = 4.48$   $\vec{X}_{21} = 5.12$   $\vec{X}_{22} = 4.96$ 

5. For each of the following arrangements of data, graph the interaction of factor B, sex, with factor A, "treatments." Place factor B on the abscissa so that each treatment level yields one line in the graph. Which cases show ordinal interaction (all questions of statistical significance aside)? Which show disordinal interaction?

Treatments 2 
$$\overline{X}_{21}$$
=25.60  $\overline{X}_{12}$ =21.66  $\overline{X}_{12}$ =21.67  $\overline{X}_{21}$ =21.68  $\overline{X}_{21}$ =21.48  $\overline{X}_{31}$ =16.44  $\overline{X}_{32}$ =17.89 (b)  $\overline{X}_{11}$ =19.63  $\overline{X}_{12}$ =14.81  $\overline{X}_{11}$ =19.63  $\overline{X}_{12}$ =14.81  $\overline{X}_{12}$ =13.55 (c)  $\overline{X}_{22}$ =13.55  $\overline{X}_{21}$ =10.01  $\overline{X}_{22}$ =13.55  $\overline{X}_{21}$ =10.06  $\overline{X}_{22}$ =13.55  $\overline{X}_{21}$ =10.06  $\overline{X}_{22}$ =13.55

6. Graph the interaction of "sex" and "treatments" for the data under (b) of Prob. 5. Place "treatments" on the abscissa. Is the interaction ordinal or disordinal? Was the interaction ordinal or disordinal when "sex" was placed on the abscissa of the graph? 7. A researcher is studying the effects on learning of inserting questions into instructional materials. There is some doubt whether these questions would be more effective preceding or following the passage about which the question is posed. In addition, the researcher wonders if the effect of the position of the questions is the same for factual questions and for questions that require the learner to compose a thoughful and original response. A group of 24 students is split at random into four groups of six students each. One group is assigned to each of the four combinations of factor B, "position of question (before vs. after the passage)" and factor A "type of question (factual vs. thought-provoking)." After 10 hours of studying under these conditions, the 24 students are given a 50-tiem test over the content of the instructional materials. The following test scores are obtained:

Position of question

	Be	fore	After	
	19	23	31	28
§ Fact	29	26	26	27
Types of question	30	17	35	32
٥ د	27	21	36	29
E Thought	20	26	39	31
•	15	24	41	35

Perform a two-factor ANOVA on the above data. Test the null hypotheses for both main effects and the interaction effects at the .10 level of significance.

 Analyze the following scores on a 50-item vocabulary test administered to 24 students of high and average intelligence (factor A) after one year of studying a foreign language under one of three methods (factor B):

Factor B

			Aural-oral method	Translation method	Combined methods
			36	26	19
		High (IIS end	29	23	30
	e	apove)	25	21	28
actor A	Intelligence	31	31	18	20
	T te		33	20	17
		Average (115 and	verage 19 22	22	13
		below)	37	14	23
			28	15	18

F

- Perform F-tests of the null hypotheses for rows, columns, and interaction at the .05 level.
   These same data were analyzed under Prob. 8 at the end of Chapter 15.
   However, there the data were created as a conference ANOVA throughout the conference and the same conference and the sa
- b. In these same data were analyzed under Prob. 8 at the end of Chapter 15. However, there the data were regarded as a one-factor ANOVA, "method of instruction" being the only factor. Verify that SS<sub>B</sub> and MS<sub>B</sub> are the same whether the data are considered in a one-factor or two-factor design. Also verify that SS<sub>B</sub> for the one-factor ANOVA of the data in Prob. 8 of Chapter 15 equals SS<sub>A</sub> + SS<sub>AB</sub> + SS<sub>w</sub> for the two-factor ANOVA of the same data.
- 9. D. L. Williams (1968) experimented with rewriting sixth-grade science materials so that the reading difficulty was appropriate for the third grade. (The following data are based on his study.) A group of 240 sixth-grade pupils were randomly assigned to one or the other of two levels of reading difficulty: "Grade 6 reading difficulty" or "Grade 3 reading difficulty." For three days, pupils read "Resources of the Sea," a chapter in a science text, at either the third-grade or sixth-grade reading difficulty level. A 129-item multiple-choice test was administered at the end of the experiment to assess comprehension. Within both levels of reading difficulty, pupils were classified as either high, average, or low scorers on the reading section of the Stanford Achievement Test. The following data were obtained for the two levels of reading difficulty and three levels of reading achievement:

	Reading difficulty			
	Grade 6	Grade 3		
	n=40	n=40		
_ High	$\bar{X}_{11} = 89.93$	$\bar{X}_{12} = 93.89$		
Average	s11=12.02	s <sub>12</sub> = 13.02		
e e	n=40	n = 40		
E Average	$\bar{X}_{2i} = 70.79$	$\bar{X}_{22} = 72.55$		
	s <sub>21</sub> =14.76	s <sub>22</sub> =15.90		
Reading	n ≈ 40	n=40		
E Low	$X_{31} = 52.09$	X <sub>32.</sub> ≈56.84		
	s <sub>31</sub> =11.30	s <sub>32</sub> =12.21		

Perform a two-factor ANOVA on these data. Test all three null hypotheses at the .05 level of significance. (Hint:  $MS_w$  is the average of the six within-cell nariances.)

10. All of the first-grade pupils in a school were given a reading readiness test in September. According to the test manual, a total of 12 girls and 18 boys had scores so low that "formal reading instruction should be postponed and "reading readiness activities" should be substituted." A reading researcher randomly assigned the 12 girls and 18 boys in equal numbers to one of the following three conditions: (1) give 18 weeks of reading readiness activities before beginning instruction; (2) give 9 weeks of readiness activities; (3) commence reading

(a)

instruction immediately (0 weeks of readiness activities). A reading achievement test was administered to all 30 punils at the end of the third grade. The following grade-placement scores were obtained:

Reading readiness activities

		18 weeks		9 weeks		None	
	(	34	4.0	3 7	4.3	4,1	4.4
Sex	Bays	4.1	3.8	4.2	3.3	3.7	3.2
		3.9	4.4	3.8	3.1	3 4	4.4 3.2 4 0
	Girts	4.0	38	46	4.2	44	4.3 4.6
	Girss	43	4.7	3.9	3.6	4.0	4.6

Perform E-tests at the .05 level of the null hypotheses of no difference among amounts of readiness activities, no differences between the sexes, and no interaction between the two factors

11. In which of the following designs are the cell frequencies proportional, i.e., in which instances are  $n_{ij} = (n_i n_j)/n$ ?

12. From which one cell could two observations be discarded to achieve proportional cell frequencies:

n <sub>II</sub> = 10	A <sub>12</sub> = 15	n <sub>3</sub> =10
n <sub>21</sub> = 6	n <sub>22</sub> =9	n <sub>23</sub> ≃6
n <sub>31</sub> = 2	n32=3	n33=4

13. A sample of 40 tenth-grade pupils drawn randomly from either public or parochial schools (factor B) can be classified as of either high, middle, or low socio-economic status (factor A). The pupils earned the following grade-placement scores on a mathematics achievement test:

Type of echool

			i ype i	JI SUIDOI	
		Put	blic	Pare	ochiał
		9 5	10 1	10 4	8.5
	High	8 7	10 4	11 6	8.5 10.3 10.2 11 1 10 3 10 6 10.7 9 9
				9.3	10.2
Ş		8.4	11.4	10 4	11 1
ş	Middle	10 5	10 6	9.4	8.5 10.3 10.2 11 1 10 3 10 6 10.7
Ĕ	міадіе	98	10.4	10 6	10 6
Sacioeconomic status		10 6		11.0	10.7
		8.6	8.9	10.0	99
		7.3	97	9 5	10.6
	Low	10 2	10.0	8 9	104
		9 5	7 1		ł
	,	98	J		)

Because the cell frequencies are disproportional, perform an unweighted-means analysis of variance to test the null hypotheses for rows, columns, and interaction.

## 18

## ONE-FACTOR AND MULTI-FACTOR ANALYSIS OF VARIANCE: RANDOM, MIXED, AND FIXED EFFECTS

#### IB.I INTRODUCTION

This chapter has three objectives: (1) to introduce an alternative to the analysis of variance model that underlies Chapters 15 through 17; (2) to show a combination of this new model and the previous ANOVA model that is of great importance in experimental research; (3) to present rules for generating the fundamental elements (sums of squares, degrees of freedom, expected values of mean squares, etc.) of most analyses of variance one is apt to meet in practice. These three topics will be dealt with in the order indicated.

#### 18.2 THE RANDOM-EFFECTS ANOVA MODEL

In this section we shall introduce an analysis of variance model and applications of the model that are fundamentally different from those of the fixedeffects ANOVA model of Chapter 15. Fortunately, many similarities between the techniques of Chapter 15 and those to be presented here should substantially facilitate learning the material in this chapter. The model which we shall now develop is called the *random-effects* analysis of variance model.

The reasons for having spoken of a "fixed-effects" ANOVA model in Chapter 15 were never made clear. It is simpler to comprehend the meaning of "fixed effects" when they can be contrasted with "random effects." In the ANOVA model of Chapter 15, we were primarily interested in making statistical inferences about the set of main effects  $\alpha_1,\ldots,\alpha_J$ . The inference to be made was from a set of sample data U groups of n persons) to J populations. We conceived of an infinite sequence of replications of the experiment in each of which there were n persons under each of J levels. Our interest in the J population means—or equivalently the J main effects—led us to consider only replications of the experiment in which the same J treatment levels appeared. In a sense, then, we "fixed" (as opposed to allowing to vary) the J populations so that each one—and only these ones—yielded one sample of size n in each replication of the experiment.

Now we wish to allow the populations that give rise to the samples in each replication of the experiment to vary across replications; this time we shall sample populations A, B, and C, and next time populations D, E, and F will be sampled. This procedure would be adopted if our interest were in the variance of the means of a large collection of populations only a few of which could be observed in one experiment. We might, then, randomly select for a given experiment both a sample of J populations from a large collection of populations and then samples of size n observations from each population. If we ran the experiment again, we would not "fix" the factor levels so that the same J populations were again represented; rather we would randomly sample a different set of J populations. As there were "effects" of the form  $\alpha_j = \mu_j - \mu$  in the fixed-effects model, so there are effects of the form  $a_i = \mu_i - \mu$  for this new model. However, whereas formerly the complete set of a,'s was present in every replication of the fixed-effects analysis, now we want to consider the case of having only a random sample of the a, effects present in a replication of the experiment. Hence, the name randomeffects ANOVA model. Figure 18.1 is an attempt to capture graphically this distinction between the fixed-effects and random-effects models.

As part of its testing program, a school system administers the Metropolitan Achievement Tests to the 7th, 8th, and 9th grades of all junior high schools in the system each May. Often these data are used to make comparisons. Is school A showing higher performance than school B? Has school C improved since last May? The administrators realize that not all sepects of the testing are standardized. In particular, the tests are given at aspects of the day in different schools. If there is some substantial different times of the day in different schools If there is some substantial variability in test performance associated with the time of day at administration of the test, then a comparison of schools A and B might be telling more something about all the  $\mu_i$ 's. The variance of the  $\mu_i$ 's tells us what sort of differences to expect in the  $\mu_i$ 's, the mean test scores for the population of students across all 360 testing periods.

We shall now formalize the random-effects model. Suppose we have a virtually infinite population of  $\mu_i$ 's. (Again, we are not worried, because in our example we have only 360  $\mu_i$ 's.) Denote the average of all these  $\mu_i$ 's by  $\mu$ . The ith student earns a score of  $X_{ij}$  during the jth testing period. The difference between the ith student's score at period j and the mean of all students' scores in that period is denoted by ei; hence,

$$X_{ij} = \mu_j + e_{ij}, \tag{18.1}$$

where  $X_{ij}$  is the score for student i in level j of the factor "time of day,"

 $\mu_i$  is the mean of all students' scores for the jth level of the

 $e_{tt}$  is the deviation of the ith student's score from the mean of all students' scores at the jth level of the factor.

The model in Eq. (18.1) will be altered in form slightly by deviating the  $\mu_i$ 's around  $\mu_i$ , the mean of all the  $\mu_i$ 's, as follows:

$$X_{ij} = \mu + (\mu_i - \mu) + e_{ij}. \tag{18.2}$$

The model in Eq. (18.2) follows by adding in and subtracting out  $\mu$ from Eq. (18.1). A notation simplification of Eq. (18.2) will put the randomeffects ANOVA model into its customary form:

$$X_{ij} = \mu + a_i + e_{ij}, (18.3)$$

where

$$a_i = \mu_i - \mu$$
 and  $e_{ij} = X_{ij} - \mu_j$ .

Suppose that  $\mu$ , the mean test score of all students over all 360 testing periods, is 30; and suppose that all students average 4 points above 30 when they take the test during the 36th testing period (9:35-10:25 a.m.); and finally suppose that the fourth student scores 8 points below the average of all students tested at this period. If the model in Eq. (18.3) holds, what test score will the fourth student obtain during the 36th testing period?

$$X_{4,36} = \mu + a_i + e_{ij} = 30 + 4 - 8 = 26.$$

Initially, we were interested in the variance of the  $\mu_i$ 's. How has this concern changed now that the  $\mu$ ,'s have been "eliminated" from the model? Since  $a_j$  is simply  $\mu_j$  minus a constant,  $\mu_j$ , the variance of the  $\mu_j$ 's is the same as the variance of the  $a_i$ 's. Thus, differences in test scores due to the time of as the variance of the fest will be reflected in the variance of the  $a_j$ 's, day of administration of the test will be reflected in the variance of the  $a_j$ 's, i.e., in  $\sigma^2$ .

Before we can proceed to estimate  $\sigma_a^2$  and make statistical inferential

statements about it, it will be necessary to make some assumptions about the model in Eq. (18.3):

- The random-effects a<sub>j</sub> are normally distributed with a mean of zero and a variance of σ<sup>a</sup><sub>o</sub>, i.e., a ~ N(0, σ<sup>a</sup><sub>o</sub>).
- 2. The error component  $e_{ij}$  is normally distributed with mean zero and variance  $\sigma^{z}$ , i.e.,  $e \sim N(0, \sigma^{z})$ .
- 3. The random-effects  $a_i$  and the error components  $e_{ii}$  are independent, from which it follows that  $\rho_{es}=0$ . Furthermore, the  $e_{ii}$  are all independent of one another.

These assumptions have several implications for the testing study. If the random-effects model is an adequate description of the data in the testing study, then:

- 1. The means  $(\mu, s)$  of populations of students' scores for all possible testing times during the day should appear to be normally distributed around  $\mu$  and have a variance of  $\sigma_s^2$ . Since  $a_j = \mu_j \mu$ , the  $a_j$  should have a normal distribution around zero with a variance of  $\sigma_s^2$ .
- 2. The test scores X<sub>ii</sub> for all students for a single time period (the fith time period) should be normally distributed around a mean of μ<sub>i</sub> with variance σ<sup>i</sup>. Since q<sub>i</sub> = X<sub>ii</sub> − μ<sub>i</sub>, σ<sub>i</sub> will be normally distributed around zero with variance σ<sup>i</sup>. This is assumed to be true for all values of f<sub>i</sub>. This assumption is the counterpart of the homogeneous variances assumption in the fixed-effects model.
- 3. If samples of n test scores were taken from the populations of scores for a randomly designated J time periods, the value of a, would give no information about whether ê, was above or below zero and similarly, the value of ê, would give no information about whether a different ê, n was above or below zero.

After these preliminaries, we come at last to the techniques for actually estimating both of and of from data. The problem to be solved is one of test, of, and the variability in test scores due to the time of administering the test, of, and the variance in test scores due to differences among students taking the test at the same time, of.

Data are gathered in a two-phased sampling plan. First, J levels of the random-effects factor are drawn at random. For example, J = 5 time periods from the population of 360 periods are randomly chosen. Just as random sampling of persons from populations permits generalizations to the tion to a population of levels. Second, a random sample of n observations is drawn from the populations in each of the J levels. In our example, this

procedure is simulated by administering the Metropolitan Reading Test at each of the five time periods to n=7 students randomly drawn from the population of all students. Restricting the ns to be equal is not a matter of concenience here; the methods of this section have never been fully developed for the unequal n's case.

In our study, we have sampled J=5 a's and Jn=35 e's. The scores on the 44-item reading test may be arranged as in Table 18.1.

TABLE 18.1 DATA FROM A STUDY OF THE EFFECTS OF TESTING TIME ON TEST SCORES

(11:09-11:59 a m.)	(10:14-11.04 a m.)	Testing time period 3 (1:31-2.21 p.m.)	(9:50-10 40 a m)	(1:53-2·43 p m.)
$X_{11} = 35$ $X_{12} = 32$ $X_{23} = 41$ $X_{41} = 42$ $X_{41} = 31$ $X_{41} = 36$ $X_{71} = 33$ $X_{71} = 36.00$ $X_{71}^{2} = 17.33$	$X_{12} = 36$ $X_{23} = 32$ $X_{21} = 34$ $X_{22} = 40$ $X_{23} = 37$ $X_{23} = 44$ $X_{23} = 42$ $X_{23} = 37$ $X_{34} = 48$ $X_{25} = 88$	$\begin{array}{c} \chi_{13} = 28 \\ \chi_{13} = 27 \\ \chi_{33} = 23 \\ \chi_{43} = 15 \\ \chi_{45} = 15 \\ \chi_{45} = 29 \\ \chi_{45} = 28 \\ \chi_{73} = 33 \\ \chi_{73} = 32.81 \end{array}$	$\begin{array}{c} X_{14} = 27 \\ X_{24} = 16 \\ X_{34} = 40 \\ X_{44} = 32 \\ X_{45} = 36 \\ X_{47} = 38 \\ X_{77} = 22 \\ X_{74} = 30.14 \\ x_{8}^{2} = 78.81 \end{array}$	$X_{15} = 32$ $X_{15} = 34$ $X_{25} = 26$ $X_{45} = 28$ $X_{45} = 23$ $X_{45} = 36$ $X_{75} = 28$ $X_{75} = 29$ $S_{5} = 21.29$

It should come as no great surprise that the variance of the J sample means,  $\overline{X}_1, \ldots, \overline{X}_J$ , bears on the question of how large the variance of the  $\mu_i$ 's is. Nor should it be unexpected that the variance of the  $X_{ij}$ 's within each sample yields an estimate of  $\sigma^2$ .

As in the fixed-effects ANOVA model, each of the J groups of scores produces a sample variance  $s_i^2$  that is an estimator of  $\sigma^2$ , the variance of the population of scores for the jth factor level. The average of these J sample variances is the best estimator available for  $\sigma^2$ . For the fixed-effects model, the average within-sample variance was called  $MS_w$ , the mean-square within:

$$MS_{w} = \frac{s_{1}^{2} + \ldots + s_{J}^{2}}{J} = \frac{\sum_{j=1}^{J} \sum_{i=1}^{n} (X_{ij} - \bar{X}_{,i})^{2}}{J(n-1)}.$$
 (18.4)

 $MS_{\omega}$  is an estimator of  $\sigma^2$  in the following senses:

- The expected value of MS<sub>w</sub> is equal to σ², i.e., E(MS<sub>w</sub>) = σ².
   Thus, MS<sub>w</sub> is an unbiased estimator of σ², the variance of scores within each factor level.
- Over replications of the study with J factor levels and n observations within each level, the sampling distribution of MS<sub>w</sub> is given by:

$$MS_w \sim \sigma^2 \frac{\chi_{J(n-1)}^2}{J(n-1)}$$
,

i.e., the sampling distribution of  $MS_n$  is  $\sigma^2$  times a chi-square distribution with J(n-1) degrees of freedom divided by J(n-1).

It is simple to prove that the expected value of  $MS_{\omega}$  is  $\sigma^2$  if one remembers that the expected value of  $\chi^2_{\omega}$  is simply n:

$$E(MS_n) = E\left[\frac{\sigma^2 \chi_{J(n-1)}^2}{J(n-1)}\right] = \frac{\sigma^2}{J(n-1)}[J(n-1)] = \sigma^2.$$

As in Chapter 15, the definitional formula for  $MS_{\pi}$  in Eq. (18.4) is not convenient for computations. In its place, one should use the following formula to compute  $MS_{\pi}$ :

$$MS_{\bullet} = \frac{\sum_{i=1}^{J} \sum_{i=1}^{N} X_{ii}^{*} - \sum_{j=1}^{J} \left( \sum_{i=1}^{N} X_{ij} \right)^{*}}{J(n-1)}.$$
 (18.5)

The application of Eq. (18.5) to the data in Table 18.1 will be illustrated in Table 18.2.

TABLE 18.2 ILLUSTRATION OF CALCULATION OF MS, AND MS, ON THE DATA IN TABLE 18.1

$$\frac{1}{2} \frac{7 \text{ Fitting time period}}{3} \frac{3}{4} \frac{4}{5}$$

$$\frac{7}{5} X_{11} = 252 \sum_{i=1}^{7} X_{i1} = 265 \sum_{i=1}^{7} X_{i1} = 183 \sum_{i=1}^{7} X_{i1} = 211 \sum_{i=1}^{7} X_{i1} = 207$$

$$\frac{7}{5} X_{11}^{2} = 9176 \sum_{i=1}^{7} X_{11}^{2} = 10.145 \sum_{i=1}^{7} X_{11}^{2} = 4981 \sum_{i=1}^{7} X_{11}^{2} = 6833 \sum_{i=1}^{7} X_{11}^{2} = 6249$$

$$\sum_{i=1}^{5} \sum_{i=1}^{7} X_{i1} = 1118 \sum_{i=1}^{5} \sum_{i=1}^{7} X_{i1}^{2} = 37,384$$

$$55_{1} = \frac{(232)^{3} + (265)^{3} + (183)^{3} + (211)^{3} + (211)^{3} + (211)^{3}}{35} = \frac{254,588}{7} - \frac{1,249,924}{35}$$

$$145_{2} = \frac{55_{1}}{17,1} - \frac{697,40}{40} = 164,49$$

$$55_{2} = \frac{35_{1}}{17,384} - \frac{(252)^{3} + (265)^{3} + (181)^{3} + (211)^{3} + (211)^{3} + (207)^{3}}{17,384} = 37,384 - 36,349.71$$

$$1014,27 = \frac{55_{2}}{10} - \frac{1014,27}{10} = 31.81$$

459

The function  $MS_b$ , mean-square between groups, with which you became familiar in Chapter 15 was there seen to be closely related to the variance of J sample means. Having defined  $MS_b$  before, we can use it now in estimating  $\sigma_{ab}^2$  the variance of the population of a's or, what is equivalent, the variance of the population of  $\mu_c$ 's. You will recall that

$$MS_b = \frac{n \sum_{j=1}^{J} (\vec{X}_{,j} - \vec{X}_{,.})^2}{J - 1},$$
 (18.6)

i.e.,  $MS_b$  is n times the sample variance of the J sample means,  $\bar{X}_1, \dots, \bar{X}_J$ .

Again, the definitional formula for a mean square makes for clumsy calculation. A more efficient computational formula for  $MS_b$  is:

$$MS_b = \frac{\sum_{i=1}^{J} \frac{\left(\sum_{t=1}^{n} X_{it}\right)^2}{n} - \frac{\left(\sum_{t=1}^{J} \sum_{t=1}^{n} X_{tt}\right)^2}{Jn}}{J-1}.$$
 (18.7)

It can be shown—though we shall not do so here—that the expected value of  $MS_b$  has the following form:

$$E(MS_b) = \sigma^2 + n\sigma_a^2, \qquad (18.8)$$

i.e., on the average—the average across an infinite collection of independent replications of the same study with I randomly chosen levels with I randomly chosen observations at each level— $MS_b$  will equal  $\sigma^2$ , the same variance that is estimated by  $MS_w$ , plus I into the variance of the population of I, so (which is the same as the variance of the population of I, so

Thus we see that  $MS_b$  estimates all that  $MS_w$  estimates and more. The something "more" is n times the quantity of interest,  $\sigma_a^2$ .

It happens that the sampling distribution of MSb is given by the following:

$$MS_b \sim (\sigma^2 + n\sigma_a^2) \frac{\chi^2_{J-1}}{J-1}$$
, (18.9)

i.e., over repeated samplings of  $J \times n$  layouts of data in which both factor levels and observations within levels are randomly sampled, the sampling distribution of  $MS_0$  is that of chi-square with J-1 degrees of freedom multiplied by the constant  $\sigma^2 + n\sigma_0^2$  over J-1. Unlike the fixed-effects model,  $MS_0$  has a sampling distribution that is a constant times the chi-square distribution even when there are differences among the  $\mu_1$ 's.

In Table 18.2 the calculation of  $MS_o$  and  $MS_o$  is illustrated using the data in Table 18.1. We see in Table 18.2 that the best estimate we have of  $\sigma^2$ , the variance of Metropolitan Reading Test scores in the population of junior-high-school students, is  $MS_o$ , which equals 33.81. We shall denote this estimate by  $\sigma^2$ , the """ indicating an estimate of the parameter below it.

The mean of all five groups is approximately 32; we would then expect that Metropolitan Reading Test scores for junior-high-school students would be normally distributed around a mean of 32 with a standard deviation of approximately √33.81 = 5.82.

The value of  $MS_b$  is an estimate of  $\sigma^2 + n\sigma_{\sigma}^2$ . We can obtain an estimate

of of alone from MS, and MS, in the following manner:

$$E\left[\frac{MS_b - MS_w}{n}\right] = \frac{E(MS_b) - E(MS_w)}{n} = \frac{\sigma^2 + n\sigma_n^2 - \sigma^2}{n} = \frac{n\sigma_n^2}{n} = \sigma_n^2.$$
(18.10)

The best estimate we can attain of  $\sigma_s^*$  is given by  $(MS_b - MS_b)/n$ . This function estimates  $\sigma_s^*$  unbiasedly. If  $MS_b$  were larger than  $MS_b$ —which could happen—the estimate  $\sigma_s^*$  of  $\sigma_s^*$  would be negative. We know, of course, that  $\sigma_s^*$  is positive, or at the very least, zero. Thus, if ever a negative estimate of  $\sigma_s^*$  were obtained, it would automatically be set equal to zero instead. For the data in Table 18.1, we have

$$\hat{\sigma}_a^2 = \frac{MS_b - MS_m}{n} = \frac{164.40 - 33.81}{7} = 18.66.$$

Thus the best estimate,  $\hat{\sigma}_{a}^{*}$ , of the variance of the means of populations of test scores obtained at each of the 360 possible testing periods during the day is 18.66. The standard deviation of these population means is approximately 4.32.

This estimate of  $\sigma_a$  has some meaning in and of itself. For example, if schools A and B chose to administer the Metropolitan Reading Test at times 1 and 2, respectively, and if  $a_1=+4$  and  $a_2=-4$  (not an unusual occurrence since each time-period effect lies only one standard deviation from zero, the mean of all  $a_i$ 's), then the two schools would be expected to differ by 8 points on the test even if they were achieving at the same level. An obtained difference between them of 8 points might easily be due to the fact that they administered the test at different times of the day. Note that this would be highly improbable if  $\sigma_a$  were 25, for example; it would be highly unlikely that an 8 point difference between school means could be attributed to "time of day" effects if  $\sigma_a$  were .25. An estimated variance component of 18.66 in our example indicates that for test scores to be comparable from school to school in the system, an effort must be made by the administration to see that the tests are administered at the same time of day in each school.

While this "absolute" interpretation of a is occasionally possible, more often a only takes on meaning when compared with a. For example, we might find it more informative to know that the ratio of a it of its j or 3 or .005. In our example, did is 0.55. In the parlance popular among statisticians, it is said that "the variance due to 'time of day' is about half as great as the variance among students' test scores."

One can identify at least three inferential questions in the one-way random-effects model: (1) How can a confidence interval on  $\sigma_0^2$  be established around  $\sigma_0^2$ ? (2) How can a confidence interval on  $\sigma_0^2/\sigma^2$  be established around  $\sigma_0^2/\sigma^2$ ? (3) How can the hypothesis that  $\sigma_0^2 = 0$  be tested?

Unfortunately, the techniques for setting a confidence interval around  $\hat{\sigma}_a^2$  on  $\sigma_a^2$  cannot be derived in a straightforward manner from the original model. Approximate techniques are available; however, they are quite complex (see Scheffé, 1959, pp. 231-35).

It is a relatively simple matter to establish a confidence interval on  $\sigma_a^2/\sigma^2$  around  $\hat{\sigma}_a^2/\hat{\sigma}^2$ . The  $1 - \alpha$  confidence interval on  $\sigma_a^2/\sigma^2$  is given in Eq. (18.11);

$$\operatorname{prob}\left\{\frac{1}{n}\left[\frac{MS_{b}}{MS_{w}}\left(\frac{1}{1-(a/2)F_{J-1,J(n-1)}}-1\right)\right] < \frac{\sigma_{x}^{2}}{\sigma^{2}}\right\}$$

$$<\frac{1}{n}\left[\frac{MS_{b}}{MS_{w}}\left(\frac{1}{1-\sigma^{2}F_{J-1,J(n-1)}}-1\right)\right]\right\} = 1 - \alpha. \quad (18.11)$$

Both the  $100[1 - (\alpha/2)]$  and the  $100(\alpha/2)$  percentile points in  $F_{J-1,J(n-1)}$  are required in Eq. (18,11). Recall that

$$_{\alpha/2}F_{J\sim 1,\,J(n-1)}=\frac{1}{_{1-(\alpha/2)}F_{J(n-1),\,J-1}}.$$

Suppose we wish to construct the 95% confidence interval on  $\sigma_a^{3/o^2}$  for the data in Table 18.1. From Table E in Appendix A we first find the value of  $_{.97}F_{4.30} = 3.25$ . Next the value of  $_{.97}F_{30.4}$  is found from which the other required percentile is calculated as follows:

$$_{.025}F_{4,30} = \frac{1}{_{.025}F_{30,4}} = \frac{1}{8,46} = 0.12.$$

Substituting the two percentiles along with the values of  $MS_b$ ,  $MS_w$ , and n into Eq. (18.11) yields the 95% confidence interval on  $\sigma_a^2/\sigma^2$ :

$$\frac{1}{7} \left( \frac{164.40}{33.81} \frac{1}{3.25} - 1 \right); \qquad \frac{1}{7} \left( \frac{164.40}{33.81} \frac{1}{0.12} - 1 \right) = (.97, 5.64).$$

The 95% confidence interval on  $\sigma_s^a/\sigma^2$  extends from .07 to 5.64. This interval tells how very uninformative our study has been. According to our data, we have little reason not to believe that  $\sigma_s^2$  could be anywhere from one-tenth as large as  $\sigma^a$  to about 5½ times larger than  $\sigma^a$ . A study that produces stable estimates of  $\sigma_s^a$  and  $\sigma_s^a$ , i.e., short confidence intervals on  $\sigma_s^a/\sigma^a$ , cannot be done cheaply; both J and n must be fairly large.

Testing the null hypothesis  $H_0: \sigma_0^2 = 0$  is not of as much interest as was the test of  $H_0: \sum_{i=0}^{n} \sigma_i^2 = 0$  in the fixed model. We can often become sufficiently skeptical to entertain the possibility that a small set of means are equal—as in the fixed-effects model ANOVA—in which case we wish to test the hy-

pothesis of no differences. Certainly far less frequently does it strike us as even remotely possible that an infinite number of levels of a factor all have the same population mean, as must be true if  $H_0$ :  $\sigma_a^2 = 0$  is true. Hence, when the random-effects ANOVA model is applied, interest will center on the estimation of  $\sigma_a^2$  more often than on testing whether  $\sigma_a^2 = 0$ .

However, if compelling reasons exist for testing  $H_0$ :  $\sigma_a^2 = 0$ , it can be tested readily as follows: If  $F = MS_b/MS_w$  exceeds the  $100(1 - \alpha)$  percentile point of the F-distribution with J-1 and J(n-1) degrees of freedom,  $H_0$ :  $\sigma_a^2 = 0$  can be rejected at the  $\alpha$ -level of significance. For example, with the data in Table 18.1,  $F = MS_b/MS_w = 164.40/33.81 = 4.86$ , which exceeds 4.02, the 99th percentile in  $F_{4,30}$ . Thus,  $H_0$ :  $\sigma_a^2 = 0$  can be rejected with  $\alpha = .01$ .

For all values of  $\sigma_a^2$ , the sampling distribution of  $F = MS_b/MS_w$  is given by:

$$F = \frac{MS_b}{MS_w} \sim \left(1 + \frac{\hbar \sigma_\sigma^2}{\sigma^2}\right) F_{J-1, J(n-1)}.$$
 (18.12)

Several major points developed in this section about the one-factor random-effects ANOVA model are summarized in Table 18.3.

TABLE 18.3

3.3	SUMMARY Source of	OF THE C	ONE-FACTOR RANDOM-EFFEC	FOR RANDOM-EFFECTS ANOVA				
variation		df	ss	E(MS)	Estimated variances			
1	Between levels	J-1	$\sum_{i=1}^{J} \frac{\left(\sum_{i=1}^{n} X_{ii}\right)^{t}}{n} - \frac{\left(\sum_{j=1}^{J} \sum_{i=1}^{n} X_{ij}\right)^{t}}{J_{n}}$	$\sigma^z + n\sigma_z^z$	$\hat{\sigma}_{a}^{1} = \frac{MS_{b} - MS_{a}}{n}$			
,	Within levels	J(n - 1)	$\sum_{j=1}^{J} \sum_{i=1}^{n} X_{ij}^{t} - \sum_{j=1}^{J} \frac{\left(\sum_{i=1}^{n} X_{ij}\right)^{t}}{n}$	a <sup>‡</sup>	$\hat{\sigma}^z = MS_w$			
					_			

A final word is in order on the question of the consequences of violating the assumptions of the random-effects ANOVA model. In Chapter 15 we saw that the consequences for the validity of the fixed-effects ANOVA of violation of the normality assumption were negligible; in addition, heterogeneous variances are immaterial in the fixed-effects model if all n's are equal. However, the situation is different for the random-effects ANOVA model. In particular, if the kurtosis of the random-effects  $a_j=\mu_j-\mu$  deviates from the kurtosis of the normal distribution, the validity of confidence intervals on  $\sigma_a^{ij}\sigma^{i}$  or of the test of  $H_a$ :  $\sigma_a^{i}=0$  may be open to serious question. Nonnormality of the observations within the factor levels of the random-effects

## 18.3 THE MIXED-EFFECTS ANOVA MODEL

The third and final distinctly different analysis of variance model that will be dealt with is a coalition of the fixed-effects and the random-effects models. This union of the two models into a mixed-effects model is particularly useful in experimental research.

As the name "mixed-effects model" might suggest, the mixed model involves two sets of effects: one fixed and the other random. Naturally, then, the model describes data gathered in a two-factor design, similar in appearance only to the two-factor fixed model of Chapter 17. One factor, e.g., the row factor, comprises a set of I fixed effects; the column factor is a random sample of J random effects from a supposedly infinite population of normally distributed effects. Consider a hypothetical experiment in which evaluators are comparing three elementary science curricula: (1) the American Association for the Advancement of Science curriculum (AAAS): (2) the Elementary Science Study curriculum (ESS); (3) the Science Curriculum Improvement Study (SCIS) curriculum. The dependent variable of the experiment is "knowledge of the processes of scientific inquiry," which is measured by a 75-item objective test. The experiment is designed as follows: 10 elementary schools are chosen to participate; six experimental classrooms are available within each school; by random methods, two classrooms are assigned to each curriculum within each school. Observations of the dependent variable are taken by administering the "processes of science" test to all 60 experimental classrooms and averaging the scores of the students within each class; thus, the 60 observations in the experiment are classroom means. The data from the experiment may be laid out as in Table 18.4.

TABLE 18.4 LAYOUT OF DATA FROM AN EXPERIMENT IN WHICH THREE ELEMENTARY SCIENCE CURRICULA ARE COMPARED (SCORES ARE CLASS MEANS TO THE NEAREST WHOLE AUBINARE ON A 37-JEM TEST).

		MENINES		OLL 11	01100				/				
School (j)								Row	Row				
		A	В	c	D	E	F	G	H	1		sums	means
	AAAS	31,27	21, 26			35, 28						603	30.15
Curriculum	ESS	39, 46	32, 30			42, 47					30, 27	747	37,35
(f)	SCIS	35, 28	31, 25	42, 39	36,38	41,37	38, 38	27,25	29,31	45, 41	31,26	683	34 15
Column	ums	206	165	258	193	230	227	150	192	248	164	2033	

The general observation in Table 18.4 is denoted by  $X_{ik}$ , where i ranges over rows (curricula) from 1 to 3, j ranges over columns (schools) from 1 to 10, and k ranges over observations within cells (classrooms) from 1 to 2. In

general, i = 1, ..., I; j = 1, ..., J; and k = 1, ..., n. The notation is equivalent to that for the two-factor fixed-effects ANOVA of Chapter 17.

The two-factor design in Table 18.4 presents two sets of main effects plus one set of interaction effects. The two main effects are "curricula," which will be called factor A, and "schools," factor B. Clearly the three science curricula were not sampled from a large population of such curricula, nor are the evaluators interested in generalizing to a hypothetical population of other curricula from which they could have conceivably been sampled. Interest focuses on the question of which one of these curricula is superior to the other two. Hence, the three main effects of factor A are considered "fixed." On the other hand, the 10 schools represented in the experiment can be considered to have been sampled from a population of schools; or more importantly, the evaluators do not want the results of their study to be limited to the 10 schools of the experiment. Their conclusion about the relative superiority of the curricula must be generalized beyond the 10 experimental schools to have any utility. Hence, the 10 schools are regarded as randomly sampled levels of the random-effects factor B. We have, then, a fixed-effects and a random-effects factor crossed in the same design. The structural model postulated for the observations in this design is aptly named the mixed-

$$X_{ijk} = \mu + \alpha_i + b_j + \alpha b_{ij} + \epsilon_{ijk}, \qquad (18.13)$$

where  $X_{ijk}$  is the kth observation in the ijth cell,

is the grand population mean of all observations,

 $\alpha_i$  is the main effect  $(\mu_i - \mu)$  of the ith level of the fixed factor,

 $b_i$  is the main effect  $(\mu_i - \mu)$  of the jth level of the random factor,  $\alpha b_{ii}$  is the interaction effect  $(\mu_{ii} - \mu_i - \mu_j + \mu)$  of the ijth com-

bination of the fixed and random factor, and  $e_{ijk}$  is the error, or "residual," component that accounts for variation

of observations within the ijth cell.

The following restrictions (not assumptions) are placed on the terms of the mixed-effects model in Eq. (18.13):

$$1. \alpha_1 + \cdots + \alpha_I = 0.$$

2. The population mean of the infinite number of b,'s—only 10 of which are present in the science curriculum experiment-is zero.

3.  $\alpha b_{1i} + \alpha b_{2j} + \ldots + \alpha b_{1i} = 0$  for all j.

4. The population of the infinite set of  $ab_{ij}$ 's for a single i (row) has a

The above restrictions imply that if we sum across the rows of the data layout in Table 18.4 the fixed-effects and interaction effects will "add out," i.e., sum to zero; however, summing across the columns of the design to obtain a particular row mean, for example, will not cause the J values of  $b_i$  or the J values of  $\alpha b_{ii}$  to sum to zero.

Suppose we wish to compare  $\vec{X}_{1...}$  and  $\vec{X}_{2...}$ , i.e., the means of the class-rooms under the AAAS and ESS curricula. These two means have the following structure in terms of the model in Eq. (18.13):

$$\begin{split} \tilde{X}_{1..} &= \frac{1}{JK} \bigg[ \sum_{1}^{J} \sum_{1}^{K} \left( \mu + \alpha_1 + b_J + \alpha b_{1J} + e_{1Jk} \right) \bigg] \\ &= \mu + \alpha_1 + \tilde{b}_. + \overline{\alpha b_1}_. + \tilde{\epsilon}_{1..} \, . \\ \tilde{X}_{2..} &= \mu + \alpha_2 + \tilde{b}_. + \overline{\alpha b_2}_. + \tilde{\epsilon}_{2..} \, . \end{split}$$

The difference between  $\vec{X}_{1..}$  and  $\vec{X}_{2..}$  is

$$\overline{X}_{1} - \overline{X}_{2} = (\alpha_{1} - \alpha_{2}) + (\overline{\alpha b}_{1} - \overline{\alpha b}_{2}) + (\overline{e}_{1} - \overline{e}_{2}).$$

Because the ab's do not sum to zero across the J columns and because a replication of the experiment with a different set of J random effects would produce different values of  $ab_1$ , and  $ab_2$ , the sampling variance of the difference between  $X_1, \dots X_L$ , will contain a component for the interaction effects, ab. This fact will be fully appreciated when we discuss the expected values of mean squares for the mixed model. But before entering upon that subject, we must state the assumptions which must be made about the mixed model of Eq. (18.13).

The following assumptions are made about the terms of the model

$$X_{iii} = \mu + \alpha_i + b_i + \alpha b_{ii} + e_{iii}$$

- The random effects, b<sub>j</sub> = μ<sub>j</sub> μ, are normally distributed with a mean of zero and a variance of σ<sup>2</sup><sub>s</sub>.
- The interaction effects αb<sub>ij</sub> are normally distributed over j for each i with a mean of zero and a variance of σ<sup>2</sup><sub>ab</sub>.
- The error components e<sub>ijk</sub> are distributed normally and independently
  of the b's and αb's with mean zero and variance σ².

A fourth very important assumption is necessary to insure the validity of the hypothesis test of the fixed main effects:

4. For all pairs of levels of the fixed factor, the correlation (across the population of random effects) of the scores under one level with the scores under the other level of the pair must be the same. For example, if in the population of all schools the correlation of classroom mean test scores for the AAAS and ESS curricula is 5,0 then the population correlation for AAAS and SCIS and for ESS and SCIS must be .50 as well.

We shall indicate later how the methods of this section should be modified when assumption 4, homogeneous correlations, is suspected to be seriously violated.

With the model and its assumptions stated, we can proceed to develop the methods by which null hypotheses about the fixed and random main effects and the interaction effects may be tested. Our objective, then, is to test the following three pairs of hypotheses:

1. 
$$H_6: \sum_{i=1}^{I} \alpha_i^2 = 0$$
 vs.  $H_1: \sum_{i=1}^{I} \alpha_i^2 \neq 0$ .

2. 
$$H_0$$
:  $\sigma_b^2 = 0$  vs.  $H_1$ :  $\sigma_b^2 \neq 0$ .

3. 
$$H_0$$
:  $\sigma_{ab}^2 = 0$  vs.  $H_1$ :  $\sigma_{ab}^2 \neq 0$ .

Not surprisingly, perhaps, the road to the hypothesis tests leads through the familiar sums of squares, degrees of freedom, mean squares, and expected values of mean squares. In fact, the computations of SS, df, and MS in the mixed model are identical to the calculations in the two-factor fixed model. The two models do not part company until the expected values of the mean squares are reached. The computations in the two-factor mixed model are presented in Table 18.5 and are illustrated on the data of Table 18.4 in Table 18.6.

TABLE 18.5 COMPUTATIONAL FORMULAS FOR SUMS OF SQUARES, DEGREES OF FREEDOM, AND MEAN SQUARES IN THE TWO-FACTOR MIXED MODEL

Source of			
pariation	df	\$\$	MS
Among rows (fixed factor A)	I-1	$SS_A = \sum_{i}^{I} \frac{\left(\sum\limits_{j}^{J} \sum\limits_{k}^{n} X_{ijk}\right)^{k}}{J_{n}} - \frac{\left(\sum\limits_{j}^{J} \sum\limits_{j}^{J} \sum\limits_{k}^{n} X_{ijk}\right)^{k}}{IJ_{n}}$	<u>ss₄</u> 11
Among columns (random factor 8)	J 1	$SS_B = \sum_{j}^{J} \frac{\left(\sum_{k}^{J} \sum_{k}^{n} X_{iik}\right)^{k}}{j_{fi}} - \frac{\left(\sum_{k}^{J} \sum_{j}^{J} \sum_{k}^{n} X_{iik}\right)^{k}}{i f n}$	SS.
Interaction of A and B	(I - I)(I - I)	$SS_{AB} = \sum_{i}^{3} \sum_{j}^{3} \frac{\left(\sum_{k}^{n} X_{ijk}\right)^{2}}{n} - SS_{A} - SS_{B}$	$\frac{SS_{AB}}{(I-1)(I-1)}$
		$-\frac{\left(\sum\limits_{i}^{f}\sum\limits_{j}^{J}\sum\limits_{k}^{n}\chi_{ijk}\right)^{k}}{II_{R}}$	, ,,,
Within cells	IJ(n - 1)	$SS_{o} = \sum_{i}^{T} \sum_{j}^{J} \sum_{k}^{n} \chi_{ijk}^{2} - \sum_{i}^{T} \sum_{j}^{J} \frac{\left(\sum_{k}^{n} \chi_{ijk}\right)^{2}}{n}$	SS <sub>*</sub>

#### TABLE 18.6

ILLUSTRATION OF CALCULATION OF SUMS OF SQUARES AND MEAN SQUARES ON THE DATA IN TABLE 184

1. 
$$SS_A = \frac{(603)^2 + (747)^2 + (683)^2}{20} - \frac{(2033)^3}{60} = 520.53$$
 $MS_A = \frac{SS_A}{J-1} = \frac{520.53}{2} = 260.27$ 

2.  $SS_B = \frac{(206)^2 + (165)^3 + \dots + (164)^3 - (2033)^2}{6} = 2059.68$ 
 $MS_B = \frac{SS_B}{J-1} = \frac{2059.68}{9} = 228.85$ 

3.  $SS_{AB} = \frac{(58)^2 + (47)^2 + \dots + (57)^2}{2} - 520.53 - 2059.68 - \frac{(2033)^2}{60} = 268.47$ 
 $A(S_{AB} = \frac{SS_{AB}}{(J-1)(J-1)} = \frac{268.47}{18} = 14.92$ 

4.  $SS_B = (31)^3 + (27)^2 + \dots + (26)^3 - \frac{(58)^3 + (47)^2 + \dots + (57)^3}{2} = 71.971$ 
 $-71.733.50 = 237.50$ 
 $MS_B = \frac{273.50}{J(G-1)} = \frac{273.50}{30} = 7.92$ 

Once again it is by way of the expected values of mean squares that we can see how various ratios of mean squares bear on the question of whether or not a null hypothesis is true. The expected values—long-run average values over replications of the experiment in Table 18.4 with different schools and classrooms each time, for example—of the mean squares in Table 18.5 are given in Table 18.7.

E(MS)

TABLE 18.7 EXPECTED VALUES OF MEAN SQUARES IN THE TWO-FACTOR MIXED-EFFECTS
MODEL

Mean sauare

	$nJ\sum_{i=1}^{J}\alpha_{i}^{2}$
MSA (fixed factor)	$\sigma^2 + n\sigma_{ab}^2 + \frac{1-1}{I-1}$
MSa (random factor)	$\sigma^2 + nl\sigma^2$
MSss (mixed interaction)	$\sigma^2 + n\sigma_{\sigma^2}^2$
MS.	σ¹

The expected mean squares in Table 18.7 present a far different picture from the E(MS)'s for the two-factor fixed-effects model. Particularly, the

variance of the interaction effects,  $\sigma_{sb}^2$ , that appears in  $E(MS_A)$  looks unusual at first. The presence of  $\sigma_{sb}^2$  in  $E(MS_A)$  was anticipated in our discussion of comparing  $X_1$ , and  $X_{1n}$ , the difference between which contained  $\tilde{\sigma}_{1n}^2$ ,  $-\tilde{\sigma}_{2n}^2$ . [from which we obtain  $\sigma^2$  in  $E(MS_A)$ ] and  $\frac{1}{ab_1}$ ,  $-\frac{1}{ab_2}$ . [from which we obtain  $\sigma_{sb}^2$  in  $E(MS_A)$ ]. No comparable situation existed in the two-factor fixed model; there it was seen that summing across any one row would "add out" the J interaction effects since

$$\sum_{i=1}^{J} \alpha \beta_{ij} = 0.$$

Consider first the problem of testing the null hypothesis that all the  $\alpha_i$ 's are zero, i.e.,

$$H_0: \sum_{i=1}^I \alpha_i^2 = 0.$$

You must repress what is perhaps your first inclination to divide  $MS_A$  by  $MS_w$  and refer the ratio to the F-distribution. Notice that  $MS_A/MS_w$  does not bear directly on the question of whether

$$\sum_{i=1}^{I} \alpha_i^2$$

is zero or not. The quantity  $\sum \alpha_i^*$  could be zero and yet  $MS_A/MS_w$  might be large because  $\sigma_w^*$  is not zero. The difference between  $MS_A$  and  $MS_w$  estimates  $n_A^*$ ;  $+ nJ \sum \alpha_i^*[I-I]$ . In instead of just  $nJ \sum \alpha_i^*[I-I]$ , as it did in the two-factor fixed model. It can be seen by inspecting the above expected mean squares that  $E(MS_A)$  differs from  $E(MS_{AB})$ , on the just of the discrepancy between  $MS_A$  and  $MS_{AB}$ , or the ratio of  $MS_A$  to  $MS_{AB}$ , bears on the size of  $\sum \alpha_i^*$ . More specifically, given the assumptions of the mixed-effects model (particularly assumption 4),  $F = MS_A/MS_{AB}$ , will have the F-distribution with degrees of freedom I - I and I - IJ/J - IJ/J

$$H_0: \sum \alpha_s^2 = 0$$

is true. A positive value of  $\sum a_1^2$  will tend to inflate  $MS_A$  above  $MS_{AB}$  and give values of  $F = MS_A/MS_{AB}$  that are larger than the typical values in the  $F_{f-1,(f-1)(f-1)}$  distribution.

Though the null hypothesis  $H_s: \sigma_s^s = 0$  is of less interest than the hypothesis about the fixed main effects, it can be tested by referring the ratio  $F = MS_B/MS_a$  to the table of the F-distribution with J - 1 and II(n-1) degrees of freedom. Moreover,  $\sigma_s^s$  can be estimated by  $(MS_B - MS_a)[nI)$ , which is unbiased. The  $1 - \alpha$  confidence interval on  $\sigma_s^2[\sigma^s$  can be constructed by using Eq. (18.11) of Sec. 18.2;  $MS_B, nI$ , and  $F_{J-1,II(n-1)}$  are substituted for  $MS_a, n$ , and  $F_{J-1,II(n-1)}$  in that equation.

The null hypothesis  $H_b$ :  $\sigma_{ab}^2 = 0$  may be tested by referring F =

 $MS_{AB}/MS_{\infty}$  to the F-distribution with (I-1)(J-1) and IJ(n-1) degrees of freedom.

For the data in Table 18.4, the three null hypotheses mentioned above have been tested at the .05 level of significance. The results appear in Table 18.8.

TABLE 18.8 NULL HYPOTHESIS TESTS FOR THE DATA IN TABLE 18.4

•••	Null hypothesis	F-ratio	Critical F $(\alpha = .05)$	Decision
	$H_0: \sum_{i=1}^l \alpha_i^t = 0$	$F = \frac{MS_A}{MS_{AB}} = \frac{260.27}{14.92} = 17.44$	$_{.96}F_{2,18}=3.55$	Reject H <sub>0</sub>
	$H_0$ : $\sigma_b^b = 0$	$F = \frac{MS_B}{MS_w} = \frac{228.85}{7.92} = 28.90$	$_{.95}F_{9,30}=2.21$	Reject Hu
	$H_0$ : $\sigma_{ab}^2 = 0$	$F = \frac{MS_{AB}}{MS_w} = \frac{14.92}{7.92} = 1.88$	$_{.95}F_{19,30}=1.95$	Do not reject H <sub>0</sub>

We see in Table 18.8 that the main effects for both the fixed and random factors are quite statistically significant. The test for interaction is significant with an  $\alpha$  of .10 but not at the .05 level. It would now be legitimate to apply the Tukey method of multiple comparisons, which was discussed in Chapter 16, to factor A ("science curricula") to determine which pairs of means differ significantly. In place of  $MS_{\alpha}$  with I(n-1) degrees of freedom that are used in the Tukey method in the one-factor ANOVA, one uses  $MS_{AB}$  with I(n-1)I(J-1) degrees of freedom (Scheffé, 1959, p. 270).

The two-factor mixed-effects ANOVA model with n=1, i.e., one observation per cell, is frequently encountered. For example, six persons (random factor B) may each be observed under four treatment conditions (fixed factor A), as shown in Fig. 18.2. This design is commonly referred to as a regeated measures design, because observations of persons are made several times instead of once. In general, an observation in the repeated measures design is denoted by  $X_{ij}$ , i indicating the row in which an observation lies and i indicating its column. The mean squares between rows (the fixed

FIG. 18.2

		Persons (/)						
		1	2	3	4_	5	_6	
	1	XII	X12	X <sub>13</sub>	X <sub>14</sub>	X <sub>15</sub>	X16	
Treatments	2	X21	X22	X <sub>23</sub>	X24	X <sub>25</sub>	X26	
(/)	3	X31	X32	X <sub>33</sub>	X34	X <sub>35</sub>	X <sub>36</sub>	
	4	X41	X42	X43	X44	X <sub>45</sub>	X46	

factor), between columns (the random factor), and for the interaction of rows and columns can all be calculated by means of the computational formulas in Table 18.5 by letting n=1 and dropping out the index k and all summations over it. Since n=1, there are II(n-1)=0 degrees of freedom for variation within cells; thus, the variance of scores around the mean of the ijth cell cannot be estimated. The analysis of variance table for the sources of variation that do exist appears as Table 18.9.

TABLE 18.9 ANOVA TABLE FOR THE TWO-WAY MIXED-EFFECTS ANOVA WITH n = 1


Source of variation	df	E(M5)		
		J∑ a}		
Factor A (fixed factor)	1-1	$\sigma^2 + \sigma_{s_1}^2 + \frac{1}{I - I}$		
Factor B (random factor) Interaction of A and B	J-1 (I-1)(J-1)	$\sigma^1 + \sigma^1_2$ $\sigma^1 + \sigma^1_{24}$		

If we assume that the I levels of the fixed factor would all correlate equally in the population of persons (the random factor), then  $F=MS_A | MS_A m$  will have an F-distribution with I-1 and (I-1)(I-1) degrees of freedom when  $H_2: \sum_{i=1}^{n} = 0$  is true. Hence,  $F=MS_A | MS_A m$  can be compared with  $_{1-1}E_{1-1,1-1,1-1}U_2$  to test  $H_1$  at the  $_{1}E_{2}$ -level of significance.

No tests of the null hypotheses  $H_0$ :  $\sigma_b^2$  and  $H_0$ :  $\sigma_{ab}^2 = 0$  are possible in the mixed-effects model when n = 1.

An important assumption for the test of  $H_0$ :  $\sum \alpha_i^2 = 0$  to be valid in the mixed model is that the correlations of all pairs of levels of the fixed factor across the population of random factor levels must be the same. Violations of this assumption work to increase the actual probability of a type I error above the value believed to hold, the "nominal" value. When heterogeneous correlations among the pairs of levels of the fixed factor are suspected. special measures must be taken to insure the validity of the F-test of the null hypothesis about the fixed main effects. Box (1954b) showed that the effect of heterogeneous correlations of the fixed factor levels was to produce a sampling distribution of  $F = MS_A | MS_{AB}$  that has degrees of freedom less than I-1 and (I-1)(J-1) when  $H_0$  is true. Greenhouse and Geisser (1959) showed that, at worst, the degrees of freedom could only be reduced to 1 and J-1 for the sampling distribution of  $F=MS_d/MS_{AB}$  under a true null hypothesis. (Also see Lana and Lubin, 1963.) These findings suggested a contingent hypothesis testing procedure that should be used if heterogeneous correlations are suspected:

<sup>1.</sup> If  $F = MS_A/MS_{AB}$  exceeds the  $1 - \alpha$  percentile point in the F-






FIG. 18.3 Schematic diagrams of the analysis of the repeated measures design.

distribution with 1 and J-1 degrees of freedom, reject  $H_0$ :  $\sum \alpha_i^2 = 0$  at the  $\alpha$ -level of significance. (The conservative test.)

- If F = MS<sub>A</sub>|MS<sub>AB</sub> falls below the 1 α percentile point in the F-distribution with I 1 and (I 1)(I 1) degrees of freedom, do not reject H<sub>0</sub>: ∑ α<sup>2</sup><sub>i</sub> = 0 at the α-level of significance. (The apparent test.)
- 3. If  $F = MS_A/MS_{AB}$  falls between  $_{1-}F_{1,J-1}$  and  $_{1-}F_{J-1,(I-1)(J-1)}$ , one must resort to a multivariate technique known as *Hotelling's T<sup>2</sup>* test (see Scheffé, 1959, or Winer, 1962).

The strategy for analyzing a repeated measures design is depicted in Fig. 18.3.

## I8.4 RULES OF THUMB FOR WRITING THE ANOVA TABLE

The following sections of this chapter consist of a collection of "rules of thumb" for finding all entries in an analysis of variance (ANOVA) table for a

		Factor C	Factor T Treatments				_
		Class	75	T <sub>2</sub>	T3	-T4	75
Ţ		Cı	X1, X2	اننا			
(	Pı	Cz	:				
a. S. Tune of		C3					
is type of l	P <sub>2</sub>	C4		<u> </u>		ļ	ļ
_		C <sub>5</sub>		1		<u> </u>	1
		Ce	L	<u></u>	<u> </u>	<u> </u>	لـــــــا

FIG 18.4 Data layout of the example used to illustrate the application of the ANOVA rules of thumb.

large class of ANOVA models.\* For designs in which each pair of factors is completely crossed or nested, in which each factor is random or fixed, and in which the same number of replications (observations) are taken within the smallest subdivision of the design (cell), rules are provided that specify the possible sources of variation, the associated degrees of freedom, computational formulas for sums of squares, and expectations of mean squares. If you master these simple rules you will hopefully come to regard complex analyses of variance as less inconvenient and be encouraged to attempt them when they are appropriate.

Many sources exist that provide rules of thumb for finding some of the entries in an ANOVA table from some of the designs considered here (e.g., see Bennett and Franklin, 1954; Cornfield and Tukey, 1956; Ouenther, 1964; Henderson, 1959; Scheffé, 1959; Schultz, 1955; Winer, 1962). None of them, however, duplicates the coverage of this section, although the somewhat inaccessible Henderson (1959) paper is the most similar to the presentation to follow.

The following experiment involving six classrooms will be used to illustrate the application of the rules presented. Three of six classrooms included in an experiment are from public schools  $(P_1)$  and three from private schools  $(P_2)$ . Each classroom was administered five treatments. Suppose that each class was split into 10 subgroups at random and that two subgroups responded independently under each of the five treatments. Thus, two observations,  $X_1$  and  $X_2$ , on the single dependent variable X exist for each classroom. This design is illustrated in Fig. 18.4.

These sections are revised from Millman and Glass (1967).

#### 18.5 DEFINITIONS OF TERMS

#### I-A. Crossed and Nested Factors

Two factors are crossed if every level (the different categories of a factor are called levels) of one of the factors appears with every level of the other factor. That is, there must be at least one observation for every possible combination of levels of factors that are completely crossed. Thus "type of school" and "treatments" (having two and five levels, respectively) are crossed since there are observations taken at each of the 10 school-treatment combinations.

A factor is said to be nested in a second factor if each level of the first (the nested factor) appears in exactly one level of the second factor. In our experiment, "classroom" is nested. No classroom appears in both a public  $(P_1)$  and a private  $(P_2)$  school. If, however, the same three classrooms were involved in both  $P_1$  and  $P_2$ , then C would not be nested in P. Nesting exists when one level of a variable does not appear with all levels of another variable. Since  $C_1$  under  $P_2$  is not the same class as  $C_4$  under  $P_2$ ,  $C_4$  does not appear in both levels of P. Note that  $C_1$  is combined with all levels of T ("treatments"); the factor "classroom" is not nested in T. For the purposes of this discussion, any nested factor must have the same number of levels in each level of the factor in which it is nested. In our illustration, there are three levels of "classroom" nested under both levels of "type of school."

#### II-B. Random and Fixed Factors

A factor may be considered random if the levels of that factor used in the study are a simple random sample from a population of levels with normally distributed effects, "Students' and "classrooms" are two factors frequently considered random. Results of an ANOVA may be generalized to the population of levels of a random factor. When all levels of a factor are in the study (e.g., male-female or high-average-low), when only the levels of interest to the investigator are in the study (e.g., method A and method B where the other methods are not of interest), or when a systematic selection of levels is used, the factor is considered fixed. Results of an ANOVA may be generalized only to the population of replications of the experiment in which the specific levels of the fixed factor included in the study are present. For example, a study that systematically selected grades three, six, nine, and 12 can generalize its results only to those four grades.

Actually, the status (fixed or random) of a factor depends as much upon the population of replications of a study to which one wishes to generalize as it does upon the way in which the levels were chosen. Five levels of a factor could be randomly sampled from a virtually infinite population of levels, but the factor would be "fixed" if one made inferences to replications of the

study in which only those five levels chosen appear. This is an abstruse point about which we will have no more to say.

In our example, "classrooms" is considered a random factor, and "type of school" and "treatments" are fixed. We shall also call "replication" within the smallest cell of a design a nested factor that is always random and is nested within all the other factors of the design. This point is important and allows us to specify rules of thumb that are simpler than many others proposed.

#### 18.6 DETERMINING THE POSSIBLE LINES (SOURCES OF VARIATION) OF THE ANOVA TABLE

#### II-A. Notation

- 1. The source of variation for a factor that is not nested within any other factors is denoted by a capital letter, e.g., A, B, C, . . . .
- 2. The source of variation for a nested factor is denoted by a capital letter followed by a colon and then the letter or letters denoting the factors within which it is nested, e.g., A:B for factor A nested within
- 3. The source of variation for an interaction is denoted by a combination of letters identifying the interacting factors followed by a colon and the letter or letters of the factors within which the interaction is nested, e.g., AB or AB: C.

#### 11-B. Rules

- 1. The ANOVA table has one line for each factor both crossed and nested (this includes the factor "replications").
- 2. The ANOVA table has one line for all possible (two factor, three factor, etc.) interactions among factors. To determine which interactions can exist, all possible pairs, trios, etc. of factors are formed by the following rules (if there is no nesting other than "replications," there will be  $2^k - k - 1$  such interactions, where k
  - a. In the symbol denoting the interaction, write to the left of the colon the letters to the left of the colons in the factors being combined. (If no colon appears in the notation for a factor, it is understood to be at the right of all letters.) b. Write following the colon, but with no repetition of a letter,

those letters to the right of the colons in the factors being combined.

c. Delete any combination having a letter to the left of the colon that is repeated to the right of the colon.

#### II-C. Illustration of the Rules Under II-B (for the example in Sec. 18.3)

- P (types of school), T (treatments), C:P (classrooms nested within P), and R:PCT (replications nested within P, C, T combinations) represent the crossed and nested factors and are lines in the table by rule II-B.1.
- The possible interactions among the factors above include: PT which is retained and may be written TP.

PC: P which is deleted because P appears both before and after the colon (rule II-B.2c).

TC:P which is retained and may be written CT:P.

PTC:P which is also deleted because P appears both before and after the colon.

All the interactions involving R:PCT are deleted because of rule II-B.2c.

3. Thus, the lines of the ANOVA table in our example consist of P, C:P, T, PT, CT:P, and R:PCT.

# I8.7 DETERMINING THE DEGREES OF FREEDOM FOR SOURCES OF VARIATION

#### III-A. Notation

- The number of levels of a factor not nested within any other factor is denoted by the lower case of the letter identifying the factor. In our example, p = 2 and t = 5.
- 2. The number of levels of a nested factor within each level or combination of levels of the factors in which it is nested is denoted by the lower case of the letter to the left of the colon identifying the nested factor. For example, in the nested classification C:P, the number of levels of C in each level of P is denoted by c, which is 3 in our example. In the nested classification R:PCT, r denotes the number of replications within a cell, which equals 2 in our example.
- 3. The total number of observations N equals the product of all the lower case letters for the crossed and nested classifications. In our example, this number is  $p \times c \times t \times r = 2 \times 3 \times 5 \times 2 = 60$ .

#### III-B. Rules

476

- Let a lower case letter correspond to each capital letter. The degrees of freedom for any line in the ANOVA table are found by subtracting one from each lower case letter to the left of the colon and multiplying the grand product of these differences by the grand product of the lower case letters to the right of the colon.
- 2. As a check, the computed degrees of freedom should add to N-1.

III-C. Illustration of the Rules Under III-B (for the example in Sec. 18.3)

> 1. The degrees of freedom for P=p-1=(2-1)=1. The degrees of freedom for C:P=(c-1)p=(3-1)2=4. The degrees of freedom for T=(t-1)=(5-1)=4. The degrees of freedom for

$$PT = (p-1)(t-1) = (2-1)(5-1) = 4.$$

The degrees of freedom for

$$CT:P = (c-1)(t-1)p = (3-1)(5-1)2 = 16.$$

The degrees of freedom for

$$R:PCT = (r-1)pct = (2-1)(2)(3)(5) = 30.$$

2. As a check, 1+4+4+4+16+30=N-1=59.

#### 18.8 COMPUTING SUMS OF SQUARES

#### IV-A. Notation

- Capital letter X will be used to denote an observation on the dependent variable and will have as subscripts all the different lower case letters used in expressing degrees of freedom. For example, X<sub>petr</sub> is used to denote a general observation in our example.
- Lower case letters will also be used to denote the upper limits of subscripts. For example, t = 1, 2, ..., t, which means that the levels of variable t are 1, 2, up to the tth level, which is five in our

#### IV-B. Rules

1. For each line in the ANOVA table write down the degrees of

freedom (df) in their symbolic form (i.e., using lower case letters) and expand algebraically. For example, the line PT has degrees of freedom (p-1)(t-1) which equals +pt-p-t+1. The computational formula for the sum of squares of a source of variation will consist of as many terms as there are terms in the expanded symbolic expression for the degrees of freedom (four terms in the case of the PT interaction), and these terms will have the same algebraic signs as their corresponding terms in the symbolic expression for the df[+,-,-, and + in the case of the expansion of (p-1)(t-1) shown above].

2. For each term in the expanded algebraic representation for the df write a multiple summation corresponding to each subscript of the general observation. Precede the summation with the algebraic sign of the term to which it corresponds. For example, for the term +pt in the algebraic expansion of (p-1)(t-1) one would write

$$+\sum_{1}^{p}\sum_{1}^{t}\sum_{1}^{c}\sum_{1}^{r}X_{pter}.$$

3. For each multiple summation expression, place within parentheses X and those summation signs whose upper limits do not appear in the corresponding term in the expanded expression for the df. For example, for the term +pt one would place the parentheses as follows:

$$+ \sum_{1}^{p} \sum_{1}^{t} \left( \sum_{1}^{s} \sum_{1}^{r} X_{pter} \right).$$

4. Square the expression inside the parentheses and divide by the total number of observations summed over to get the quantity inside the parentheses. This number will be the product of the upper limits of the summation signs inside the parentheses. If no summation sign appears inside the parentheses, one "sums" over 1 value, so the term is divided by 1. The part of the computational formula for the PT interaction that corresponds to the +pt term is then

$$+\sum_{1}^{p}\sum_{1}^{t}\left(\sum_{1}^{e}\sum_{1}^{r}X_{pter}\right)^{2}$$

IV-C. Illustration of the Rules Under IV-B (for the example in Sec. 18,3)

In Sec. 18.7 above, six sources of variation were identified for the example problem. Thus, six sums of squares must be calculated in the analysis of the design. Only the formulas for the sums of squares PT and R:PCT will be

demonstrated. Attempt to write the remaining formulas by following rules IV-B.1 through IV-B.4. Four subscripts are needed to denote a general observation:  $X_{\text{setr}}$ .

1. Sum of squares for PT.

Rule 1. df for (PT) = (p - 1)(t - 1) = pt - p - t + 1.

Rule 2. pt - p - t + 1:  $\sum_{k=1}^{\infty} \sum_{i=1}^{k} \sum_{j=1}^{k} X_{petr} - \sum_{i=1}^{\infty} \sum_{j=1}^{k} \sum_{j=1}^{k} X_{petr} - \sum_{i=1}^{\infty} \sum_{j=1}^{k} \sum_{j=1}^{k} X_{petr} + \sum_{j=1}^{\infty} \sum_{j=1}^{k} \sum_{j=1}^{k} \sum_{j=1}^{k} X_{petr} + \sum_{j=1}^{\infty} \sum_{j=1}^{k} \sum_{j=1}^{k} \sum_{j=1}^{k} X_{petr}$ Rule 3. pt - p - t + 1:  $\sum_{j=1}^{\infty} \sum_{j=1}^{k} \sum_{j=1}^{k} X_{petr} - \sum_{j=1}^{\infty} \sum_{j=1}^{k} \sum_{j=1}^{k} \sum_{j=1}^{\infty} X_{petr} - \sum_{j=1}^{\infty} \sum_{j=1}^{k} \sum_{j=1}^{\infty} X_{petr} - \sum_{j=1}^{\infty} \sum_{j=1}^{\infty} X_{petr} - \sum_{j=1}^{\infty} \sum$ 

The result of the application of rule 4 above is the computational formula for the sum of squares for the interaction of factors P and T.

2. Sum of squares for R:PCT.

Rule 1. If for (R:PCT) = (r - 1)pct = pctr - pct.

Rule 2. pctr - pct:

$$\sum_{1}^{2} \sum_{1}^{c} \sum_{1}^{c} \sum_{1}^{c} X_{setr} - \sum_{1}^{c} \sum_{1}^{c} \sum_{1}^{c} \sum_{1}^{c} X_{setr}.$$
Rule 3. pctr - pct:

$$\sum_{1}^{2} \sum_{1}^{c} \sum_{1}^{c} \sum_{1}^{c} (X_{setr}) - \sum_{1}^{2} \sum_{1}^{c} \sum_{1}^{c} \sum_{1}^{c} (\sum_{1}^{c} X_{setr}).$$
Rule 4. pctr - pct:

$$\sum_{1}^{2} \sum_{1}^{c} \sum_{1}^{c} \sum_{1}^{c} (X_{setr})^{2} - \sum_{1}^{2} \sum_{1}^{c} \sum_{1}^{c} (\sum_{1}^{c} X_{setr})^{2}.$$

The preceding formula yields the sum of squares for replications (or "within").

# 18.9 DETERMINING THE EXPECTATIONS OF MEAN SQUARES

#### V-A. Notation

- 1. The symbol o<sup>2</sup>, having to the left of the colon in its subscript only lower case letters corresponding to random or finite factors, denotes the variance of a random variable underlying those random and finite factors. For example, o<sup>2</sup><sub>c</sub>, denotes the variance of the effects associated with all the classrooms (C) included in the population of classrooms found in a particular type of school (P).
- 2. The symbol σ², having included to the left of the colon in its subscript lower case letters corresponding to fixed factors, denotes a function of the sum of the squared effects of the variables represented to the left of the colon.\* For example, σ² denotes a function of squared fixed effects associated with treatments, e.g., σ² = ∑ α²/<sub>1</sub> (f 1).

#### V-B. Rules

- Unless deleted by rule V-B.2 below, the expectation of a mean square of any factor contains a o<sup>2</sup> for each line in the ANOVA table that has in its denotation all the letters denoting the mean square under consideration.
- 2. Certain of the σ² components of an expectation for a mean square given in V-B.1 above vanish according to the following rule: any σ² having to the left of the colon a letter denoting a fixed classification disappears except when the source of variation of the mean square includes this letter. Remember, if there is no colon in the subscript, the colon is by definition at the right of all letters.
- The coefficient of a particular σ² in a particular mean square includes the product of all the lower case letters not found in the subscript of σ².

Although it has become customary to adopt the o<sup>2</sup> notation, the reader should keep in mind that for factors or combinations of factors involving fixed factors, the o<sup>2</sup> is not a variance of a random variable; it is related to a sum of squared constants (the fixed effects).

V-C. Illustrations of the Rules Under V-B (for the example in Sec. 18.3)

One way of determining the expected values of mean squares is first to list all possible  $\sigma^1$ , then to eliminate selected  $\sigma^2$  by rule V-B.1, then eliminate more  $\sigma^2$  by rule V-B.2, and finally to attach coefficients to the remaining components by rule V-B.3. This procedure will be followed here.

The expected mean square for any line could contain σ<sup>2</sup><sub>p</sub>, σ<sup>2</sup><sub>t</sub>, σ<sup>2</sup><sub>t</sub>, σ<sup>2</sup><sub>t</sub>, σ<sup>2</sup><sub>t</sub>, and σ<sup>2</sup><sub>et</sub>, as well as σ<sup>2</sup><sub>t</sub> pet. In the E(MS) for P, delete σ<sup>2</sup><sub>t</sub> because σ<sup>2</sup><sub>t</sub> does not have a p among its subscripts (rule V-B.1).
 In the E(MS) for C:P, eliminate σ<sup>2</sup><sub>t</sub>, σ<sup>2</sup><sub>t</sub>, and σ<sup>2</sup><sub>tt</sub> because none

of these has both a c and a p among its subscripts (rule V-B.1). In the E(MS) for T, eliminate  $\sigma_p^2$  and  $\sigma_{e,p}^2$  (rule V-B.1).

In the E(MS) for PT, eliminate  $\sigma_p^2$ ,  $\sigma_e^2$ , and  $\sigma_e^2$ , (rule V-B.1). In the E(MS) for PT, eliminate  $\sigma_p^2$ ,  $\sigma_e^2$ , and  $\sigma_e^2$  (rule V-B.1). In the E(MS) for CT: P eliminate all  $\sigma^2$  except  $\sigma_{cl:p}^2$  and  $\sigma^2$  since

none contains all the letters c, t, and p in its subscripts (rule V-B.1). In the E(MS) for R:CTP, eliminate all  $\sigma^2$  except  $\sigma^2$ , since none contains all the letters c, t, p, and r (rule V-B.1).

2. Recall that C and R are considered random, P and T fixed. In addition to σ<sup>2</sup><sub>t</sub>, the E(MS) for P so far contains σ<sup>2</sup><sub>t</sub>, σ<sup>2</sup><sub>t</sub>, σ<sup>2</sup><sub>t</sub>, and σ<sup>2</sup><sub>t+P</sub>. Now σ<sup>2</sup><sub>t</sub> contains fixed factor P to the left of the colon, but the mean square under consideration (P) contains (is) this letter, so σ<sup>2</sup><sub>t</sub> stays, σ<sup>2</sup><sub>t+P</sub> contains only random factor C to the left of the colon, so σ<sup>2</sup><sub>t+P</sub> also stays. σ<sup>2</sup><sub>t+P</sub> contains fixed factor T to the left of the colon and t is not part of the mean square under consideration (P), so σ<sup>2</sup><sub>t+P</sub> is eliminated. For the same reason σ<sup>2</sup><sub>t+P</sub> is eliminated. σ<sup>2</sup> (that is, σ<sup>2</sup><sub>t+P</sub>) always remains since only the random factor R is to the left of the colon.

In addition to  $\sigma^2$ , the E(MS) for C:P contains so far only  $\sigma^2_{rp}$  and  $\sigma^2_{trp}$ ,  $\sigma^2_{rp}$  is retained because C is random, and it is also retained since c is part of C:P. However,  $\sigma^2_{trp}$  contains t to the left of the colon and T is both fixed and t is not part of the mean square under consideration (C:P). Thus  $\sigma^2_{trp}$  is eliminated.

In addition to  $\sigma^2$ , the E(MS) for T contains so far  $\sigma^2_i$ ,  $\sigma^2_i$ , and  $\sigma^2_{\sigma P}$ ,  $\sigma^2_i$  is retained because t is part of T.  $\sigma^2_p$  is eliminated because T is part of T. The mean square under consideration.  $\sigma^2_{\sigma P}$  survives because C is random. Note that P is to the right of the colon, and thus not affected by rule V-B.2.

<sup>\*</sup> The expected mean square E(MS) for the random factor replications within the smallest cell is usually denoted simply as  $\sigma^s$  or  $\sigma^s$  ( $\epsilon$  for "error"). This practice has been followed here.  $\sigma^s$  always appears as a source of variation even if r=1, in which case there are no degrees of freedom for this source.

48 I

Similarly, it can be shown that the E(MS) for PT is  $\sigma^2$ ,  $\sigma_{n_1}^2$ , and  $\sigma_{cl:n}^2$ ; and the E(MS) for CT:P is  $\sigma^2$  and  $\sigma_{cl:n}^2$ .

3. The coefficients of the surviving components are found by a straightforward application of rule V-B.3. For example, the coefficient of  $\sigma_{ep}^2$  is  $(t \times r) = (5)(2) = 10$ . See Table 18.10 for the final results,

**TABLE 18.10** 

10.10	Source of variation	OVA TABLE (FOR df	THE EXA	AMPLE IN SEC. 18.3)  E(MS)		
_	P	(p-1)	= 1	$\sigma^2 + 10\sigma^2$	+ 30σ <sup>2</sup>	
	C:P	p(c-1)	= 4	$\sigma^2 + 10\sigma^2$		
	T	(t - 1	1) = 4		$\sigma_{ot;p}^{2} + 12\sigma_{i}^{2}$	
	PΤ	(p-1)(t-1)	1) = 4	$\sigma^2 + 2\epsilon$	$\sigma_{et}^{2} + 6\sigma_{et}^{2}$	
	CT:P	p(c-1)(t-1)		$\sigma^2 + 2c$	72	
	R:PCT	pct(r-1)	<b>≃ 30</b>	$\sigma^2$	9	
	Total	$pctr - 1 = N \sim$	1 = 59			

#### 18.10 SIGNIFICANCE TESTING

It is usually the researcher's purpose to test one or more null hypotheses, i.e., that particular variance components, fixed effects, or mixed effects are zero. To test any one such hypothesis, one first identifies the source of variation corresponding to the variance component, fixed effect, or mixed effect in question. The mean square for this source of variation becomes the numerator of the F-ratio, the test statistic used in testing the null hypothesis. One then determines what the expected value of this mean square is when the variance component, fixed effect, or mixed effect in question is zero. Call this new expected value, E(MS | Ho true). The denominator of the F-ratio is the mean square that has the expected value  $E(MS \mid H_0 \text{ true})$ . In other words, the null hypothesis is tested with the ratio of two mean squares such that both would have the same expected value if the null hypothesis in question were true. (The mean square in the numerator corresponds to the source of variation being tested.)

Suppose in our example we were interested in the hypothesis that  $\sigma_{\rm s}^2 = 0$ . Since the expected value of the mean square for P contains  $\sigma_{\rm s}^2$  and since the expected value of P equals the expected value of C:P when  $\sigma_{z}^{2} = 0$ , the ratio of the mean square for P to the mean square for C:P is appropriate for testing the hypothesis that  $\sigma_n^2 = 0$ .

If no appropriate F-ratio exists for testing an effect, approximate methods are sometimes applicable. (See, e.g., Winer, 1962, pp. 199-202.)

In spite of the fact that the above significance testing procedures are common practice, a somewhat more enlightened skepticism concerning the distribution properties of F-ratios in mixed designs (i.e., having both fixed and random factors) would seem to be called for by the considerations raised in Sec. 18.2 concerning mixed model assumptions and in Scheffe (1959, chap. 8) for example.

#### PROBLEMS AND EXERCISES

1. A random sample of J = 10 judges was drawn from a population of judges. Each judge rated an independent random sample of n = 20 children on a 7 point scale of "emotional adjustment." The analysis of variance yielded the following

Source of variance	df	MS	E(MS)
Between judges Within judges	9	10.48 9.64	$\frac{\sigma^2 + 20\sigma_a^2}{\sigma^2}$

- a. Estimate the variance of judges,  $\sigma_a^2$ , from the above data.
- b. Compare the sizes of  $\delta_a^2$  and  $\delta^2$ . Who differs more, children being rated by the same judge or judges on the average across all children?
- c. Establish the 95% confidence interval on  $\sigma_0^2/\sigma^2$ .
- 2. A population of 30,000 spelling words is identified. A measurement researcher is interested in the variability to be expected among 50-item spelling tests that could be formed from the pool of 30,000 words. The total number of possible 50-item tests is equal to the combinations of 30,000 items taken 50 at a time, i e.,  $C_{50}^{20,000}$ , which is a staggeringly large number. The researcher constructs six 50item spelling tests by randomly sampling items from the item pool. Thus, the six tests can be considered to be randomly sampled from the population of all possible 50-item tests. Each test is given to an independent random sample of n=12 pupils. The following test scores (total number of correct spellings out of

1		Test				
<u> </u>	2	3	4	5	6	
21 14	34	44	12	30		
11	19 26	31 36	26	18	27 32	
27 19	31	24	22 30	14 29	26	
32	39 42	40 38	35	36	29 34	
21 23	27 14	42	14 19	27 30	20	
18	25	35 27	24	25	15 25	
25 24	29 33	29	23 31	25 20	31	
23	36	30 33	27 25	32	39 23	
ing the or	ne-factor ran	dom .e	43	19	29	

g the one-factor random-effects ANOVA, calculate the estimate of the

variance among all possible spelling means,  $\hat{\sigma}_a^2$ , and the estimate of the variance of pupils' scores within any spelling test,  $\hat{\sigma}^2$ . That is, calculate

$$\hat{\sigma}_a^2 = \frac{MS_A - MS_w}{12} \quad \text{and} \quad \hat{\sigma}^2 = MS_w.$$

b. Calculate the 95% confidence interval around  $\hat{\sigma}_{\alpha}^{2}/\hat{\sigma}^{2}$  on  $\sigma_{\alpha}^{2}/\sigma^{2}$ .

 $E(MS_{-}) = \sigma^2$ 

 In a two-factor random-effects ANOVA with I levels of factor A, J levels of factor B, and n observations per cell, the expected values of the mean squares are as follows:

$$\begin{split} E(MS_A) &= \sigma^2 + n\sigma_{ab}^2 + nJ\sigma_a^2, \\ E(MS_B) &= \sigma^2 + n\sigma_{ab}^2 + nI\sigma_b^2, \\ E(MS_{AB}) &= \sigma^2 + n\sigma_{ab}^2, \end{split}$$

Find a linear combination of the mean squares that provides an unbiased estimator of  $\sigma_{\sigma}^2$ . [Hint: Notice how  $\sigma_{\delta}^2$  can be estimated:

$$\begin{split} E\frac{MS_B - MS_{AB}}{nI} &= \frac{E(MS_B) - E(MS_{AB})}{nI} \\ &= \frac{\sigma^2 + n\sigma_{ab}^2 + nI\sigma_b^2 - \sigma^2 - n\sigma_{ab}^2}{nI} = \frac{nI\sigma_b^2}{nI} = \sigma_b^4. \end{split}$$

4. It has often been maintained that neurologically handicapped children evidence a lower Performance IQ than Verbal IQ on the Wechsler Intelligence Scale for Children (WISC). Hopkins (1964) compared the Verbal and Performance IQ's for a group of about 30 children ranging in age from six years to 12 years who were diagnosed as neurologically handicapped independently of the intelligence testing. The following data were obtained:

Person	Verbal I <u>Q</u>	Performance IQ	Person	Verbal IQ	Performance IQ
1	87	83	16	83	85
2	80	89	17	83	77
3	95	100	18	92	84
4	116	117	19	95	85
5	77	86	20	100	95
6	81	97	21	85	99
7	106	114	22	89	90
8	97	90	23	86	93
9	103	89	24	86	100
10	109	80	25	103	94
11	79	106	26	80	100
12	103	96	27	99	107
13	126	121	28	101	82
14	101	93	29	72	106
15	113	82	30	96	108

Consider the "Persons" to be a random factor and the "Verbal IQ vs. Performance IQ" to be a fixed factor. Using the mixed-ellects NNOVA model, text the null hypothesis at the .05 level that populations of Verbal and Performance IQ's have the same mean for neurologically handicapped children. (The Fiest will lead to the same decision that would be arrived at if a dependent-groups t-test of the hypothesis #1 = #1 & were made.)

Scores on the Information, Vocabulary, Digit Span, and Block Design subtests
of the Wechsler Intelligence Scale for Children are tabulated below for a group of
12 neurologically handicapped children (Hopkins, 1964);

	Person	Infor.	Test Voc.	Dig. S.	Bl. Des.
-	1	7	8	7	7
	2	5	10	8	12
	3	9	11	9	11
	4	17	18	9	13
	5	4	7	7	9
	6	6	9	8	11
	7	11	11	7	7
	8	10	14	12	7
	9	8	11	7	13
	10	12	11	5	9
	11	13	16	6	18
	12	11	10	11	5

WISC subtest scores are scaled to a mean of 10 and standard deviation of 3 for the general population. It has often been asserted that patterns of subtest scores on the WISC can be used to diagnose neurological handicaps. Test the null hypothesis that the twelve test scores above were randomly sampled from four normal distributions with the same mean.

The design should be regarded as a repeated measures design; hence, in performing the F-test use the flow chart in Fig. 18.3.

- 6. Twenty raters are drawn at random from a population of 10,000 raters. Thirty-two rates are drawn at random from a population of 20,000 rates. Eight traits are "drawn at random" from a population of eight traits (i.e., all traits of interest to the investigator for this particular study are used). Each of the 20 raters rated each of the 32 rates once on each of the eight traits in a well-conducted study.
  - a. How many ratings does this yield?
  - b. Which factors are "random" and which "fixed"? Are any factors "nested" within any other factors? Which?
  - c. How many sources of variation are there in these ratings? List them.
  - d. Using the rules of thumb given in the chapter, work out all the expected mean squares for this design, indicate the appropriate ratio of mean squares and the degrees of freedom for each F, and provide the formulas for estimating all estimable components of variance.

For another way to analyze data from this particular design, see Stanley (1961b).

7. Two raters who were Democrats and two raters who were Republicans each rated three ratees who were prominent Democrats and three ratees who were prominent Republicans on each of four different traits. This yielded a total of 2 x 2 x 2 x 3 x 4 = 96 ratings. The design is shown below. There are five factors, two of which are "nested."

			Trait rated, Party of rater, Rater number															
	Party Ratee		Intelligence				Honesty			F	Friendliness				Generosity			
ratee number		DR		D		R	D		R		D		R					
L		1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	
	1	7	6	8	7	8	2	8	7	5	1	8	2	5	0	10	5	
R	2	7	7	6	7	7	6	6	6	6	6	9	1	5	6	8	2	
	3	6	6	4	5	8	9	8	6	10	5	9	8	9	5	10	6	
	4	8	8	6	6	9	10	3	1	8	8	7	6	9	7	4	4	
D	5	8	5	6	7	9	5	4	4	5	1	4	1	9	1	3	1	
	6	5	4	3	3	7	8	1	0	7	7	8	4	7	8	0	0	

- a. Identify the two nested factors. Within what is each such factor nested?
- b. Do the nested factors "cross" any other factors? Which ones?
- c. Which factors does the "political party of rater" factor cross?
- d. Which of the factors were probably considered as each having had its levels drawn randomly from an infinite (hypothetical) population of levels?
- e. 62 (partyofrater x partyofratee) contributed most to variation of the ratings. Would you have expected this result in advance of the rating procedure? What does this interaction probably mean?
- f. The second largest estimated component of variance was that for party of rater × party of ratee × trait. What does this three-factor interaction mean? (One sometimes sees a three-factor interaction referred to as a "second-order" interaction, because a zero-order "interaction" would be a main effect, not interacting with anything. Thus a two-factor interaction such as that in (e), above, may be called a "first-order" interaction.)

For further results, see Stanley (1961a).

### 19

# FUNDAMENTALS OF FXPERIMENTAL DESIGN\*

#### 19.1 INTRODUCTION

The word "experimentation" has come to have many meanings for behavioral scientists. The most common meaning night be termed "experientation," trying new approaches and subjectively evaluating their effectiveness. In this chapter we are concerned with a more structured inquiry, akin to that carried out in many of the sciences. It involves control by the experimenter of at least one variable, such as method of teaching arithmetic, that he can manipulate. Thus, experimentation of this kind differs from observation of naturally occurring events in that the stage has been set by the experimenter so that the possibly differential effects of at least two "treatments" can be observed in a situation where assignment of experimental unity (often, pupils or classes) to the several treatments has been made without bias. "Nature" almost always makes biased assignments to its treatments; even before a natural experiment, the experimental units to be subjected to one treatment are usually not comparable to those to be subjected to another treatment.

<sup>\*</sup> For a simple, general approach to this topic see Stanley (1967b).

For example, as a group, smokers differ in many ways from nonsmokers, any one of which ways might be the cause of a greater incidence of lung cancer among smokers.

In brief, nature rarely assigns experimental units to treatments randomly, whereas the careful experimenter almost always does. One can define a controlled, variable-manipulating, comparative experiment as a study in which the available experimental units are assigned at random (either simply or restrictively) to the various treatments. More generally, if we consider each factor to be manipulated (e.g., several ways to teach arithmetic; sex; overt versus covert response in programmed instruction; or 100% reinforcement versus 50% reinforcement) as having two or more feets or categories (e.g., SMSG\* curriculum versus two other ways to teach mathematics would constitute three levels of the teaching-of-mathematics factor), we can then talk about factor-level combinations, such as the six generated by three ways to teach mathematics crossed with two levels of sex (male-female). This is a 3 × 2 factorial design—one factor at three levels crossed with a second factor at two levels.

The basic experimental design involves one or more factors that are either manipulated by the experimenter, as illustrated above, or not manipulated by him (e.g., male or female sex, day of the week, height above average or below average). The levels of one factor may be crossed with those of another, or the levels of one or more factors may be nested within the levels of another factor. For instance, when three male raters and three female raters rate each of 10 ratees on each of seven traits, raters are nested within sex, because no male rater is also a female rater (i.e., the rater levels do not cross the sex levels, though they do cross the ratee and the trait levels). You have already seen examples of nesting in Chapter 18, including Prob. 7 at the end of that chapter.

Another example would be an experiment in which the manipulated factors were immediate reinforcement versus delayed reinforcement and 50% reinforcement versus 100% reinforcement, all levels of one factor crossing all levels of the other, with the experimental subjects (i.e., persons) classified as male versus female, blond versus brunette versus redhead, and volunteers versus novolunteers. The subjects would be nested within the 12 "nests" created by the intersection of sex with hair color with volunteering. One-fourth of the subjects in each nest would be assigned randomly to a given factor-level combination of the manipulated factors.

Hierarchical nesting occurs when some factors are subclasses of others; e.g., cities could be nested within counties, counties within states, and states within regions. An experimenter might assign cities at random to experimental treatments. He would then have a partly nested and partly crossed

SMSG means "School Mathematics Study Group," a group composed of mathematics educators who pioneered in bringing modern mathematics to secondary-school courses in the United States.

design, with the geographical units not crossing each other but crossing the levels of the treatment factor. One then has no basis for concluding that a certain city would have done relatively better in the experiment had it been in another state.

Nesting and crossing can occur only when there are at least two factors. In a given study there may be no nesting, no crossing, all of one or the other, or a mixture of nesting and crossing, as in the above examples. Note, though, that nesting rarely occurs when all the factors are manipulated.

The term "factorial design" is taken by some statisticians to mean a fully crossed set of two or more factors, with an equal or unequal number of experimental units assigned completely randomly to each of the factor-level combinations. (See Kendall and Buckland, 1957, p. 106.) For one factor, with random assignment of n, experimental units to each of the J levels of the factor, there would be no crossing, of course. Other definitions of factorial design include cases where only restricted assignment of the experimental units is possible, as when the N available experimental units must be considered as having been drawn from two or more different populations (e.g., men versus women); each such population defines a level of a nonmanipulated factor.

(See Stanley, 1967, pp. 204-205, for an argument about whether "randomized-block designs" are factorial designs. We shall see subsequently in this chapter that they involve full crossing of factor levels, but with the restricted randomization that necessarily results when a given experimental unit already has "tied" to it a level of a factor such as sex or school attended. No harm will be done if one refers to any design created by the crossing and/or nesting of factor levels as a factorial design, provided that distinctions among such designs that are important for interpreting the results of the experiment or status study are not overlooked.)

#### 19.2 AN EXAMPLE

For concreteness, let us consider a controlled experiment involving four styles of printing type crossed with three sizes of printing type, a 4 × 3 factoral design yielding 12 factor-level combinations: each style is tried with every size, and each size is tried with every style. This is a complete design. If we take 12 pupils and assign one at random to each of the 12 assign 24 pupils, two at random to each of the 12 combinations, we produce one replicate. Usually, as a minimum, we shall excrete two replicates. In order to have a powerful enough experiment, we may need to assign more than two pupils to each combination. Some number of replicates, n, will yield the power we require. Methods for determining n exist.

SEC. 19.2

How do we conduct this factorial-design experiment involving four styles crossed with these sizes? Of course, we must decide which styles and sizes to use. Probably we have firmly in mind four styles that are candidates for use in the textbook or test we plan to prepare. We also know which three sizes to try for our purposes—perhaps 8 point, 12 point, and 16 point. These four styles and three sizes produce 12 factor-level combinations that are the only ones of interest to us in this experiment. In the jargon of experimental design, we have two fixed-effects factors and therefore employ a fixed-effects model, because we have "drawn" the four styles from a target population of just three sizes.

Where do we get the experimental units with which to do the experiment? We might secure a "grab-group" consisting of the first 12n individuals who happen our way, or we might define a population of individuals, such as all fourth graders in a large school system, and draw 12n individuals at random from that population. Using a grab-group limits rigorous statistical generalization from the outcome of the experiment to just those 12n persons. whereas drawing the experimental units randomly from a population permits statistical generalization to that population, thereby increasing external validity, i.e., generalizability. (We may, however, be able to generalize nonstatistically from the members of the grab-group to other persons "like them" if we know enough about the adventitiously chosen individuals to be reasonably sure that none of their characteristics determining the outcome of the experiment differ enough from those of the target population to change the results there. This is difficult to ascertain, and in any event we have no probabilistic warrant for generalizing from the grab-group to anyone else whatsoever.)

After we secure the 12n experimental units by one of the above methods, we randomly assign n of them to each of the factor-level combinations, which have common content but different combinations of style and size. We wish to vary only style and size. All other variables should be held constant (as for example, by using just males in the experiment, thereby keeping the sex factor at one level) or randomized over all 12 factor-level combinations. This is where experimental control becomes crucial. It might be practicable, for instance, to seat all 12n pupils randomly in the room. (That could be done by passing out random seat assignments at the door as the pupils arrived.) If a nonrandom room arrangement were used, this would have to be considered as part of the design, making it more complex than a 4 × 3 factorial. Control of extraneous variables calls for great care and ingenuity so that the experiment will be internally valid, i.e., produce comparisons that are free from bias.

After the 12n pupils have read the same passage with the same time limit under the same conditions except for style and size of printing type, they will

be given a common test to determine how much each learned from the particular combination of style and size of type to which he was exposed. The total score of each pupil on this outcome test will constitute the observations on the dependent variable to be analyzed.

#### 19.3 ADVANTAGES AND DISADVANTAGES OF THE FACTORIAL DESIGN

Complete, balanced factorial designs (designs where each of the possible factor-level combinations occurs and has  $n \ge 1$  experimental units assigned to it) permit testing more than one hypothesis about main effects (e.g., the influence of type size or the influence of style) efficiently in the same experiment. They also make it possible, where two or more factors are used, to study how the factors interact. Perhaps the least effective of the three sizes when combined with the least effective of the four styles does not produce the least effective of the twelve factor-level combinations. If the effects are additive, so that knowing the effectiveness of a certain size factor level and a certain style factor level one can predict the effectiveness of that factor-level combination as well as "chance" permits, then we say that the two factors do not interact. One cannot study interaction statistically in one-factor studies, and yet appreciable interaction may greatly limit generalization of the results of one-factor experiments.

The factorial design is relatively simple. It does not require any premeasurements or "matching" of the experimental units, because bias is
avoided (in the probabilistic sense) by the randown assignment of the experimental units to the factor-level combinations. On the other hand, it permits
the within-factor-level-combination variability to be as great as true-score
variation among individuals treated alike plus errors of measurement dictate.
Thus the signals (the genuine effects) may be drowned out by the noise (the
within-combination variability) if the signal-to-noise ratio is low. This noise
(or error, as it is usually called) lowers the power of the significance tests used
and increases the width of the confidence intervals computed. True-score
error can be lessened by various techniques such as blocking, stratifying,
lereling, and corarying, all of which depend on classification or pre-measurement of the experimental units and are used to reduce the within-combination
variability of true scores. Measurement error can be reduced somewhat by
using a more reliable outcome measure.

But more important, sampling error in the estimation of main and interaction effects can be reduced (and thus power of significance tests increased) simply by making n larger, provided that the larger experiment can be conducted as efficiently (i.e., no greater error per observation) as the smaller one.

#### 19.4 BLOCKING

Of course, one can often lower within-factor-level variability of the outcome measures by drawing the experimental units from a homogeneous subpopulation-e.g., persons all the same age, sex, IO, and socio-economic level. This may reduce "error" considerably, but it does so at the expense of limiting the generalizability of the findings to other persons of the same age, sex. IO. and socio-economic level as the ones used in the experiment. It will usually be better to set up explicitly as factors in the experiment those characteristics thought likely to be most closely related to the outcome measure(s) of the study-i.e., to the dependent variable(s). This factorialization will reduce error almost as well as would the subpopulation method. One can then test the interactions of the status variables with the manipulated variable(s) to determine whether or not the findings can be generalized over age groups, sexes, etc. Indeed, the thoughtful experiment designer often has his cake and eats it, too, as Ronald Fisher showed convincingly long ago.

One of the earliest crossed classifications developed by Fisher for agricultural research was the randomized-block design. (See Fisher, 1925, pp. 226-29.) If V different varieties of wheat were to be planted in a field, he suggested that the field first be divided into B blocks of ground, the fertility within a given block being as homogeneous as possible. Then each block was divided into V plots, one for each variety of wheat. One variety was assigned at random to each plot within a given block, so that on each block all V varieties were planted once each. This, then, produced  $B \times V$  factorlevel combinations and BV observations when the wheat matured. See Table 19.1 for the layout of the observations.

Note that varieties were assigned randomly to plots within blocks. Fisher showed that this randomization was crucial. Also, the B blocks in the experiment might be considered a random sample drawn from a hypothetical population of blocks "like these," whereas the V varieties were probably the "target population" of varieties of wheat, i.e., all the varieties in which the experimenter was interested. This is a mixed model:random block effects and fixed variety effects. The E(MS)'s shown in Table 19.1 are such that

MS and the MS (North V parieties)

is distributed as

 $F_{V-1,(R-1)(V-1)}$ 

under the null hypothesis that the V varieties produce equal quantities of wheat.

By choosing homogeneous parts of the field (the blocks) rather than merely assigning the varieties wholly at random throughout the field, Fisher removed the between-block variance from the mean square for error and

TABLE 19.1 OUTUNE OF DATA FROM A RANDOMIZED-BLOCK-DESIGN EXPERIMENT:  $X_{**}$ NOTATION, WHERE  $b=1, 2, \ldots, 8$  BLOCKS AND  $v=1, 2, \ldots, v$  VARIETIES
OF WHEAT; RANDOM BLOCKS, FIXED VARIETIES

	Variety of wheat						
Block of ground	1 2	v					
1	X <sub>11</sub> X <sub>21</sub>	X <sub>1</sub> ,					
2	X21 X15	X <sub>17</sub>					
<del></del>		:					
:							
В	X21 X21	Xer					
Source of variation	ql	E(MS)					
Between blocks (a)	B 1	$\sigma^2 + V\sigma$ $\sigma^1 + \sigma^2_{\alpha\beta} + B\sigma$					
Between varieties ( $\beta$ ) Blocks × varieties ( $\alpha\beta$ )							

increased the power of the significance test for varieties. He was not much interested in MS<sub>stets</sub>, except that it should be as large as possible under the conditions of a particular experiment. The general principle of homogeneous subsorting of experimental material can be extended beyond agriculture to a number of situations of great interest to behavioral scientists. In some of these, blocks cannot reasonably be considered a random-effects factor. Also, in many of them there will be more than one observation per factor-level combination. (Fisher allowed for that in some situations, but the more plots there were per block, the less the within-block homogeneity was in many agricultural situations.)

Our discussion below is based on three scales of measyrement: nominal-star factor, which we call a blocking variable; ordinal, which we call straighting; and interval or ratio, which we call kerding. These three terms are our own coinage for this situation. We believe that they serve behavioral science more usefully than does the single, ancestral expression "randomized-block design." Blocking, stratifying, and leveling variables are classificatory, rather than manipulated. They are antecedent to the beginning of the experiment itself, as for example when the field was divided into blocks before the varieties were planted on plots within these blocks. Let us now consider each of these three types of factorialization that are useful for reducing error and improving generalizationability.

SEC. 19.4 BLOCKING 493

### Blocking on a Nominal-Scale

A familiar example of blocking is the classification of each pupil as being either male or female and the introduction of this two-level factor explicitly into the experimental design. Physiological sex is not a manipulated variable, but in the experimental design it is treated statistically like the manipulated factors. With two levels of sex, three levels of type size, and four levels of type style one has  $2\times 3\times 4=24$  factor-level combinations and needs 24n experimental units.

Main effects of sex, size, and style can be estimated. Also, one can study the interaction of sex with size, sex with style, and size with style, and the three-factor interaction of sex, size, and style. If the main effect of sex is significant, then by having a sex factor one has reduced the error variance significantly. If sex interacts with either or both of the manipulated factors, then by having sex as an explicit factor one has learned how to limit his generalizations appropriately. For example, it might be discovered that women find Style 3 easiest to read, whereas men find Style 1 easiest. When pursued further, this might have practical consequences for the design of textual materials.

Another example of blocking is the use of identical twins in an experiment where one factor is manipulated at two levels. Twin A of each pair is assigned at random to a level of the manipulated factor, and twin B of that same pair then receives the other level. With P pairs of twins, one has 2P experimental units. The three sources of variation are between twin pairs, between the two treatments, and interaction of pairs with treatments. Note that this is just one replicate, so no direct statistical test of the interaction is afforded. If the variation among the twin pairs is significant, one has reduced the error term significantly, but because one sacrifices half his degrees of freedom for error in so doing, the power of the statistical test of the two-level treatment effect may not be improved or may even be lessened. In this design, the factor "twin pairs" would almost surely be regarded as random, since one would probably want the comparison of A and B to be generalized to a population of twins. Incidentally, this is a social- or biological-science version of Fisher's randomized-block design. The twin pairs are unordered, having been "measured" on a nominal scale.

### Stratifying on Ordinal-Scale Variables

We call an ordinal-scale variable used as an explicit factor in the experimental design, such as socio-economic status of each experimental unit, a stratifying variable. There might be five levels, such as high, upper-middle,

middle-middle, lower-middle, and low socio-economic status, creating a five-level classificatory (i.e., not manipulated) factor.

#### Leveling on Interval- or Ratio-Scale Variables

An interval or nearly interval scale or a ratio scale can be used to yield what is called a leveling variable. To reduce within-factor-level-combination variability one may group the experimental units before the experiment begins

TABLE 19.2 SCHEMA OF DATA FOR THE LEVELING DESIGN, WITH REPLICATION:  $X_{tt}$  NOTATION,  $I=1,2,\dots,L$  LEVELS,  $t=1,2,\dots,T$  TREATHENTS, AND  $I=\dots,n$  INDIVIDUALS FOR EACH TREATHENT WITHIN EACH LEVEL\*

Levels		Treatme	ents	
Zetis	1	2	····	T
ı	X <sub>111</sub> X <sub>113</sub>	X <sub>111</sub> X <sub>112</sub>		X <sub>171</sub> X <sub>172</sub>  X <sub>17a</sub>
2	X <sub>111</sub> X <sub>111</sub>	X <sub>111</sub> X <sub>111</sub>	:::	X <sub>171</sub> X <sub>172</sub>  
:	:	:		:
L	X <sub>L11</sub> X <sub>L12</sub> X <sub>L14</sub>	X <sub>L11</sub> X <sub>L11</sub>	:::	X <sub>LT1</sub> X <sub>LT1</sub>
	A 414	Xtta		Xir.

One can let i = 1,2..., n<sub>n</sub>, instead of constant n, especially when levels and treatment are both facel effects factors, as they usually will be. This may be preferable when the number of individuous about a certain levels of a natural or desired classification differs from that variable at other a Becal off consult again the discussion of proportional cell-frequencies analyses. Here the proportionality condution would be that n<sub>n</sub> = (n, n, n, n).

on something, such as measured reading comprehension or height, that is expected to correlate well within treatments with the outcome measure of the experiment. If there are T levels of a treatment factor and LT experimental units, one would arrange the experimental units from highest to lowest on the pre-measured factor into L levels. Within each such level, one experimental unit would be assigned at random to each of the T treatments (i.e., the T levels of the manipulated variable), creating one replicate of an  $L \times T$  design. If the measures of the leveling factor do correlate significantly greater than zero with the outcome measures, then (as in the twin design outlined above) the within-treatment variability will be reduced significantly

Alternatively, one might choose to use N=nLT experimental units, where n is greater than 1. (In the above paragraph, n=1.) Then one would group the N experimental units, from highest to lowest, into  $L=N_{\parallel}nT$  sets, and would assign at random n experimental units to each treatment within each level. This would permit testing the interaction of levels with treatments, which the n=1 design does not allow directly. See Table 19.2 for an outline of this design.

#### 19.5 ORDERED LEVELS OF FACTORS

If one has three equally spaced sizes of printing type, such as 8, 12, and 16 point, he has three equally spaced levels of an ordered factor. A significant trend for size of type might be linear, representing an equal increment (or decrement) as one goes from 8 point type to 12 point type and from 12 point type to 16 point type. The trend might be quadratic (i.e., second-order), as when 8 and 16 are equally effective but 12 is much better, or it might combine both linear and quadratic components.

Style of printing type, you note, is a nominal-scale factor, not ordered. It is quite possible to have two or more ordered factors in the same study, however, as for example if one introduced five equally spaced weights of paper into the print study; the various trends that result could then be evaluated. (E.g., see Winer, 1962, and Edwards, 1968.)

# 19.6 RANDOM SELECTION OF FACTOR LEVELS

Earlier in this chapter it was hinted that the four styles of printing type might have been drawn at random from a larger target population of printing styles to which one wished to generalize. If that population contained, say, 40 styles, then the four drawn would be 10% of the entire population, a small but hardly negligible percentage. If, on the other hand, one drew four schools

out of a target population of 4000 schools, the one-tenth of 1% that the schools in the experiment constitute of the population would be tiny, so one might choose to consider that essentially the four schools had been drawn from an infinite population of schools, in which case one would (using the jargon of the field) say that the schools are a random-effects factor.

If one has both fixed-effects and random-effects factors in an experiment, we say that he should use a mixed-model analysis of his results. (See Chapter 18 for the mixed-effects ANOVA model.) Genuine random-effects factors seem rare in educational and psychological research, but often we choose to act as if the levels of a factor such as teachers, schools, classes, ratees, or twin pairs had been drawn randomly from a virtually infinite population of such levels. We do this by arguing that the particular levels used are plausibly a random sample from a hypothetical population to which we wish to generalize: teachers "like these," schools "like these," etc. This logic has the endorsement of several top-level mathematical statisticians who have debated it rather hotly with each other. (E.g., see Cornfield and Tukey, 1956.) If we are going to generalize to other teachers, other schools, and the like anyway, then we should use the analytical model that fits such generalization, rather than using the model that applies just to the particular levels in

#### 19.7 NATURAL AND CONTROLLED EXPERIMENTS

Sampling factor levels from a population of factor levels larger than the number to be used in the experiment brings controlled experimentation closer to the methodology of sample surveys than it was until the early 1950's. A difference that persists is the manipulation by the experimenter of the levels of one or more factors, along with the random assignment of the experimental units to the factor-level combinations. Sampling of occupations might occur in a survey of salaries, where respondents are classified by occupation, sex, and marital status, as when the occupations to be studied are drawn at random from a large list of occupations. The analysis of the results of such a study might proceed formally in much the same way as that for an experiment involving styles of printing type, sex, and marital status, but in the status study nothing was manipulated, whereas in the experiment style of type was. Of course, one can conceive of an experiment in which experimental units representing the crossing of the two sexes and three marital statuses are assigned at random to occupations, thereby distributing ability, interests, education, age, and the like randomly across the occupations and in this way removing the confounding of occupation with those personal characteristics.

Generally, interpretation of the results of a controlled experiment will

be easier than that of the analogous "natural experiment." A more familiar example than the occupational one is investigation of the effects on general English vocabulary of studying Latin in high school. If students elect (i.e., volunteer) to take Latin or not to take it, the inputs for the two conditions will almost always be substantially different, the students taking Latin being better initially on English vocabulary, IQ, and a host of other cognitive and affective variables. If, however, half the prospective enrollees in Latin could be assigned at random to Latin in the ninth and tenth grades and the other half in, say, the eleventh and twelfth grades, nonreactively so that disappointment and frustration did not upset the operation of the school, it should be possible to compare both groups unbiasedly at the end of the tenth grade, after half had completed two years of Latin and the other half had taken none. Because of the random assignment, there would be no systematic confounding of any antecedent variables with the experimental variable (i.e., took Latin versus did not take it). Results should be much more readily interpretable than in the natural experiment.

One has only to recall the great difficulties encountered by statisticians when analyzing the results of the vast natural-smoking experiment that has been going on for many years. Is cigarette smoking one of the potent "causes" of lung cancer? Of other ills? After much comparison of human subgroups and animal experimentation in order to discredit plausible alternative hypotheses, most researchers in this area have concluded that smoking cigarettes does increase the probability that a person will develop lung cancer and have certain other ailments, but because the work with humans was not controlled experimentation, no proof overwhelmingly convincing to all intelligent persons has yet been provided. Indeed, associational analyses cannot eliminate all plausible alternative hypotheses, whereas controlled experiments if conducted impeccably can rule out all systematic ones, leaving only chance fluctuations (usually of quite low probability and largely under the experimenter's control) as the alternative explanation. This is not to say that a single experiment can be definitive or perfect; none ever is. Often an experiment will raise more new questions than it answers old ones, but at least the process of randomized assignment of experimental units to factor-level combinations removes the chief source of systematic bias that afflicts most natural experiments.

Control may exact a high price in terms of lowered external validity (i.e., generalizability), however. For example, one probably cannot assign persons at random to unidentified-flying-object (UFO) versus non-UFO clubs and preserve the sense of the distinction as it occurs naturally. One might try paying some persons to smoke and others to refrain from smoking, but very likely this would not simulate well enough the natural situation where persons of certain temperaments and backgrounds cannot resist moking several packages of cigarettes daily, whereas other types of individuals

are not tempted. One cannot very well assign occupations randomly in a meaningful way. Even assigning Latin versus no Latin to eligible high-school students by deferring this subject for half of them has never been done, so far as we are aware. However, this technique of delaying treatment for a randomly selected group of control subjects has been employed frequently in evaluating the effects of psychotherapy.

But in general, researchers in the behavioral and social sciences and education have not made use of the simple and powerful expedient of random assignment to comparison groups even when to do so would be uncomplicated. There do not even seem to be any controlled experiments involving cursive handwriting versus manuscript handwriting in the primary grades, despite the fact that all school children throughout the country are taught to write by some combination of such methods.

In this chapter we have dealt only with attempts to make causal inferences from comparative data. Questions such as "What is the distribution of the ages of female first-grade teachers in California?" and "Are there more male than female elementary-school principals in the United States?" are examples of perfectly reputable inquiries not directly concerned with causal influences. Without a great store of such information, one is hardly ready to design controlled experiments. We can get answers to important questions by the use of questionnaires, interviews, and case studies. Survey researchers have developed powerful analytical techniques for inferring causation from status data, too. For examples, see Bialock (1964).

#### 19.8 OTHER EXPERIMENTAL DESIGNS

Because this is meant to be a systematic textbook, rather than a handbook, we have not attempted to explain or even mention previously a number of experimental designs that are sometimes of value to behavioral scientists, including educational researchers. Among these are Latin and Greco-Latin squares, balanced and partially balanced incomplete-block designs, and fractional-factorial designs. The interested reader is at this point equipped to consult specialized books for these. The most comprehensive manual is probably Cochran and Cox (1957). A simpler treatment, less statistical but still rather comprehensive, is by Cox (1958). Winer (1962) deals with many squares, fractional replication, and randomized-block designs, see Edwards applied-statistics textbook that covers some of these special topics. Also see Brownlee (1965) and McLean (1966, 1967).

#### 19.9 THE ANALYSIS OF COVARIANCE

Fisher developed the analysis of variance and covariance as a procedure for analyzing the results from factorial designs of many types. Most reluctantly, we have not discussed the analysis of covariance (ANCOVA) in this textbook because to have done so systematically and thoroughly would have required many pages and made an already-long book appreciably longer. For experimentation, the basic principle of ANCOVA is that there are measures of one or more antecedent variables, i.e., measures secured before the random assignment of experimental units to treatments is made. These are chosen with the hope that the regression of the outcome measures on these antecedent measures will be considerable (i.e., that the linear association of the X's, the premeasures, with the Y's, the postmeasures, will be appreciable).

In effect, an analysis of variance is performed on the (Y - Y)'s, where the Y's are predicted from the X's in the usual  $b_1X + b_0$  way described earlier in this book. Some complexity arises because, even with only a single antecedent measure, one has at least two levels of the treatment factor, and hence at least two columns of Y's. With two or more antecedent measures multiple-regression methods must be used, because one is securing the bestweighted composite of the several antecedent variables for predicting the Y's.

A hypothetical illustration of use of the analysis of covariance may make its purpose clearer. Suppose that one is studying four different ways to teach computation of one-way ANOVAs. As his antecedent variables the experimenter has, for each person later to be used in the experiment, a verbalaptitude score (V), a quantitative-aptitude score (Q), and previous gradepoint average (G) in quantitative courses. He assigns  $n_i$  (preferably n =N/4) of the persons randomly to each of the four teaching methods, without any reference to their test scores or GPAs. Then he carries out the experiment and at its end administers to all persons a test of computing one-way ANOVAs; this yields the outcome measures, Y's. Finally, he performs an analysis of covariance on the data to determine whether, after he has used the antecedent information statistically, the adjusted means of the four methods differ significantly. If the regression of the final-test scores on the three predictors is significant, his adjusted within-method mean square will be smaller than if he had not used this antecedent information, thus giving him a more powerful significance test and permitting smaller confidence limits to be constructed around differences between method means. The data layout is sketched in Table 193

For details of such analyses (especially of the case where there is just one predictor variable) see Edwards (1968), Brownlee (1965, pp. 376-96), Winer (1962), McNemar (1962), Lindquist (1953), and Scheffé (1959). Also, for a

fair dice and independent tosses—of 15 straight "passes" in craps is irrelevant in a crap game with loaded dice. Similarly, the probability of a type I error may be substantially incorrect if the data of the experiment do not conform to the statistical-mathematical model from which that probability was derived. We are here concerned primarily with the assumption of independence of the replications of a comparative experiment.

#### Illustration of the Invalidating Effect of Nonindependence of Replications of an Experiment


Imagine that methods A and B of teaching students how to balance chemical equations are being compared. The design for comparing the two methods involves the pairing of 120 pupils into 60 pairs of similar ability and the random assignment of the members of each pair to either method A or method B. All 120 pupils are taught by the same teacher, but they must be divided into four groups of 30 each for instruction. The design can be depicted as shown in Fig. 19.1. Each of the four squares in Fig. 19.1 encloses 30 pupils who study together in a single class.

Suppose that methods A and B will be compared by observing which member of each matched pair (the A pupil or the B pupil) scores higher on a test of proficiency in balancing chemical equations. This is a legitimate, though not very powerful, method of comparing A and B and making a hypothesis test; you may recognize it as the method used in the nonparametric "sign test."

Now let us imagine that a researcher has carried out the above experiment and that in each of the 60 pairs of pupils, the pupil studying under method B scored higher on the proficiency test than his matched partner who studied under method A. (Of course this is an improbable set of results even if method B was far superior to method A. We use this exaggerated example because it portrays more clearly the point we wish to make.) The researcher reasons as follows:

"Suppose that the null hypothesis that methods A and B were equally effective is, in fact, true. What, then, is the probability that pupil B will score higher than pupil A in each of the 60 pairs? If in fact method A and method B are no different, then the probability is 1/2 in any single pair that pupil B will score higher than pupil A (Random assignment of pupils in a pair to methods A and B assures us of this.) The probability that chance alone conspired to make pupil B better than pupil A in all 60 pairs is equal to(1/2)", which is an incredibly small number. Hence, we can safely reject the hypothesis that method A and method B are equally effective."


Implicit in the researcher's calculations which caused him to reject the null hypothesis because of its lack of plausibility in view of the obtained data was the hypothesis that under a true null hypothesis each matched pair



represents an independent opportunity for the occurrence of an event (namely, that pupil B exceeds pupil A) that has probability 1/2. Actually, each matched pair is an experiment: one pupil is taught by method A; the other, by method B. One pair of pupils is an unreplicated experiment. Observation of the relative superiority of methods A and B on several pairs of pupils constitutes replication of the experiment. There are as many replications of the experiment as there are matched pairs whose performance is observed. Our researcher made the assumption—gratuious, perhaps—that the replications of his experiment were independent.

Let us examine this assumption by digging beneath the outward appearances of the experiment—the obtained data—down to the dynamics of the experiment that produced the data. Is it plausible that if pupil B scored higher than pupil A in pair I that this does not affect the chances of pupil B scoring higher than pupil A in pair 2? As most classrooms are now constituted, it does not seem plausible. Recall that the 120 pupils were placed into four separate and intact classrooms. Anyone who knows anything about instruction knows that the members of an intact class interact during instruction in a way that enhances or interferes with the learning in the group.

Again, let us exaggerate so as to make the point more clearly. In one sense we will exaggerate by describing a sort of nonindependence of replications that is far stronger than any that would be met in practice. In another sense, we will exaggerate, or "fantasize" might be a better word, by assuming that the researcher knows how the nonindependence operates, which he seldom will know. Suppose that in the group of 120 pupils there are two troublemakers who are so obnoxious and disruptive that they will successfully inhibit the instruction in any class of which they are a part. These two bait their teachers, annoy their classmates, and generally cause a ruckus. If a method A classroom has one of these troublemakers and the matched method B classroom does not, he will so depress the scores on the final proficiency test that each of his classmates in method A will perform more



pairs and B superior to A for the Data will show method A superior to remaining 30 pairs. method B in all 60 pairs of pupils.

#### FIG. 19.2

poorly than their matched pair in method B. (Of course, if this troublemaker was assigned to method B instead, he would cause a similar appearance of

To guard against the possibility that both of these troublemakers should happen to be assigned to the same classroom (Heaven forbid!), our experimenter made certain that one of them would be assigned at random to either classroom I or II and the other to either classroom III or IV. In any event, the assignment of the pupils in each pair to either method A or B was made at random so that the comparison was not intentionally biased.

In light of what we now know about the activities in the classrooms that produced the data, let us calculate the probability of the obtained results-60 points" for method B and none for method A-assuming that the null hypothesis of no difference is true. In the random assignment of the single troublemaker to either method A or method B for classrooms I and II, there

is probability 1/2 that he will be assigned to classroom I (and cause its 30 pupils to do more poorly than their 30 matched pairs in classroom II) and a probability of 1/2 that he will be assigned to classroom II. The same considerations apply to the assignment of the second troublemaker to either classroom III or classroom IV. The random assignments of the two troublemakers are independent. Hence there are four possible outcomes of the experiment (see Fig. 19.2). (Recall that the presence of a troublemaker in a class will cause the 30 members of that class to do more poorly than their matched partners in the class receiving the other treatment.)

Each of the four outcomes is equally likely. The probability of any one of them is 1/4. In particular, outcome 1 will occur one-fourth of the time this same experiment is executed. Consequently, given that methods A and B are equally effective, the probability is not (1/2)<sup>66</sup> that the B pupil will perform better than the A pupil in all 60 matched pairs (as our researcher originally calculated); the probability of this overwhelming apparent superiority of method B is 1/4 when, in fact, A and B are equally effective. The probability of falsely concluding that method B was superior to method A was 1/4 instead of (1/2)<sup>66</sup>. In fact, the experiment was no more sensitive to the discovery of superiority of one method than an experiment in which only two independent matched pairs of punils were used.

The researcher was so completely in error in his first analysis of the probability of a false conclusion because he failed to recognize that conducting his experiment with intact classrooms did not produce 60 independent replications of his experiment.

### Experimental Unit Versus Unit of Statistical Analysis

In the analysis of experiments a distinction must be made between the unit of statistical analysis and the experimental unit. Before valid probability statements can be made about types of errors, these two units must coincide, i.e., the statistical analysis must be carried out on the legitimate experimental units.

Definition: The units of statistical analysis are the data (the actual numbers) that we consider to be the outcomes of independent replications of our experiment. If you will, the units of statistical analysis are the numbers that we count when we count up degrees of freedom "within" or "for replications."

Imagine that 10 pupils study method I and ten pupils study method II.

A t-test is performed on the 20 scores on the dependent variable; hence, the

experiment.

"pupil" is the unit of statistical analysis. For the sake of analysis, each pupil is considered to be a replication of the experiment; method I is replicated 10 times, and method II is replicated 10 times.

By looking at a researcher's statistical analysis, we can easily determine which unit he chose as the unit of statistical analysis.

It is somewhat more difficult to state an adequate definition of the experimental unit.

Definition: The experimental units are the smallest divisions of the collection of experimental subjects that have been randomly assigned to the different conditions in the experiment and that have responded independently of each other for the duration of the

(This definition clearly reflects the fact that it was proposed with experimentation on living organisms in mind-note the word "respond.")

Independence is the crux of the matter; the following remarks are an attempt to illustrate the notion of independent replications. Suppose that the effects of methods A and B are equal when measured on a particular dependent variable. The probability that subject 1 under method A will score higher on the dependent variable than subject 1 under method B is 1/2 when subjects are randomly assigned to methods. Now consider subject 2 under A and subject 2 under B, i.e., the second replication of the experiment. If the probability that subject 2 under method A will score higher than subject 2 under method B is 1/2 regardless of the outcome of the experiment on the first replication (with subjects 1), then the two replications are independent.

Judging the validity of the assumption that the replications of an experiment are independent is no easy matter. The assumption of homogeneous variances is easily tested with Bartlett's test (if we are confident the normality assumption is met) or Levene's test. The assumption of normality is easily tested with a chi-square or Kolmogorov-Smirnov test if nis reasonably large. However, the researcher will usually be faced with the task of making a considered judgment of the degree of independence of the replications rather than the task of applying a particular statistical test. His judgment must be based on an intimate knowledge of the dynamics of the experimental setting. In some instances, blatant nonindependence of the replications will be impossible to overlook, as in our example. In other experiments, nonindependence of replications will be subtle and can go

In the example at the beginning of this section, the experimental subjects were the 120 pupils. The smallest division of this group of subjects that was randomly assigned to methods A and B was an individual pupil. However, the individual pupils did not respond to instruction independently. In two classrooms, a troublemaker caused the remaining pupils to learn nothing about balancing chemical equations.

Each classroom did respond independently of the other three classrooms, we might assume; and the classrooms were randomly assigned to methods A and B. Therefore, the "classroom" is the smallest division of the 120 pupils that satisfies the conditions of random assignment and independent responding. Therefore, the "classroom" is the experimental unit in our example.

À valid analysis of the illustrative experiment would be carried out on four observations: the average proficiency test scores of the four classrooms. In other words, a valid analysis would use the four classroom means as the units of statistical analysis. The experiment comparing methods A and B was only replicated twice (once for each classroom) for each method. "Classroom" is the experimental unit; hence, the classroom means must be the unit of statistical analysis. A valid statistical inferential analysis is possible, but it is hardly worth the bother. The results of the two replications of the experiment give us no more evidence to conclude that method B is better than method A, than we have evidence to conclude that a coin is biased in favor of "heads" because the first two times we flipped it we got "heads." (However, means based on 30 pupils each should be more stable than means based on one pupil each, i.e., the experimental results should be better for the 120 pupils than they would have been for just four, one in each class.)

Educational researchers are especially prone to making the error of analyzing data in terms of units other than the legitimate experimental unit. Almost all of the comparative experiments carried out under actual school conditions face the same dilemma. At most, no more than five or six intact classrooms have been involved in the experiment. Perhaps pupils have been assigned to classrooms at random, perhaps not. At least, the classrooms have been assigned at random to the experimental conditions being compared. The researcher has two alternatives, though he is seldom aware of the second one: (1) he can run a potentially illegitimate analysis of the experiment by using the "pupil" as the unit of statistical analysis, or (2) he can run a legitimate analysis on the means of the five or six classrooms, "classroom" being the actual experimental unit, in which case he is almost certain to obtain statistically nonsignificant results (with only five or six replications, the power of his significance test is low).

If the researcher chooses the first alternative and is led into error, he can find solace in the knowledge that methodologists themselves have long sanctioned his actions either explicitly or by example.

As early as 1940, Lindquist presented a legitimate analysis of variance of data gathered from an experiment involving intact groups (see Lindquist, 1940, pp. 107ff.). Lindquist's thinking was far ahead of that of his colleagues,

if McNemar was representative of the latter. In a review of Lindquist's text McNemar (1940) wrote as follows:

We next raise a puzzling question for which we have no definite answer. Regioning on page 107, the analysis of variance technique is applied to an educational-methods experiment involving five schools and three methods, with twenty nunits in each of lifteen classes. The analysis is carried through on the basis of the fifteen class means in such a way that neither the number of pupils nor the pupil variation enters into the analysis. This at first struck the reviewer as being indefensible, but a few pages later (p. 117) one finds that the pupils have their inning via the 'within classes' variation, but this latter variation is not utilized because the 'interaction variance' is larger than the 'within classes' variance. The reviewer suspects that something is wrong with a test of significance which does not involve the variation of the individuals upon which the means are based. We are unable to locate the fallacy here, if there be such, but we have in mind a worked-out example in sex differences which shows no significant difference in length at birth when analyzed by the variance technique using means, but which yields highly significant differences when analyzed by the ordinary critical ratio procedure. We are not arguing that the author is wrong in arguing that intact groups are the proper sampling units, but rather that the case is not convincingly stated.

Notice that McNemar's worked-out example, which he considered to be contrary to Lindquist's example, is quite unlike a classroom experiment of the sort Lindquist discussed. What if McNemar's example involved 20 boys (four from each of five families)?

#### Further Reading

Fortunately, there now exist about a half-dozen discussions of the problem of determining the appropriate experimental unit and unit of statistical analysis:

- Campbell, D. T. and J. C. Stanley. "Experimental and quasiexperimental designs in research on teaching." Chapter 5 in Handbook of Research on Teaching, ed. N. L. Gage. Chicago: Rand-McNally, 1963. Relevant pages: 192, passim.
- 2. Cos, David R. Planning of Experiments. New York: Wiley, 1958. Relevant pages; 2-3, 155, 171-73, 191, 196. The designated pages deal with the problem of experimenting with intact groups. The context is agricultural experimentation and the discussion is at a high level. Save this one to read after you have mastered some preliminary material.
- Hovland, C. I., A. A. Lumsdaine, and F. D. Sheffield. Experiments on Mass Communication. Princeton, N.J.: Princeton University Press, 1949.

- 4. Lindquist, E. F. Design and Analysis of Experiments in Psychology and Education. Boston: Houghton Mifflin, 1953. Relevant pages: 192-93. A short discussion of experimenting with intact class-tooms. Lindquist discusses the problem in an educational context. An important reference. However, it must be read carefully and critically. When Lindquist prepared this textbook, the procedures we presented for finding E(MS)'s had not yet been worked out fully.
- 5. Lumsdaine, A. A. "Instruments and media of instruction." Chapter 12 in Handbook of Research on Teaching, ed. N. L. Gage. Chicago: Rand-McNally, 1963. See especially pages 656-59. Pages 656-59 are probably the best readily available discussion of this crucial problem of running comparative experiments with intact groups of persons instead of individual persons.
- Peckham, Perc D., Gene V. Glass, and Kenneth D. Hopkins. "The experimental unit in statistical analysis: comparative experiments with intact groups." *Journal of Special Education*, 3 (1969).

### **TABLES**

### TABLE A SOOO RANDOM DIGITS

36 55 59 55 54 17 54 67 37 04 32 64 35 28 61 69 57 26 87 77 24 12 26 65 91	93 71 61 68 94 66 08 32 46 53 32 88 65 97 80 08 35 56 08 60 92 05 24 62 15 55 12 12 92 81 95 81 90 68 31 00 91 19 89 36 39 51 03 59 05 14 06 04 06 19 27 69 90 64 94 14 84 54 66 59	29 73 54 77 62 71 29 92 38 59 07 60 79 36 27 95 45 8 76 35 59 37 79 80 86 30 05 29 54 96 96 16 33 56 46 07	53 09 14
61 19 63 02 31 30 53 22 17 04 03 78 89 75 99	27 69 90 64 94 14 84 54 66 72 92 96 26 17 73 41 83 95 53 82 10 27 41 22 02 39 68 52 33 07 75 86 72 07 17 74 41 65 31 66 85 78 34 76 19 53 15 26 74 33	17 26 77 09 43 78 03 87 02 10 06 16 88 29 55 98 66 64	67 85 23

Regranted from A Million Random Digits With 100,000 Normal Deciates (New York: The Free Press, 1955), by permassion of the RAND Corporation and the publisher.

_																											
A 60	36	59	46	53	35	07	5	3 3	9 49	- 4	2 6	51 4	12 9	92	97	0	1 9	91	82	83	16	-	98	95	37	32	31
83	79	94	24	02					4 42			19 4							11								65
32	96	00	74	05			98					8 5			00				30								53
19	32	25	38	45					6 06			9 7							86								13
11	22	09	47	47					\$ 08			0 9							24						43		
31	75	15	72	60	68	98	ο	) 5	3 39	1	5 4	7 0	4 8	33	55	8	8 6	5 :	12	25	96	c	13	15	21	91	21
88	49	29	93	82					5 04			94				7.	4 8	4 3	39	34	13	2	2	10	97	85	08
30	93	44	77	41			18					8 8				2	3 5	9 :	72	04	05	9	4	20	52	03	80
22	88	84	88	93					48			30				74	1 0	2 2	28	46	17	8	2	03	71	02	68
78	21	21	69	93					86			7 2				6.	5 7	4 1	11 -	40	14	8	7 .	48	13	72	20
41	84	98	45	47	46	85	05	23	26			7 7							6 .						58		
46	35	23	30	49	69	24	89	34	60	4:	5 30	0 5	0 7	5	21				33						37		
11	08	79	62	94	14	01	33	17	92	59	7.	4 76	67	2	77				3 4						37		
52	70	10	83	37	56	30	38	73	15			2 00							9 9						16		
		53			81	30	44	85	85	68	3 6:	5 22	2 7	3	76	92	8:	5 2	5 :	8	66	-			80		
20	85	77	31	56	70	28	42	43	26	79	37	7 59	5	2	20	01	1.	59	6 3	2	67				24		
15	63	38	49	24					14			2 12							4 7						53		
92	69	44	82	97					15	59	58	94	91	0 1	67				4 1						70		
77	61	31	90	19					80	20	55	49	14	4 (	09				48						69 '		
38	68	83	24	86					45			47				43	94	7:	5 1	6 8	80	43	8	5 :	25 !	6 :	93
25	16	30	18	89	70	01	41	50	21	41	29	06	73	3	12		85		1 5						16 :		
65	25	10	76	29	37	23	93	32	95	05	87	00	11	ا ا	19				2 6						19 2		
36	81	54	36	25	18	63	73	75	09	82	. 44	49	90	) (	)5				7 3						2 2		
04	39	71	16	92	05	32	78	21	62	20	24	78	17	7 5	59	45	19	1/2	2 5	3 3 7 0	12				3 3		
		52			95	09	66	79	46			08							8								
83	76	16	08	73	43	25	38	41	45	60	83	32	59	8	13				1						07		
14	38	70	63	45			40			43	52	90	63	1	8	38	38	47	4	7 6	1				31		
21	32	19	22	45	80	08	87	70	74	88	72	25	67	3	6	66	16	44	94	3	1				54		
72	47	20	00	08	80	89	10	80	02	94	81	33	19	0	ю	54	15	58	34		4				68		
		65			93	12	81	84	64	74	45	79	05	6	1				18								
39	52	87	24	R4	82	47	42	55	03	48	54	53	52	4	7	18	61	91	36	7.	4	18	61	1	1 9	4	1
81	61	61	87	11			24			75	12	21	17	2	4	74	62	77	37	0		58	31	9.	5 79	, ,	,
07	58	61	61	20			12			92	90	41	31	4	1	32	39	21	97	6					08		
90	76	70	42	25			41		00	69	90	26	37	4	2	78	46	42	25	01					89		
40	18	82	81	93			38			94	97	21	15	9	8	62											
34	41	48	21	57	06	00	75	50	07	10	15	20	00	2	3	12	30	28	07	83					86		
03	43	97	52	62	86 44					36	02	40	08	6	7	76	37	84	16	05		65	96	17	34	- 82	3
67	04	90	on.	70	93					94	45	87	42	84	1	05	04	14	98	07		20	28	83	40	D(	,
79	49	50	41	10	52					54	15	83	42	43	3	46	97	83	54	82		9	36 30	29	59 34	70	
91	70	43	05	52	04					75	05	19	30	29	•	47	66	56	43	82							
09	10	82								04		00	c2	0.4		83	38	98	73	74		54	27	85	80	44	
90	04	58	00	97	32					94	02	00	19	97	,	91	87	07	61	50		8	47	66	46	59	
73	18	95	24 02	97 07	51					94 62	20	06	44	64		27	12	46	70	18	- 4	11	36	18	27	50	
13	76	87	64	00	47					90	42	91	22	72		95 3	37	50	58	71	9	3 1	52	34	31 78	18	
54	01	64	40	5U 54	20					00	68	22				20	71 -	45	32	95	(	7	ľŪ	61	18	13	
	-	-4	70	20	66	28	13	10	U.S	vu	u																

	78 54 24		13 66 15			89 75 :	
	81 33 31		40 51 00			46 04 1	
53 84 08 62 33	81 59 41		51 21 59			66 87 9	
91 75 75 37 41	61 61 36		50 26 39			17 65 1	
89 41 59 26 94	00 39 75	83 91	12 60 71	76 46	48 94	97 23 (	06 94 54 13 74 08
77 51 30 38 20	86 83 42		68 41 48			8I 39 8	
19 50 23 71 74	69 97 92		55 21 02	2 97 73	74 28	77 52 5	51 65 34 46 74 15
21 81 85 93 13	93 27 88		05 68 67		07 08	28 50 4	46 31 85 33 84 52
51 47 46 64 99	68 10 72		94 04 99	9 13 45	42 83	60 91 9	91 08 00 74 54 49
99 55 96 83 31	62 53 52	41 70	69 77 7	1 28 30	74 81	97 81 4	42 43 86 07 28 34
33 71 34 80 07	93 58 47	28 69	51 92 6	6 47 21	58 30	32 98 2	22 93 17 49 39 72
85 27 48 68 93	11 30 32		28 83 43			59 04 6	
84 13 38 96 40	44 03 55		73 85 2	7 00 91		26 05	
56 73 21 62 34	17 39 59		10 12 3	9 16 22	85 49	65 75	60 81 60 41 88 80
65 13 85 68 06	87 64 88	3 52 61	34 31 3	6 58 61	45 87	52 10	69 85 64 44 72 77
38 00 10 21 76	81 71 91	1 17 11	71 60 2	0 20 22	74 21		
37 40 29 63 97	01 30 4		56 27 1		14 21	96 40 4	
97 12 54 03 48	87 08 3	3 14 17	21 81 5		75 32	46 26 0 76 20 4	
21 82 64 11 34	47 14 3		64 63 8		73 Z3	90 64 4	
73 13 54 27 42	95 71 9	0 90 35	85 79 4	7 42 06	09 70	98 81 3	
					00 /8	98 81 .	56 64 69 11 92 02
07 63 87 79 29 60 52 88 34 41	03 06 1	1 80 72	96 20 7	4 41 56	23 82	19 95 ;	38 04 71 36 69 94
83 59 63 56 55	07 95 4		59 17 5	2 06 95	05 53	35 21	39 61 21 20 64 55
10 85 06 27 46	06 95 8		05 12 8	0 97 19	77 43	35 37	83 92 30 15 04 98
39 82 09 89 52	99 59 9	1 05 07	13 49 9	0 63 19	53 07	57 18	39 06 41 01 93 62
37 82 07 89 32	43 62 2	6 31 47	64 42 1	8 08 14	43 80	00 93	51 31 02 47 31 67
59 58 00 64 78	75 56 9	7 88 00	22 01 e	5 44 86			
38 50 80 73 41	23 79 3	4 87 63	00 00 0	9 70 22	23 76	80 61 :	56 04 11 10 84 08
30 69 27 06 68	94 68 8	1 61 27	56 10 4	8 00 91	17 71	90 42 (	07 95 95 44 99 57
65 44 39 56 59	18 28 8	2 74 37	49 63 2	2 40 41	82 06	76 34 6	00 05 46 26 02 00
27 26 75 02 64	13 19 2	7 22 94	07 47 7	4 46 06	08 33	76 56	
91 30 70 69 91					17 98	54 89	11 97 34 13 03 58
68 43 49 46 88	19 07 2	22 42 10	36 69 9	5 37 28	20 02	53 57 9	
48 90 81 58 77		1 36 22	62 12 6	9 84 02	17 04	38 25	
06 91 34 51 97		52 45 91	35 70 C	00 47 54	83 63	45 26	
10 45 51 60 19		27 86 01	11 88 3	0 95 72	63 01	19 89 (	
		3 37 12	91 34 2	78 21	88 32	58 08	
12 88 39 73 43	65 02	76 11 84					
21 77 83 09 76	38 80 '	73 69 61	11 (4	0 13 92	17 97	41 50	77 90 71 22 67 69
19 52 35 95 15	65 12 :	25 96 50	96 00	4 20 96	63 28	10 20 3	71 09 91 64 74 40
67 24 55 26 70	35 58	31 65 62	20 24	36 82 58	69 57	21 37 4	98 16 42 60 16 00
60 58 44 73 77	07 50	03 79 92		58 66 86 12 65 29	76 46	33 42 -	22 26 65 60 00 02
53 85 34 13 77				4 63 29	26 76	08 36	37 41 32 64 43 44
24 63 73 87 36		69 48 50		37 38 59			
83 03 01 24 51	11 00	48 93 42	52 62 3	30 70 pr	49 36	47 33 3	
16 44 42 43 14	36 14	22 28 15	07 75 9	25 17 27	12 36	91 86 (	01 03 74 28 38 73
60 79 01 81 57		19 90 73 86 57 62	27 49 1	37 na na	95 13	72 75 1	85 51 97 23 78 67
		oo oo 62	11 16 1	17 85 76	45 01	03 25	52 54 84 65 47 59
					-2 81	95 29	79 65 13 00 48 60

TABLE B AREAS AND ORDINATES OF THE UNIT NORMAL DISTRIBUTION



z	Area	u Ordinate	z	Area	Ordinate
-3.00	.0013	,0044			
-5.00	100.12			.0036	.0107
-2.99	.0014	,0046	-2 69	.0037	.0110
-2.98	.0014	.0047	-2.68	.0038	.0113
-2.97	.0015	.0048	-2.67	.0039	.0116
-2.96	.0015	.0050	-2.66	,0040	.0119
-2.95 -2.95	.0016	.0051	-2.65	,0040	.0117
-2.93	.00.0			.0041	.0122
-2.94	.0016	.0053	-2.64	.0043	.0126
-2.94	,0017	.0055	-2.63	.0044	.0129
-2.93	.0018	.0056	-2.62	.0044	.0132
-2.92	.0018	,0058	-2.61	.0047	.0136
2.90	.0019	.0060	-2.60	.00-17	
2.90	,0017		2.50	.0048	.0139
-2.89	.0019	.0061	-2.59	.0049	.0143
-2.89 -2.88	.0020	.0063	2.58	.0051	.0147
-2.87	.0021	.0065	-2.57	0052	.0151
2.86	.0021	.0067	-2.56	.0054	.0154
-2.85	.0022	.0069	-2.55	,000,4	
-2.83	.0022		-2.54	.0055	.0158
-2.84	.0023	.0071	-2.53	.0057	.0163
-2.83	.0023	.0073	-2.52	.0059	.0167
-2.83 -2.82	.0024	.0075	-2.52 -2.51	.0060	.0171
-2.81	.0025	.0077	-2.50	.0062	.0175
2.80	.0026	.0079	-2.50		
2-2.80			-2.49	.0064	.0180
-2.79	.0026	.0081	-2.48	.0066	.0184
-2.78	.0027	,0084 .0086	-2.47	,0068	.0189
-2.77	.0028		-2,46	,0069	.0194
-2.76	.0029	.0088	-2,45	.0071	.0198
-2.75	.0030	.0091			
2		.0093	-2.44	,0073	.0203
2.74	.0031	.0096	-2.43	,0075	
-2.73	.0032	,0099	-2.42	.0078	.0213
-2.72	.0033	.0101	-2.41	.0080	.0219
-2.71	.0034	.0104	-2.40	.0082	.0224
-2.70	.0035		<u> </u>		

				_																				
08 3	35 1	36 9	99	10	78	54	24	27	85	13	66	15	88	73	04	61	89	75	53	31	22	30	84	20
28 3	10	SO :	32	64	81	33	31	05	91	40	51	00	78	93	32	60	46	04	75				18	
53 8	34 (	38 (	62	33	81	59	41	36	28	51	21	59	02	90	28	46	66	87	95				07	
91 1					61	61	36	22	69	50	26	39	02	12	55	78	17	65	14	83	48	34	70	55
89	11	59	26	94	00	39	75	83	91	12	60	71	76	46					06				74	
77							42			68	41	48	27	74	51	90	81	39	80	72	89	35	55	07
19	50	23	71	74			92			55	21	02	97	73					51				74	
21					93	27	88	17	57	05	68	67	31	56					46				84	
51							72			94	04	99	13	45					91				54	
99	55	96	83	31	62	53	52	41	70	69	77	71	28	30	74	81	97	81	42				28	
33					93	58	47	28	69	51	92	66	47	21	58	30	27	0.0	22	02		40	39	
85							32					43							00				36	
84							55					27			61	22	26	05	61	(1	22	71	84	43
	73						59					39			85	49	65	75	60	02	52	41	84	23
65	13	85	68	06	87	64	88	52	61			36			45	87	52	10	69	81 85	64	41	88 72	77
38	00	10	21	76	81	71	91	17	11	71	٤0	29	20											
37	40	29	63	97	01	30	47	75	86	56	27	11	29	31	14	21	96	40	49				96	
97	12	54	03	48	87	08	33	14	17	71	91	53	01	50	4/	32	46	26	05	40	03	03	74	38
21	82	64	11	34	47	14	33	40	72	6.1	67	88	92	20	/3	23	76	20	47	15	50	12	95	78
73	13	54	27	42	95	71	90	90	35	84	70	47	29	02	49	13	90	64	41	03	85	65	45	52
															08	78	98	81	56	64	69	11	92	02
				29 41	03	06	11	80	72	96	20	74	41	56	23	82	19	95	38	04	71	20	69	
91	50	61	54	55	07	95	41	98	14	59	17	52	06	95	05	53	35	21	39	61	71	20	64	94
10	37	03	20	46	06	95	89	29	83	05	12	80	97	19	77	43	35	37	83	01	21	20	04	33
				52	99	39	91	05	07	13	49	90	63	19	53	07	57	19	39	22	30	13	93	98
	-	٠,	0,		43	02	26	31	47	64	42	18	08	14	43	80	00	93	51	31	02	47	31	62
				78	75	56	97	88	00	90	01	55								21	02	*/	31	0/
38	50	80	73	41	23	79	34	87	63	60	0.7	29	44	86	23	76	80	61	56	04	11	10	84	លន
30	69	27	06	68	94	68	81	61	27	30	10	68	/0	22	17	71	90	42	07	95	95	44	99	53
65	44	39	56	5 59	18	28	82	74	37	40	61	22	00	91	82	06	76	34	00	05	46	26	92	00
27	26	75	0	2 64	13	19	27	22	94	07	47	74	40	41	08	33	76	56	76	96	29	99	08	36
91	30	70		9 91											17	98	54	89	11	97	34	13	03	58
68	43	49	4	6 88	15	47	22	42	10		69	95	37	28	28	27	53							
45	90	1 81	- 51	8 77		4/	31	36	22	62	12	69	84	Uδ	12	94	38	3/	93	28	97	66	62	52
O.	5 91	34	5	1 97		14	32	45	91	35	70	00	47	54	83	27	45	23	90	09	81	59	31	46
10	45	51	64	19		31	21	86	10	- 11	88	30	95	79	63	01	19	20	92	54	13	os	51	60
					-	- 21	U	31	12	91	34	23	78	21	88	32	58	69	01	14	97	44	03	44
13	2 88	39	7	3 43	65	i n	, 74		84							-	30	08	21	43	66	77	08	83
2	7	7 8	0	2 76	20	80	7	66	61		28	50	13	92	17	97	41	50	27	-00	٠.			
11	9 5	2 3	5 9	5 15		5 12	2 2	0.	59		64	94	20	96	63	28	10	20	22	90	/1	22	67	69
6	1 2	1 5	5 2	6 70	3:	5 51	8 31	1 69	63	70	28	36	82	58	69	57	21	37	0.5	16	٥l	04	74	49
6	U 5	5 4	7	3 77	0	7 50	0 0	70	92		24	68	66	86	76	46	33	42	22	10	45	39	15	29
	١.	٠.		3 77							13	42	65	29	26	76	08	36	37	41	03	39	08	υZ
,		, ,	: !	3 77 17 36		6 0	6 6	49	50	49		87								41	32	04	43	44
	, p	2 0		4 51	7.	4 3:	3 4	8 0	, ,,		63	30	38	59	49	36	47	33	31	96	24	0.1	36	
ĭ	6 4		; ;	3 34	- 3	8 9	2:	2 20			75	30 95	19	92	12	36	91	28	01	03	74	28	38	72
6	0 7	9 n		1 57	ر ن	0 13	5 19	9 00	77		40	37	17	77	97	37	72	75	25	51	97	23	78	67
			_ •	,	- 5	1 1	7 8	5 57	62		16	17	09	39	85	13	03	25	52	54	84	65	78 47	50
					_		-	_	_				83	76	45	81	95	29	79	65	13	m	48	60
												_	_	_	_	_						~~	70	w

					.7
		и		1 .	u
z	Area	Ordinate	Į z	Area	Ordinate
-1.49	.0681	.1315	-1.04	.1492	.2323
-1.48	.0694	.1334	1.03	.1515	.2347
-1.47	.0708	.1354	-1.02	.1539	.2371
-1.46	.0721	.1374	-1.01	.1562	.2396
		.1394	-1.00	.1587	.2420
-1.45	.0735	.1394	-1.00	.1307	.2420
-1.44	.0749	.1415	-0 99	.1611	,2444
-1.43	.0764	.1435	-0.98	.1635	.2468
-1.42	.0778	.1456	-0.97	.1660	.2492
-1.41	.0793	.1476	-0.96	.1685	.2516
-1.40	.0808	.1497	-0.95	.1711	.2541
			-0.94	.1736	.2565
-1.39	.0823	.1518	-0.94	.1762	.2589
-1.38	.0838	.1539		.1788	.2613
-1.37	.0853	.1561	-0.92	.1814	.2637
-1.36	.0869	.1582	-0.91	.1841	.2661
-1.35	.0885	.1604	-0.90	.1041	.2001
-1.34	.0901	.1626	-0.89	.1867	.2685
-1.33	.0918	.1647	-0.88	.1894	.2709
-1.32	.0934	.1669	-0.87	.1922	.2732
÷1.32	.0934	.1691	-0.86	.1949	.2756
-1.30		.1714	-0.85	.1977	,2780
-1.30	.0968	.1714		,2005	.2803
1.29	.0985	.1736	-0.84		.2827
-1.28	.1003	.1758	-0.83	.2033	.2850
-1.27	.1020	.1781	-0.82		.2874
-1.26	.1038	.1804	-0.81	.2090	.2897
-1.25	.1056	.1826	-0.80	.2119	
-1.24	.1075	.1849	-0.79	.2148	.2920
-1.24 -1.23	.1073	.1872	-0.78	.2177	.2943
-1.23 -1.22		.1895	-0.77	.2206	.2966
	.1112	.1919	0.76	,2236	.2989
-1.21 -1.20	.1131 .1151	.1942	-0.75	.2266	.3011
		1055	-0.74	.2296	.3034
-1.19	.1170	.1965	-0.73	,2327	.3056
-1.18	.1190	.1989	-0.72	.2358	.3079
-1.17	.1210	.2012	-0.71	.2389	.3101
-1.16	.1230	.2036	-0.70	.2420	.3123
1.15	.1251	.2059			,3144
-1.14	,1271	,2083	-0.69	.2451	.3166
-1.14 -1.13	.1271	.2107	-0.68	.2483	.3187
-1.13	.1314	.2131	-0.67		.3209
-1.12 -1.11	.1335	,2155	-0 66	.2546 .2578	,3230
-1.11 -1.10	.1357	.2179	-0.65	.2318	
		.2203	0.64	.2611	.3251 .3271
1.09	,1379	.2203	0.63	.2643	.3292
-1.08	.1401	.2251	-0.62	.2676	,3292
-1.07	.1423	.2275	-0.61	,2709	.3312
-1.06	.1446	.2273	0.60	.2743	.5
-1.05	.1469	.2299			

z	Area	u Ordinate	z	Area	u Ordinate
-2.39	,0084	.0229	-1.94	.0262	.0608
-2.38	.0037	.0235	-1.93	.0268	
-2.37	.0039	.0241	-1.92	.0274	.0620
-2.36	.0091	.0246	-1.91	.02/4	.0632
-2.35	.0094	.0252	~1.90		.0644
		.0232	~1.90	.0287	.0656
-2.34 -2.33	.0096	.0258	-1.89	.0294	.0669
-2.33	.0099	.0264	-1.88	.0301	.0681
-2.32	.0102	.0270	-1.87	.0307	.0694
	.0104	.0277	-1.86	.0314	.0707
-2.30	.0107	.0283	-1.85	.0322	.0721
-2,29	.0110	.0290	-1.84	.0329	
-2.28	.0113	.0297	-1.83		.0734
~2.27	.0116	.0303	-1.82	.0336	.0748
-2.26	.0119	.0310	-1.81	.0344	.0761
~2.25	.0122	.0317		.0351	.0775
~2.24		.0317	-1.80	.0359	.0790
-2.24	.0125	.0325	-1,79	.0367	
-2.23 -2.22	.0129	.0332	-1.78	.0375	.0804
-2.22 -2.21	.0132	.0339	-1.77	.0373	.0818
-2.21 -2.20	.0136	.0347	~1.76	.0392	.0833
-2.20	.0139	.0355	-1.75	.0392	.0848
-2.19	.0143	.0363	f .		.0003
-2.18	.0146	.0371	-1.74	.0409	.0878
-2.17	.0150	.0371	-1.73	.0418	.0891
~2.16	.0154	.0379	-1.72	.0427	.0909
-2.15	.0158	.0396	-1.71	.0436	.0925
• • •		.0396	~1.70	.0446	.0940
-2.14 -2.13	.0162	.0404	~1.69		
-2.13	.0166	.0413	-1.68	.0455	.0957
-2.12	.0170	.0422	-1.67	.0465	.0973
-2.11	.0174	.043[	-1.66	.0475	.0989
-2.10	.0179	.0440	-1.65	.0485 .0495	.1006
-2.09	.0183	.0449	i i	.04,55	.1023
-2.08	.0188	.0459	-1.64	.0505	.1040
-2.07	.0192	.0468	-1.63	.0516	.1057
-2.06	.0197	.0478	~1.62	.0526	.1074
-2.05	.0202	.0488	-1.61	.0537	.1092
-201	.0207		-160	.0548	-1109
-201	.0212	,0498	-1.59		
-202	.0212	.0503	-1.58	.0559	.1127
-201	.0222	.0519	-1.57	.0571	.1145
-200	.0228	.0529	-1.56	.0582	.1163
		.0540	-1.55	.0594	.1182
-1.99 -1.98	.0233	.0551			.1200
-1.93	.0239	.0562	~1.54	.0618	.1219
-1.96	.0244	.0571	-1.53	.0630	.1238
-1.95	.0250	.0584	-1.52	.0643	.1257
,	.0256	.0596	-1.51 -1.50	.0655	.1276

	1 .	u	Jj.	Area	u Ordinate
z	Area	Ordinate	z z	Area	Orainale
0.31	.6217	.3802	0.76	.7764	.2989
0.32	.6255	.3790	0.77	.7794	.2966
0.33	.6293	.3778	0.78	.7823	.2943
0.34	.6331	.3765	0.79	.7852	.2920
0.35	.6368	.3752	0.80	.7881	.2897
				.7910	.2874
0.36	.6406	.3739	18.0	.7939	.2850
0.37	.6443	.3725	0.82	.7967	.2827
0.38	.6480	.3712		.7995	.2803
0.39	.6517	.3697	0.84	.8023	.2780
0.40	.6554	.3683	0.85	.8023	1 2700
0.41	.6591	.3668	0.86	.8051	.2756
0.42	.6628	.3653	0.87	.8078	.2732
0.43	.6664	.3637	0.88	.8106	.2709
0.44	.6700	.3621	0.89	.8133	.2685
0.45	.6736	.3605	0.90	.8159	.2661
			0.91	.8186	.2637
0.46	.6772	.3589	0.92	,8212	.2613
0.47	.6808	.3572	0.93	.8238	.2589
0.48	.6844	.3555	0.94	.8264	.2565
0.49	.6879	.3538	0.95	.8289	.2541
0.50	.6915	.3521	0.93	1	.2516
0.51	.6950	.3503	0.96	.8315	.2492
0.52	.6985	.3485	0.97	.8340	.2492
0.52	.7019	.3467	0.98	.8365	.2444
0.54	.7054	.3448	0.99	.8389	.2420
0.55	.7088	.3429	1.00	.8413	.2420
		1	1.01	.8438	.2396
0.56	.7123	.3410	1.01	,8461	.2371
0.57	.7157	.3391	1.02	.8485	.2347
0.58	.7190	.3372	1.04	.8508	.2323
0.59	.7224	.3352	1.05	.8531	.2299
0.60	.7257	.3332	1.03		,2275
0.61	7701	,3312	1.06	.8554	.2251
0.61	.7291 .7324	.3292	1.07	.8577	2227
0.62 0.63		.3271	1.08	.8599	.2203
0.64	.7357 .7389	.3251	1.09	.8621	.2179
0.64	.7422	.3230	1.10	.8643	-
	1		1.11	.8665	.2155
0.66	.7454	.3209	1.12	.8686	.2131
0.67	.7486	.3187	1.13	.8708	,2107 ,2083
0.68	.7517	,3166	1.14	.8729	.2083
0.69	.7549	.3144	1.15	.8749	.2039
0.70	.7580	.3123		.8770	.2036
0.71	.7611	1018,	1.16	,8790	.2012
0.72	.7642	.3079	1.17	.8810	.1989
0.72	.7673	,3056	1.18	.8830	.1965
0.74	.7704	.3034	1.19 1.20	8849	.1942
0.75	,7734	.3011	1,20		

z	Area	u Ordinate	2	Area	u Ordinate
-0.59	.2776	.3352	-0.14	.4443	.3951
-0.58	.2810	.3372	-0.13	.4483	.3956
-0.57	.2843	.3391	-0.12	.4522	
-0.56	.2877	,3410	-0.11		.3961
-0.55	.2912	.3429		.4562	.3965
0.55	.2712	.3429	-0.10	.4602	.3970
-0.54	.2946	.3448	-0.09	.4641	.3973
-0.53	.2981	.3467	-0.08	.4681	.3977
-0.52	.3015	.3485	-0.07	.4721	
-0.51	.3050	.3503	-0.06	.4761	.3980
-0.50	.3085	.3521	-0.05	.4801	.3982
-0.49	.3121	.3538			.5964
-0.48	.3156		-0.04	.4840	.3986
-0.47	.3192	.3555	-0.03	.4880	,3988
-0.46		.3572	~0.02	-4920	.3989
	.3228	.3589	-0.01	.4960	.3989
0.45	.3264	.3605	0 00	.5000	.3989
-0.44	.3300	.3621	0.01	****	
-0.43	.3336	.3637	0.02	.5040	.3989
-0.42	.3372	.3653	0.02	.5080	.3989
-0.41	.3409	.3668		.5120	.3988
-0,40	.3446	.3683	0.04 0.05	.5160 .5199	.3986
-0.39	.3483			.3199	.3984
-0.38	.3520	.3697	0.06	.5239	.3982
-0.37	.3557	.3712	0.07	.5279	.3980
-0.36	.3594	.3725	0.08	.5319	
-0.35		.3739	0.09	.5359	.3977
	.3632	.3752	0.10	.5398	.3973
-0.34	.3669	.3765			10,770
-0.33	.3707	.3778	0.11	.5438	.3965
-0.32	.3745	.3790	0.12	.5478	.3961
-0.31	.3783	.3802	0.13	.5517	.3956
-0.30	.3821		0.14	.5557	.3951
		.3814	0.15	.5596	.3931
-0.29 -0.28	.3859	.3825	اميدا		1
	.3897	.3836	0.16	.5636	.3939
-0.27	.3936	.3847	0.17	.5675	.3932
-0.26	.3974	.3857	0.18	.5714	.3925
-0.25	.4013	.3867	0.19	.5753	.3918
~0.24	.4052		0.20	.5793	.3910
-0.23	.4090	.3876	0.21	5035	
-0.22	.4129	.3885	0.22	.5832	.3902
-0.21		.3894	0.23	.5871	.3894
~0.20	.4168	.3902	0.23	.5910	.3885
	A207	.3910	0.25	.5948 .5987	.3876
-0.19	.4247			.5,6/	.3867
~0.18	.4286	.3918	0.26	.6026	
-0.17	.4325	.3925	0.27	.6064	.3857
-0.16	4364	.3932	0.28	.6103	.3847
-0.15	.4404	.3939	0.29		.3836
		.3945	0.30	.6141	.3825 .3814

z         Area         Undinate         z         Area         Undinate           2.11         .9826         .0431         2.56         .9948         .0151           2.12         .9830         .0422         2.57         .9948         .0151           2.13         .9834         .0413         2.58         .9951         .0143           2.14         .9838         .0404         2.59         .9952         .0139           2.16         .9846         .0387         2.61         .9955         .0132           2.17         .9850         .0379         2.62         .9936         .0129           2.18         .9854         .0317         2.61         .9955         .0132           2.19         .9857         .0363         2.64         .9959         .0122           2.20         .9861         .0335         2.65         .9960         .0119           2.21         .9864         .0347         2.66         .9961         .0116           2.22         .99868         .0339         2.67         .9962         .0113           2.23         .9875         .0225         2.69         .9964         .0107           2.24										
2.12	z	Area		.	z		Area	,		
2.12         .9830         .0422         2.57         .9949         .0147           2.13         .9834         .0413         2.58         .9951         .0147           2.14         .9838         .0404         2.59         .9952         .0139           2.15         .9846         .0387         2.61         .9955         .0132           2.16         .9846         .0387         2.61         .9955         .0122           2.17         .9850         .0379         2.62         .9956         .0129           2.18         .9854         .0371         2.63         .9957         .0126           2.19         .9857         .0363         2.64         .9959         .0122           2.20         .9861         .0355         2.65         .9960         .0119           2.21         .9864         .0347         2.66         .9961         .0116           2.22         .9863         .0319         2.67         .9962         .0111           2.22         .9865         .0319         2.68         .9961         .0116           2.22         .9875         .0217         2.70         .9966         .0101           2.24 <td>2.11</td> <td>.9826</td> <td>0431</td> <td> </td> <td>2.5</td> <td>6</td> <td>0049</td> <td>,</td> <td>0151</td> <td>_</td>	2.11	.9826	0431		2.5	6	0049	,	0151	_
2.13         .98344         .0413         2.58         .9951         .0143           2.14         .9838         .0404         2.59         .9952         .0139           2.15         .9842         .0396         2.69         .9953         .0139           2.16         .9846         .0387         2.61         .9955         .0132           2.17         .9850         .0379         2.62         .9956         .0129           2.18         .9854         .0371         2.63         .9959         .0122           2.19         .9857         .0363         2.64         .9959         .0122           2.20         .9864         .0339         2.67         .9962         .0116           2.22         .9868         .0339         2.67         .9962         .0113           2.23         .9871         .0312         2.68         .9963         .0110           2.24         .9868         .0339         2.67         .9962         .0113           2.25         .9878         .0317         2.70         .9965         .0104           2.26         .9981         .0310         2.71         .9966         .0104           2.26 <td>2.12</td> <td>.9830</td> <td></td> <td>- 1</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>	2.12	.9830		- 1						
2.14         9818         O404         2.59         .9952         .0139           2.15         .9846         .0396         2.60         .9953         .0139           2.16         .9846         .0387         2.61         .9955         .0132           2.17         .9850         .0379         2.62         .9956         .0129           2.18         .9854         .0371         2.63         .9957         .0126           2.19         .9857         .0363         2.64         .9959         .0122           2.20         .9861         .0355         2.65         .9960         .0119           2.21         .9864         .0347         2.66         .9961         .0116           2.221         .9868         .0339         2.67         .9962         .0119           2.221         .9875         .0252         2.69         .9964         .0101           2.222         .9878         .0317         2.70         .9965         .0104           2.25         .9878         .0310         2.71         .9966         .0101           2.26         .9881         .0310         2.71         .9966         .0010           2.27 <td>2.13</td> <td>.9834</td> <td></td> <td>- 1</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>	2.13	.9834		- 1						
2.15         .9842         .0.396         2.60         .9953         .0136           2.16         .9846         .0.387         2.61         .9955         .0132           2.17         .9850         .0.379         2.62         .9956         .0129           2.18         .9854         .0.317         2.63         .9957         .0163           2.19         .9857         .0.363         2.64         .9959         .0122           2.20         .9868         .0.339         2.67         .9962         .0116           2.21         .9868         .0.339         2.67         .9962         .0113           2.23         .9875         .0.017         2.70         .9965         .0110           2.24         .9875         .0.025         2.68         .9961         .0116           2.22         .9878         .0.0317         2.70         .9965         .0107           2.24         .9875         .0.0217         2.70         .9966         .0104           2.25         .9881         .0.031         2.71         .9966         .0104           2.26         .9881         .0.030         2.72         .9967         .0099	2.14	.9838		İ						
2.16         .9846         .0387         2.61         .9955         .0132           2.17         .9850         .0379         2.62         .9956         .0129           2.18         .9854         .0371         2.63         .9937         .0125           2.19         .9857         .0363         .264         .9959         .0126           2.20         .9861         .0355         2.65         .9960         .0119           2.21         .9864         .0347         2.66         .9961         .0116           2.22         .9868         .0339         2.67         .9962         .0113           2.23         .9871         .0332         2.68         .9963         .0110           2.24         .9875         .0255         2.69         .9964         .0107           2.24         .9875         .0252         2.69         .9963         .0110           2.25         .9878         .0317         2.70         .9965         .0104           2.26         .9881         .0310         2.71         .9966         .0101           2.27         .9884         .0330         2.72         .9966         .0033           2.29 <td>2.15</td> <td>.9842</td> <td></td> <td>- 1</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>	2.15	.9842		- 1						
2.17		İ			2,0		.,,,,,		1 .0150	
2.17         .9850         0.0379         2.62         .9956         0.129           2.18         .9854         0.0371         2.63         .9957         0.126           2.19         .9857         0.0363         2.64         .9959         0.0122           2.20         .9861         0.0355         2.65         .9960         0.0119           2.21         .9864         .0347         2.66         .9961         .0116           2.22         .9868         .0339         2.67         .9962         .0113           2.23         .9871         .0332         2.68         .9963         .0110           2.24         .9875         .0225         2.69         .9964         .0107           2.25         .9878         .0317         2.70         .9965         .0104           2.26         .9881         .0310         2.71         .9966         .0101           2.27         .9887         .0297         2.73         .9968         .0096           2.23         .9893         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0086           2.		.9846	.0387	- il	2.61	ı	.9955		.0132	
2.18         .9854         .0371         2.63         .9957         .0126           2.19         .9857         .0363         2.64         .9959         .0122           2.20         .9861         .0355         2.65         .9960         .0119           2.21         .9864         .0347         2.66         .9961         .0116           2.22         .9888         .0339         2.67         .9962         .0113           2.22         .9875         .0325         2.68         .9963         .0110           2.24         .9875         .02125         2.69         .9964         .0107           2.25         .9878         .0317         2.70         .9965         .0104           2.22         .9887         .0303         2.72         .9967         .0099           2.23         .9887         .0297         2.73         .9969         .0033           2.23         .9890         .0290         2.74         .9969         .0033           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .5898         .0270         2.77         .9972         .0086           2.33 <td></td> <td>.9850</td> <td>.0379</td> <td>ı II</td> <td>2.62</td> <td>2</td> <td>.9956</td> <td></td> <td></td> <td></td>		.9850	.0379	ı II	2.62	2	.9956			
2.20         .9861         .0355         2.65         .9969         .0112           2.21         .9864         .0347         2.66         .9961         .0116           2.22         .9868         .0339         2.67         .9962         .0113           2.23         .9871         .0332         2.68         .9963         .0110           2.24         .9875         .0225         2.69         .9964         .0107           2.25         .9878         .0317         2.70         .9965         .0104           2.22         .9881         .0310         2.71         .9966         .0101           2.22         .9887         .0297         2.73         .9968         .0096           2.23         .9890         .0290         2.74         .9969         .0093           2.30         .9893         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0079           2.35 <td></td> <td>.9854</td> <td>.0371</td> <td>- 11</td> <td>2.63</td> <td>, (</td> <td>.9957</td> <td></td> <td></td> <td></td>		.9854	.0371	- 11	2.63	, (	.9957			
2.21         .9864         .0347         2.66         .9961         .0116           2.22         .9868         .0349         2.67         .9962         .0113           2.23         .9871         .0332         2.68         .9963         .0110           2.24         .9875         .0325         2.69         .9964         .0107           2.25         .9878         .0317         2.70         .9965         .0104           2.26         .9881         .0310         2.71         .9966         .0101           2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .9968         .0093           2.29         .9890         .0280         2.74         .9969         .0093           2.30         .9893         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0086           2.31         .9901         .0264         2.78         .9973         .0086           2.33         .9901         .0258         2.79         .9974         .0079           2.36 <td></td> <td>.9857</td> <td>.0363</td> <td>- lì</td> <td>2.64</td> <td>۱ ا</td> <td>.9959</td> <td></td> <td>.0122</td> <td></td>		.9857	.0363	- lì	2.64	۱ ا	.9959		.0122	
2.22         9868         .0339         2.67         .9962         .0113           2.23         .9871         .0332         2.68         .9963         .0110           2.24         .9875         .0325         2.69         .9964         .0107           2.25         .9878         .0317         2.70         .9965         .0104           2.25         .9881         .0310         2.71         .9966         .0101           2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .9968         .0096           2.29         .9890         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .9898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0079           2.36         .9906         .0252         2.80         .9974         .0079           2.36	2.20	.9861	.0355		2.65	i i	.9960		.0119	
2.22         9868         .0339         2.67         .9962         .0113           2.23         .9871         .0332         2.68         .9963         .0110           2.24         .9875         .0325         2.69         .9964         .0107           2.25         .9878         .0317         2.70         .9965         .0104           2.25         .9881         .0310         2.71         .9966         .0101           2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .9968         .0096           2.29         .9890         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .9898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0079           2.36         .9906         .0252         2.80         .9974         .0079           2.36		1	}	- II		- 1				
2.23         .9871         .0312         2.68         .9963         .0110           2.24         .9875         .0217         2.69         .9864         .0107           2.25         .9878         .0317         2.70         .9865         .0104           2.26         .9881         .0310         2.71         .9966         .0101           2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .9968         .0096         .0093           2.29         .9890         .0290         2.74         .9969         .0093         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .9898         .0270         2.77         .9972         .0086           2.33         .9904         .0258         2.79         .9974         .0081           2.34         .9904         .0258         2.79         .9974         .0081           2.35         .9909         .0246         2.81         .9975         .0077           2.36         .9909         .0246         2.81         .9975         .007				- 11				ı	.0116	
2.24         .9875         .0325         2.69         .9964         .0107           2.25         .9878         .0317         2.70         .9865         .0104           2.26         .9881         .0310         2.71         .9966         .0101           2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .9968         .0096           2.29         .9890         .0290         2.74         .9969         .0093           2.30         .9893         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .5898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0079           2.36         .9900         .0246         2.81         .9975         .0077           2.37         .9911         .0231         2.82         .9976         .0075           2.38 <td></td> <td></td> <td></td> <td>- il</td> <td></td> <td></td> <td></td> <td>- 1</td> <td>.0113</td> <td></td>				- il				- 1	.0113	
2.25         .9878         .0317         2.70         .9965         .0104           2.26         .9881         .0310         2.71         .9966         .0101           2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .9968         .0096           2.29         .9890         .0290         2.74         .9969         .0093           2.30         .9893         .0270         2.76         .9971         .0088           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .5898         .0270         2.77         .9972         .0086           2.33         .9904         .0258         2.79         .9974         .0081           2.34         .9904         .0258         2.79         .9974         .0079           2.36         .9909         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0214         2.82         .9976         .0077           2.39 <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>.0110</td> <td></td>									.0110	
2.26         .9881         .0310         2.71         .9966         .0101           2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .968         .0096           2.29         .9887         .0297         2.73         .968         .0096           2.29         .9890         .0283         2.75         .9970         .0091           2.30         .9893         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .9898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0079           2.35         .9906         .0252         2.80         .9974         .0079           2.37         .9911         .0241         2.82         .9976         .0077           2.38         .9913         .0235         2.83         .9977         .0073           2.38				- 11				- 1		
2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .9968         .0096           2.29         .9890         .0290         2.74         .9969         .0093           2.30         .9891         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .5898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0079           2.35         .9906         .0252         2.80         .9974         .0079           2.36         .9909         .0246         2.81         .9975         .0077           2.38         .9913         .0231         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0069           2.41 <td>2.25</td> <td>.9878</td> <td>.0317</td> <td></td> <td>2,70</td> <td></td> <td>.9965</td> <td>- 1</td> <td>,0104</td> <td></td>	2.25	.9878	.0317		2,70		.9965	- 1	,0104	
2.27         .9884         .0303         2.72         .9967         .0099           2.28         .9887         .0297         2.73         .9968         .0096           2.29         .9890         .0290         2.74         .9969         .0093           2.30         .9891         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .5898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0079           2.35         .9906         .0252         2.80         .9974         .0079           2.36         .9909         .0246         2.81         .9975         .0077           2.38         .9913         .0231         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0069           2.41 <td>2.26</td> <td>9881</td> <td>0310</td> <td>- II</td> <td>2.71</td> <td>İ</td> <td>9966</td> <td>J</td> <td>0101</td> <td></td>	2.26	9881	0310	- II	2.71	İ	9966	J	0101	
2.28         .9887         .0.297         2.73         .9968         .0096           2.29         .9890         .0.290         2.74         .9969         .0093           2.30         .9893         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .3898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0061           2.35         .9909         .0246         2.81         .9975         .0077           2.36         .9909         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0229         2.84         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9921         .0224         2.85         .9978         .0069           2.41 </td <td></td> <td></td> <td></td> <td>- 11</td> <td></td> <td>- 1</td> <td></td> <td>- J</td> <td></td> <td></td>				- 11		- 1		- J		
2.29         .9890         0.290         2.74         .9969         .0093           2.30         .9893         0.283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .5898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0071           2.35         .9906         .0252         2.80         .9974         .0079           2.36         .9909         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0235         2.83         .9977         .0073           2.39         .9918         .0224         2.85         .9978         .0069           2.40         .9918         .0224         2.85         .9978         .0069           2.41         .9920         .0219         2.86         .9979         .0067           2.42 <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>ı</td> <td></td> <td></td>								ı		
2.30         .9893         .0283         2.75         .9970         .0091           2.31         .9896         .0277         2.76         .9971         .0088           2.32         .9898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0081           2.35         .9906         .0252         2.80         .9974         .0079           2.36         .9909         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0073           2.38         .9913         .0224         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9921         .0224         2.85         .9978         .0069           2.41         .9920         .0219         2.86         .9979         .0067           2.43         .9922         .0213         2.87         .9979         .0065           2.43 <td>2.29</td> <td></td> <td></td> <td>И</td> <td></td> <td>- 1</td> <td></td> <td>- 1</td> <td></td> <td></td>	2.29			И		- 1		- 1		
2.32         .9898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0081           2.35         .9906         .0252         2.80         .9974         .0079           2.36         .9999         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0235         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0069           2.41         .9920         .0219         2.86         .9979         .0067           2.42         .9922         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0061           2.44         .9927         .0203         2.89         .9981         .0060           2.45 <td>2.30</td> <td></td> <td></td> <td>- ii</td> <td></td> <td></td> <td></td> <td>- 1</td> <td></td> <td></td>	2.30			- ii				- 1		
2.32         .9898         .0270         2.77         .9972         .0086           2.33         .9901         .0264         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0081           2.35         .9906         .0252         2.80         .9974         .0079           2.36         .9999         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0235         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0069           2.41         .9920         .0219         2.86         .9979         .0067           2.42         .9922         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0061           2.44         .9927         .0203         2.89         .9981         .0060           2.45 <td></td> <td>1</td> <td>1</td> <td>-//</td> <td>-</td> <td>İ</td> <td></td> <td>- 1</td> <td></td> <td></td>		1	1	-//	-	İ		- 1		
2.33         .9901         .0254         2.78         .9973         .0084           2.34         .9904         .0258         2.79         .9974         .0081           2.35         .9906         .0252         2.80         .9974         .0079           2.36         .9909         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0073           2.38         .9913         .0223         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0069           2.41         .9920         .0219         2.86         .9979         .0067           2.43         .9922         .0213         2.87         .9979         .0065           2.43         .9927         .0203         2.89         .9981         .0061           2.44         .9927         .0203         2.89         .9981         .0061           2.44         .9927         .0203         2.89         .9981         .0061           2.44 <td></td> <td>.9896</td> <td>.0277</td> <td>-   </td> <td></td> <td>- 1</td> <td></td> <td>- 1</td> <td></td> <td></td>		.9896	.0277	-		- 1		- 1		
2.34         .9904         .0258         2.79         .9974         .0081           2.35         .9906         .0252         2.80         .9974         .0079           2.36         .9909         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0223         2.83         .9977         .0073           2.40         .9918         .0229         2.84         .9977         .0071           2.41         .9920         .0219         2.86         .9979         .0069           2.41         .9922         .0213         2.87         .9979         .0069           2.43         .9925         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0060           2.45         .9929         .0198         2.90         .9981         .0060           2.47         .9932         .0189         2.92         .9982         .0058           2.49         .9934         .0180         2.94         .9984         .0051           2.49 <td></td> <td></td> <td></td> <td>ll .</td> <td></td> <td>- 1</td> <td></td> <td>ļ</td> <td></td> <td></td>				ll .		- 1		ļ		
2.36         .9906         .0252         2.80         .9974         .0079           2.36         .9909         .0246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0235         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .99218         .0224         2.85         .9979         .0069           2.41         .9922         .0213         2.87         .9979         .0067           2.43         .9925         .0208         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0061           2.45         .9929         .0198         2.90         .9981         .0060           2.46         .9931         .0194         2.91         .9982         .0058           2.47         .9932         .0189         2.92         .9982         .0056           2.49         .9936         .0180         2.94         .9983         .0055           2.49 <td></td> <td></td> <td></td> <td>- !!</td> <td></td> <td></td> <td></td> <td>- 1</td> <td></td> <td></td>				- !!				- 1		
2.36         9909         .0.246         2.81         .9975         .0077           2.37         .9911         .0241         2.82         .9976         .0073           2.38         .9913         .0235         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0069           2.41         .9920         .0219         2.86         .9979         .0067           2.42         .9922         .0213         2.87         .9979         .0065           2.43         .9925         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0060           2.45         .9929         .0198         2.90         .9981         .0060           2.47         .9912         .0189         2.92         .9982         .0058           2.47         .9932         .0184         2.91         .9982         .0056           2.48         .9934         .0180         2.94         .9984         .0051           2.51 <td></td> <td></td> <td></td> <td></td> <td></td> <td>- 1</td> <td></td> <td>-  </td> <td></td> <td></td>						- 1		-		
2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0235         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0669           2.41         .9920         .0219         2.86         .9979         .0067           2.42         .9922         .0213         2.87         .9979         .0065           2.43         .9925         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0060           2.45         .9929         .0198         2.90         .9981         .0060           2.47         .9912         .0189         2.92         .9982         .0058           2.47         .9912         .0180         2.91         .9984         .0051           2.49         .9936         .0175         2.95         .9984         .0051           2.51         .9940         .0171         2.96         .9985         .0050           2.51 <td>2.35</td> <td>.9906</td> <td>.0252</td> <td>1</td> <td>2.80</td> <td></td> <td>.9974</td> <td></td> <td>.0079</td> <td></td>	2.35	.9906	.0252	1	2.80		.9974		.0079	
2.37         .9911         .0241         2.82         .9976         .0075           2.38         .9913         .0235         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0669           2.41         .9920         .0219         2.86         .9979         .0067           2.42         .9922         .0213         2.87         .9979         .0065           2.43         .9925         .0203         2.88         .9980         .0063           2.45         .9929         .0198         2.90         .9981         .0060           2.45         .9921         .0194         2.91         .9982         .0058           2.47         .9931         .0184         2.91         .9982         .0058           2.48         .9934         .0184         2.93         .9983         .0055           2.49         .9936         .0175         2.94         .9984         .0031           2.50         .9938         .0175         2.95         .9984         .0051           2.51 <td>2.36</td> <td>9909</td> <td>0246</td> <td>-</td> <td>2.81</td> <td></td> <td>.9975</td> <td>1</td> <td>.0077</td> <td></td>	2.36	9909	0246	-	2.81		.9975	1	.0077	
2.38         .9913         .0235         2.83         .9977         .0073           2.39         .9916         .0229         2.84         .9977         .0071           2.40         .9918         .0224         2.85         .9978         .0069           2.41         .9920         .0219         2.86         .9979         .0067           2.42         .9922         .0213         2.87         .9979         .0065           2.43         .9927         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0060           2.45         .9929         .0198         2.90         .9981         .0060           2.46         .9931         .0164         2.91         .9982         .0058           2.47         .9932         .0189         2.92         .9982         .0056           2.48         .9934         .0180         2.24         .9984         .003           2.49         .9936         .0180         2.24         .9984         .003           2.50         .9938         .0175         2.95         .9984         .0051           2.51				11		1	.9976	1	,0075	
2.40         .9918         .0224         2.85         .9978         .0069           2.41         .9920         .0219         2.86         .9979         .0067           2.42         .9922         .0213         2.87         .9979         .0065           2.43         .9925         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0061           2.45         .9929         .0198         2.90         .9981         .0060           2.46         .9931         .0194         2.91         .9982         .0058           2.47         .9932         .0189         2.92         .9982         .0056           2.48         .9936         .0180         2.94         .9984         .003           2.49         .9936         .0180         2.94         .9984         .003           2.50         .9938         .0175         2.95         .9984         .0051           2.51         .9940         .0171         2.96         .9985         .0050           2.52         .9941         .0167         2.277         .9985         .0048           2.53	2.38		.0235	IJ	2.83	1	.9977		.0073	
2.41         .9920         .0219         2.86         .9979         .0067           2.42         .9922         .0213         2.87         .9979         .0065           2.43         .9925         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0061           2.45         .9929         .0198         2.90         .9981         .0060           2.46         .9931         .0194         2.91         .9982         .0058           2.47         .9932         .0189         2.92         .9982         .0056           2.48         .9934         .0184         2.93         .9983         .0055           2.49         .9936         .0175         2.94         .9984         .0051           2.50         .9938         .0175         2.95         .9984         .0051           2.51         .9940         .0167         2.97         .9985         .0049           2.52         .9941         .0167         2.97         .9985         .0049           2.53         .9943         .0167         2.98         .9986         .0047	2.39	.9916	.0229	li .		ļ		1		
2.42         .9922         .0213         2.87         .9979         .0065           2.43         .9925         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0061           2.45         .9929         .0198         2.90         .9981         .0060           2.46         .9931         .0194         2.91         .9982         .0058           2.47         .9932         .0189         2.92         .9982         .0056           2.48         .9934         .0184         2.93         .9983         .0055           2.49         .9936         .0175         2.95         .9984         .0051           2.50         .9938         .0175         2.95         .9984         .0051           2.51         .9940         .0167         2.97         .9985         .0049           2.52         .9941         .0167         2.97         .9985         .0049           2.53         .9943         .0163         2.98         .9986         .0047	2.40	.9918	.0224	1	2.85	1	.9978		.0069	
2.42         .9922         .0213         2.87         .9979         .0065           2.43         .9925         .0203         2.88         .9980         .0063           2.44         .9927         .0203         2.89         .9981         .0061           2.45         .9929         .0198         2.90         .9981         .0060           2.46         .9931         .0194         2.91         .9982         .0058           2.47         .9932         .0189         2.92         .9982         .0056           2.48         .9934         .0184         2.93         .9983         .0055           2.49         .9936         .0175         2.95         .9984         .0051           2.50         .9938         .0175         2.95         .9984         .0051           2.51         .9940         .0167         2.97         .9985         .0049           2.52         .9941         .0167         2.97         .9985         .0049           2.53         .9943         .0163         2.98         .9986         .0047	7.41	0070	0210	1	206	1	0070		0067	
2.43 9925 0.0208 2.88 9980 0.063 2.44 9927 0.203 2.89 9981 0.061 2.45 9929 0.198 2.90 9981 0.060 2.46 9931 0.104 2.91 9982 0.058 2.47 9932 0.189 2.92 9982 0.058 2.48 9934 0.184 2.93 9983 0.055 2.49 9936 0.180 2.94 9984 0.053 2.49 9936 0.175 2.95 9983 0.051 2.51 9940 0.171 2.96 9985 0.050 2.51 9941 0.167 2.97 9985 0.068 2.52 9941 0.167 2.97 9985 0.048 2.53 9943 0.163 2.98 9985 0.047				11				ł		
2.44         .9927         .0203         2.89         .9981         .0061           2.45         .9929         .0198         2.90         .9981         .0060           2.46         .9931         .0194         2.91         .9982         .0058           2.47         .9932         .0189         2.92         .9982         .0056           2.49         .9934         .0184         2.91         .9983         .0059           2.49         .9936         .0175         2.95         .9984         .0031           2.50         .9938         .0175         2.95         .9984         .0051           2.51         .9940         .0171         2.96         .9985         .0050           2.52         .9941         .0167         2.97         .9985         .0047           2.53         .9943         .0163         2.28         .9986         .0047						1		ĺ		
2.45         .9929         .0198         2.90         .9981         .0060           2.46         .9931         .0194         2.91         .9982         .0058           2.47         .9932         .0189         2.92         .9982         .0056           2.48         .9934         .0184         2.93         .9983         .0055           2.49         .9936         .0175         2.94         .9984         .0031           2.50         .9938         .0175         2.95         .9984         .0051           2.51         .9940         .0171         2.96         .9985         .0050           2.52         .9941         .0167         2.97         .9985         .0048           2.53         .9943         .0163         2.98         .9986         .0047				11	2.89	1		1	.0061	
2.47         .9912         .0189         2.92         .9982         .0056           2.48         .9934         .0184         2.93         .9983         .0055           2.49         .9936         .0180         2.94         .9984         .0051           2.50         .9938         .0175         2.95         .9984         .0051           2.51         .9940         .0171         2.96         .9985         .0050           2.52         .9941         .0167         2.97         .9985         .0040           2.53         .9943         .0163         2.98         .9986         .0047					2.90		.9981		.0060	
2.47         .9912         .0189         2.92         .9982         .0056           2.48         .9934         .0184         2.93         .9983         .0055           2.49         .9936         .0180         2.94         .9984         .0051           2.50         .9938         .0175         2.95         .9984         .0051           2.51         .9940         .0171         2.96         .9985         .0050           2.52         .9941         .0167         2.97         .9985         .0040           2.53         .9943         .0163         2.98         .9986         .0047		ĺ		ł l		1	0003	l	0050	
2.48				11		1		1		
2.49 9936 .0180 2.94 .9984 .0053 2.50 9938 .0175 2.95 .9984 .0051 2.51 9940 .0171 2.96 .9985 .0050 2.52 9941 .0167 2.97 .9985 .0048 2.53 .9943 .0163 2.98 .9986 .0047				1				i		
2.50 .9938 .0175 2.95 .9984 .0051 2.51 .9940 .0171 2.96 .9985 .0050 2.52 .9941 .0167 2.97 .9985 .0048 2.53 .9943 .0163 2.98 .9986 .0047						1				
2.51 .9940 .0171 2.96 .9985 .0050 2.52 .9941 .0167 2.97 .9985 .0048 2.53 .9943 .0163 2.98 .9986 .0047				lí		1		1		
2.51 .9941 .0167 2.97 .9985 .0048 2.53 .9943 .0163 2.98 .9986 .0047	2.50	.7730		1		1		ĺ		
2.52 .9941 .0163 2.98 .9986 .0047	2.51	.9940						1		
2.33				1						
2.54   00.45   .01.26    2.99   .9980   .0040				1						
	2.54	.9945								
2,55 .9946 .0154 3.00 .9987 .0044	2.55	.9946	.0134	<u>_</u>	3.00		.,,,,			

2	Area	u Ordinate		Area	u Ordinate
1.21	.8869	.1919	1.66	.9515	.1006
1.22	.8888	.1895	1.67	.9525	.0989
1.23	.8907	.1872	1.68	.9535	
1.24	.8925	.1849	1.69		.0973
1.25	.8944	.1826	1.70	.9545	.0957
	.0,44	.1020	1.70	.9554	.0940
1.26	.8962	.1804	1.71	.9564	.0925
1.27	.8980	.1781	1.72	.9573	.0909
1.28	.8997	.1758	1.73	.9582	.0893
1.29	.9015	.1736	1.74	.9591	.0878
1.30	.9032	.1714	1.75	.9599	.0863
1.31	.9049	.1691	1.76		
1.32	.9066	.1669	1.77	.9608	.0848
1.33	.9082	.1647	1.78	.9616	.0833
1.34	.9099	.1626		.9625	.0818
1.35	.9115	.1604	1.79	.9633	.0804
	1		1,80	.9641	.0790
1.36	.9131	.1582	18.1	.9649	.0775
1.37	.9147	.1561	1.82	.9656	.0761
1.38 1.39	.9162	.1539	1.83	.9664	.0748
1.40	.9177	.1518	1.84	.9671	.0734
1.40	.9192	.1497	1.85	.9678	.0734
1.41	.9207	.1476	1		
1.42	.9222	.1456	1.86	.9686	.0707
1.43	.9236	.1435	1.87	.9693	.0694
1.44	.9251	.1415	1.88	.9699	.0681
1.45	.9265	.1394	1.89	.9706	.0669
1.46		1	1.90	.9713	.0656
1.47	.9279	.1374	1,91	.9719	
1.48	.9292 .9306	.1354	1.92	.9726	.0644
1.49		.1334	1.93	.9732	.0632
1.50	.9319 .9332	.1315	1.94	.9738	.0620
	.9332	.1295	1.95	.9744	.0608
1.51	.9345	.1276	1.96		.0370
1.52 1.53	.9357	.1257	1.96	.9750	.0584
1.54	.9370	.1238	1.98	.9756	.0573
1.55	.9382	.1219	1.99	.9761	.0562
1.33	.9394	.1200	2.00	.9767 .9772	.0551
1.56	9406	.1182	'	.5,12	.0540
1.57	.9418	.1163	2.01	.9778	.0529
1.58	.9429	.1145	2.02	.9783	.0519
1.59	.9441	.1127	2.03	.9788	.0508
1.60	.9452	.1109	2.04	.9793	.0498
1.61	.9463		2.05	.9798	.0488
1 62	.9463	1092	2.06		
1.63	.9474	.1074	2.05	.9803	.0478
1.64	.9484	.1057	2.08	.9808	.0468
1.65	.9505	.1040	2.09	.9812	.0459
	1,7303	.1023	2.10	.9817 .9821	.0449
					.0440

TABLE D PERCENTILE POINTS OF E-DISTRIBUTIONS

Percentile\*

1							•							
1.158   3.25   3.10   7.27   1.000   1.376   1.963   3.078   6.314   4.301   6.3651   63.65	df	**			70	75			90	95	97.5	99	99.5	99.95
1 138 325 510 .727 1.000 1.376 1.903 5.00 2.900 4.300 6.965 9.925 31.939 3 1.17 .277 .424 .584 .765 978 1.250 1.638 2.353 3.182 4.541 5.841 12.941 4 1.34 2.71 .444 .569 .741 9.44 1.190 1.533 2.132 2.776 3.747 4.604 8.610 5 1.32 .267 .408 .559 .727 920 1.156 1.476 2.015 2.571 3.165 4.032 6.859 6 1.31 .265 .404 .553 .718 9.06 1.134 1.440 1.934 2.247 3.165 2.998 3.499 5.405 8 1.30 .262 .309 .546 .706 .889 1.108 1.397 1.860 2.366 2.898 3.355 5.041 8 1.30 .262 .309 .546 .706 .889 1.108 1.397 1.860 2.366 2.8281 3.20 4.781 10 .129 .260 .397 .542 .700 .879 1.093 1.372 1.812 2.228 2.764 3.169 4.781 11 .129 .260 .396 .540 .697 .876 1.088 1.301 .183 1.332 2.22 2.281 3.204 1.781 12 .128 .259 .395 .539 .695 .873 1.033 1.366 1.782 2.179 2.681 3.055 4.318 13 .132 .259 .394 .538 .694 .870 1.079 1.330 1.771 2.145 2.242 2.277 4.140 14 .128 .258 .393 .537 .692 .868 1.076 1.345 1.761 15 .128 .258 .392 .535 .691 .865 1.074 1.341 1.753 2.131 2.602 2.947 4.140 16 .128 .258 .392 .535 .691 .868 .861 1.074 1.341 1.753 2.131 2.602 2.947 4.140 17 .128 .257 .392 .534 .688 .861 1.064 1.325 1.725 2.093 2.914 .015 18 .127 .257 .391 .533 .687 .860 1.064 1.325 1.725 2.083 2.921 4.015 19 .127 .257 .391 .533 .687 .860 1.064 1.325 1.725 2.065 2.258 2.845 3.850 20 .127 .256 .390 .532 .686 .859 1.006 1.323 1.724 2.003 2.592 2.818 3.832 21 .127 .256 .390 .532 .686 .859 1.006 1.323 1.724 2.003 2.893 3.952 22 .127 .256 .390 .532 .686 .859 1.006 1.323 1.721 2.006 2.482 2.797 3.745 23 .127 .256 .390 .531 .684 .856 1.038 1.315 1.700 2.025 2.473 2.771 3.073 24 .127 .256 .390 .531 .684 .856 1.038 1.315 1.700 2.025 2.479 2.779 3.735 25 .127 .256 .390 .531 .684 .856 1.058 1.315 1.701 2.042 2.492 2.797 3.745 26 .127 .256 .390 .531 .684 .856 1.058 1.315 1.701 2.042 2.492 2.797 3.745 26 .127 .256 .390 .531 .684 .856 1.058 1.315 1.701 2.042 2.492 2.797 3.745 27 .127 .256 .390 .531 .684 .856 1.058 1.315 1.701 2.042 2.492 2.797 3.745 28 .127 .256 .390 .531 .684 .856 1.058 1.315 1.700 2.042 2.492 2.797 3.745 29 .127 .256 .390 .531 .684 .856 1.058 1.315 1.700 2.042 2				0.5	70	13	-			<u> </u>	12 706	31.821	63.657	636.619
2 142 289 445 .617 816 1.061 1.386 1.886 2.353 1.182 4.541 5.841 12.941   4 1.134 .271 444 .569 .741 9.41 1.190 1.513 2.132 2.776 3.747 4.061 8.610   5 1.32 .267 .408 .559 .772 9.20 1.156 1.476 .2015 2.571 3.165 4.032 6.859   7 1.30 .265 .402 .539 .711 8.96 1.119 1.415 1.895 2.365 2.998 3.499 5.405   8 1.10 .262 .399 .546 .706 .889 1.08 1.197 1.860 2.366 2.896 3.355 5.041   9 1.29 .260 .198 .543 .703 .881 1.100 1.183 1.833 2.262 2.821 3.250 4.781   10 1.29 .260 .397 .542 .700 .879 1.093 1.372 1.812 2.282 2.764 3.169 4.587   11 1.29 .260 .396 .540 .697 .876 1.088 1.156 1.796 2.201 2.718 3.066 4.371   12 1.28 .259 .394 .538 .694 .870 1.079 1.350 1.771 2.160 2.650 3.012 4.211   12 1.28 .258 .393 .535 .691 .866 1.076 1.345 1.761 2.145 2.622 2.977 4.073   15 1.28 .258 .393 .535 .691 .866 1.074 1.341 1.753   18 1.27 .257 .392 .514 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .514 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .514 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .534 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .534 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .534 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .534 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .534 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .534 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .534 .688 .862 1.067 1.331 1.740   1.27 .257 .392 .534 .688 .862 1.067 1.331 1.740   1.28 .256 .390 .531 .684   8.55 1.081 0.131 1.711   2.060 2.592 .288 3.953   20 .127 .256 .390 .531 .684   8.55 1.081 0.131 1.711   2.060 2.492 2.797 3.745   2.604 2.492 2.797 3.745   2.604 2.492 2.797 3.745   2.605 2.593 .530 .683   8.54 1.055 1.331 1.760   2.007 2.432 2.703 3.646   8.55 1.035 1.331 1.760   2.007 2.432 2.703 3.646   8.55 1.035 1.331 1.760   2.007 2.432 2.703 3.767   3.767 3.767		.158	.325	.510	.727	1.000	1.376	1.963	3.078	0.314			9.925	
31         1.17         277         424         5.84         .765         .978         1.250         1.638         2.132         2.776         3.717         4 (64)         8.610           4         1.13         2.131         .416         5.99         .777         920         1.156         1.476         2.015         2.571         3.365         4.032         6.859           6         1.131         .265         .404         .553         .718         .906         1.131         1.440         1.943         2.247         3.143         3.707         5.99           8         1.102         .625         .402         .599         .711         .896         1.191         1.415         1.895         2.366         2.998         3.499         5.043           9         1.292         .61         .398         .543         .703         .881         1.100         1.292         .62         .399         .542         .700         .879         1.931         1.322         .222         .2821         .2200         .4718         .4781           11         1.292         .260         .396         .540         .697         .876         1.088         1.361         1.792	2						1.061	1 386	1.886	2.920				12.941
4         1.134         271         414         569         741         .941         1.139         2.121         2.571         3.365         4.032         6.859           5         1.132         .267         .408         .559         .727         .920         1.156         1.476         .2015         2.217         3.365         4.032         6.859           7         .130         .263         .402         .549         .711         .896         1.191         .141         1.893         2.247         .314         3.707         .549           9         .129         .261         .398         .543         .703         .883         1.100         .138         .222         .281         3.259         .4781           10         .129         .260         .397         .542         .700         .879         1.093         1.372         1.812         .222         .281         3.259         .4781           11         .129         .260         .396         .540         .697         .876         1.088         1.531         .1796         .282         .2761         .169         .4871           12         .128 .258         .393         .535         <	3						078	1 250	1.638	2.333			4 604	8.610
5         .132         .267         .408         .559         .727         .920         1.136         .1476         .2447         .1431         3.707         .5959           6         .131         .265         .404         .553         .718         .906         1.134         1.440         1.943         .247         .1431         3.707         .5959           8         .130         .261         .399         .546         .706         .889         1.108         1.397         1.860         2.306         .289         3.355         .5041           9         .129         .261         .398         .543         .703         .881         1.100         1.831         .833         .222         .282         .2355         .505         .504           11         .129         .260         .396         .540         .697         .876         1.088         1.361         .1782         .219         .2681         .3055         .431           12         .128         .259         .394         .538         .694         .870         1.091         .130         .1771         .2160         .2681         .3055         .4318           12         .128         .25	4						041	1 190	1.533	2,134		3 365	4.032	6.859
6   1.31   2.65   AQ4   5.53   7.18   906   1.134   1.440   1.943   2.447   3.143   3.707   5.959   7.140   7.	5						.920	1.156	1.476	2.015	2.311	3.555		
6 1.31 265 404 .553 .718 .906 1.134 1.440 1.895 2.365 2.998 3.499 5.405 8 1.100 1.265 1.005 1.005 1.105 1.105 1.005 1.105 1.005 1.105 1.005 1.105 1.005 1.105 1.005 1.105 1.005 1.105 1.005 1.105 1.005 1.105 1.005 1.105 1.005 1.10				2400							2 447	3 143	3.707	
7 1.10 2.63 402 549 7.11 8.96 1.119 1.413 1.89 2.202 2.203 3.355 5.041   9 1.29 2.61 398 5.45 7.06 8.89 1.08 1.397 1.860   1.29 2.60 3.97 5.42 7.00 8.79 1.081 1.831 1.833   1.29 2.60 3.97 5.42 7.00 8.79 1.093 1.372 1.812 2.222 2.764 3.169 4.587   11	6	.131	265	401	553	.718	.906	1.134	1.440	1.943			3.499	
8         1.130         2.62         3.99         5.46         706         .859         1.198         1.39         2.262         2.821         3.250         4.781           9         1.129         2.60         1.97         3.83         1.100         1.383         1.833         2.222         2.261         3.169         4.587           11         1.129         2.60         .396         .540         .697         .876         1.083         1.136         1.782         2.222         2.261         3.169         4.587           11         1.129         .260         .396         .540         .697         .876         1.083         1.363         1.796         2.201         2.183         3.055         4.387           12         1.128         .299         .394         .537         .692         .868         1.076         1.345         1.761         2.160         2.681         3.055         4.31           15         1.282         .258         .393         .537         .691         .866         1.074         1.341         1.755         2.131         2.602         2.997         4.015           16         .1.28         .257         .392         .534	7						007	1 110	1 413	1.073			3.355	
9 1.129 2.61 398 3.43 703 8.83 1.100 1.83 1.432 2.228 2.764 3.169 4.587 11 1.129 2.60 3.96 5.40 6.97 8.76 1.083 1.361 1.796 2.201 2.118 3.105 4.387 12 1.128 2.29 3.95 5.39 6.95 8.76 1.083 1.361 1.796 2.201 2.118 3.105 4.318 13 1.28 2.29 3.95 5.39 6.97 8.76 1.083 1.361 1.796 2.201 2.118 3.105 4.318 14 1.28 2.58 3.93 5.37 6.92 8.68 1.076 1.345 1.761 2.145 2.204 2.277 4.140 15 1.28 2.58 3.93 5.37 6.92 8.66 1.074 1.341 1.753 2.131 2.602 2.947 4.073 16 1.28 2.58 3.93 5.35 6.90 8.65 1.074 1.337 1.746 2.102 2.567 2.898 3.955 17 1.28 2.57 3.92 5.34 6.68 8.62 1.076 1.333 1.740 2.110 2.567 2.898 3.955 18 1.27 2.57 3.92 5.34 6.68 8.62 1.061 3.23 1.724 2.003 2.532 2.878 3.922 2.127 2.57 3.91 5.33 6.68 8.61 1.066 1.323 1.724 2.003 2.532 2.888 3.955 19 1.27 2.57 3.91 5.33 6.68 8.61 1.066 1.323 1.724 2.003 2.532 2.881 3.830 12 1.27 2.57 3.95 5.39 6.69 8.60 1.064 1.325 1.725 2.066 2.528 2.845 3.850 12 1.27 2.57 3.95 5.39 6.68 8.86 1.061 1.321 1.717 2.004 2.502 2.287 3.850 12 1.27 2.56 3.90 5.31 6.68 8.51 1.061 1.321 1.717 2.004 2.502 2.819 3.767 2.812 2.77 2.56 3.90 5.31 6.88 8.56 1.081 1.321 1.717 2.004 2.492 2.797 3.745 2.122 2.56 3.90 5.31 6.88 8.56 1.081 1.316 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.90 5.31 6.88 8.56 1.081 1.316 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.90 5.31 6.88 8.56 1.081 1.316 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.90 5.31 6.88 8.56 1.081 1.316 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.90 5.31 6.88 8.56 1.081 1.316 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.89 5.30 6.83 8.54 1.005 1.313 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.89 5.30 6.83 8.54 1.005 1.313 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.89 5.30 6.83 8.54 1.005 1.313 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.89 5.30 6.83 8.54 1.005 1.313 1.704 2.004 2.492 2.797 3.745 2.122 2.56 3.89 5.30 6.83 8.54 1.005 1.313 1.704 2.004 2.492 2.797 3.745 2.122 2.55 3.89 5.30 6.83 8.54 1.005 1.313 1.704 2.004 2.492 2.756 3.659 2.005 2.205 2.805 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305 2.305	8	.130	262	700	546		000	1 108	1.397	1.800		7 821	3.250	
1.12   2.60   3.96   5.40   6.97   8.76   1.083   1.372   1.812   1.212   2.203   3.96   5.40   6.97   8.76   1.083   1.361   1.725   2.101   2.183   3.055   3.121   1.212   2.295   3.95   5.39   6.95   8.73   1.033   1.356   1.782   2.190   2.650   3.012   4.221   1.212   2.256   3.93   5.37   6.92   8.68   1.076   1.345   1.761   2.145   2.624   2.977   4.073   1.212   2.258   3.93   5.37   6.91   8.66   1.076   1.341   1.753   2.131   2.602   2.947   4.073   1.212   2.258   3.93   5.35   6.91   8.66   1.074   1.341   1.753   2.131   2.602   2.947   4.073   1.212   2.257   3.92   5.34   6.689   8.65   1.074   1.341   1.753   2.131   2.602   2.947   4.073   1.212   2.277   3.92   5.34   6.689   8.65   1.067   1.333   1.740   2.101   2.567   2.898   3.965   1.074   1.212   2.277   3.91   5.33   6.88   8.61   1.064   1.323   1.725   2.086   2.288   3.955   1.272   2.277   3.91   5.33   6.887   8.60   1.064   1.323   1.725   2.086   2.282   2.884   3.850   1.272   2.273   3.95   3.853   3.853   2.274   4.073   3.85	9						003	1 100	1.385	1,033		2.764	3.169	4.587
11   1.129   2.60   3.96   5.40   6.97   8.76   1.088   1.363   1.796   2.201   2.718   3.106   4.319     12   1.28   2.59   3.95   5.39   6.95   8.73   1.083   1.356   1.782   2.179   2.681   3.055   4.318     13   1.28   2.59   3.95   5.39   6.95   8.73   1.083   1.356   1.782   2.172   2.160   2.650   3.012   4.221     14   1.28   2.58   3.93   3.57   6.92   8.68   1.076   1.345   1.753   2.131   2.602   2.947   4.140     15   1.28   2.58   3.93   5.35   6.91   8.66   1.074   1.341   1.753   2.131   2.602   2.947   4.073     16   1.28   2.58   3.92   5.35   6.90   8.65   1.074   1.341   1.753   2.131   2.602   2.947   4.073     16   1.28   2.57   3.92   5.34   6.68   8.61   1.069   1.333   1.740   2.110   2.567   2.898   3.965     17   1.28   2.57   3.92   5.34   6.68   8.62   1.067   1.330   1.744   2.101   2.557   2.898   3.965     18   1.27   2.57   3.91   5.33   6.68   8.61   1.064   1.325   1.729   2.093   2.539   2.861   3.883     1.27   2.57   3.91   5.33   6.68   8.60   1.064   1.325   1.729   2.093   2.532   2.845   3.850     127   2.57   3.91   5.32   6.66   8.59   1.063   1.323   1.721   2.004   2.593   2.813   3.192     21   1.27   2.56   3.90   5.32   6.66   8.59   1.063   1.321   1.717   2.004   2.593   2.819   3.792     22   1.27   2.56   3.90   5.31   6.68   8.56   1.058   1.316   1.710   2.004   2.492   2.797   3.745     23   1.27   2.56   3.90   5.31   6.68   8.56   1.058   1.316   1.710   2.004   2.492   2.797   3.745     24   1.27   2.56   3.90   5.31   6.68   8.56   1.058   1.316   1.710   2.004   2.492   2.777   3.765     25   1.27   2.56   3.90   5.31   6.68   8.56   1.058   1.315   1.706   2.025   2.473   2.771   3.600     26   1.27   2.25   3.89   5.30   6.68   8.51   1.051   1.311   1.701   2.004   2.472   2.763   3.673     27   1.27   2.56   3.90   5.30   6.68   8.51   1.051   1.311   1.701   2.004   2.472   2.763   3.673     28   1.27   2.25   3.89   5.30   6.68   8.51   1.051   1.311   1.701   2.004   2.472   2.763   3.673     29   1.27   2.56   3.99   5.30   6.83   8.51   1.05	10						879	1 093	1.372	1.812	2,220	20.0		
11         1.19         2.60         3.66         5.40         6.97         8.76         1.088         1.366         1.79         2.681         3.055         4.318           12         1.28         2.29         1.94         5.99         8.93         1.381         1.361         1.782         2.160         2.650         3.012         4.221           13         1.28         2.29         1.94         5.99         8.68         1.076         1.345         1.762         2.131         2.602         2.947         4.073           16         1.28         2.28         3.93         .535         .690         .865         1.071         1.331         1.740         2.131         2.602         2.947         4.073           17         1.28         2.277         2.95         .595         .690         .865         1.071         1.331         1.740         2.101         2.567         2.898         3.921           18         1.27         2.57         .92         .534         .688         .861         1.069         1.333         1.740         2.101         2.557         2.898         3.922           12         1.27         2.57         .391         .533			.200	.391	.342	.700					2 201	2.718		
12	11	120	760	206	540	697	.876	1.088	1.363	1.796				
13         1.28         259         394         .538         .694         .870         1.079         1.353         1.761         2.145         2.622         2.977         .4140           14         .128         .258         .393         .537         .692         .866         1.074         1.341         1.753         2.131         2.602         2.947         4.073           15         .128         .258         .393         .535         .690         .865         1.071         1.337         .746         2.100         .2583         2.921         4.015           16         .128         .257         .392         .534         .689         .861         1.091         .1333         1.746         2.110         .2563         2.893         3.952           18         .127         .257         .192         .534         .688         .862         1.069         1.333         1.740         2.100         .2592         .2883         3.922           20         1.27         .257         .911         .533         .687         .860         1.064         1.123         1.72         2.085         2.252         2.888         3.922           21         .127	12						0.77	1 001	1 356	1./04			3.012	
14         1.28         2.58         3.93         537         692         .868         1.076         1.493         1.75         2.131         2.092         2.947         4.015           15         .128         .258         .393         .536         .691         .866         1.074         1.341         1.755         2.131         2.602         2.947         4.015           16         .128         .258         .392         .535         .600         .865         1.071         1.331         1.740         2.100         .2567         .2893         .395           17         .128         .257         .392         .534         .689         .861         1.060         1.333         1.740         2.100         .2562         .2893         .395           18         .127         .257         .392         .534         .689         .861         1.060         1.323         1.720         2.030         .2519         .2861         3.883           20         .127         .257         .391         .533         .686         .859         1.061         1.321         1.712         .2066         2.288         1.061         1.321         1.714         .2069         2.509 <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>2.977</td> <td></td>													2.977	
1.128   2.258   3.93   5.36   6.691   8.66   1.074   1.341   1.735   1.746   2.120   2.583   2.921   4.015     1.28   2.256   3.92   5.315   6.690   8.65   1.071   1.337   1.746   2.120   2.567   2.898   3.952     1.28   2.257   3.92   5.34   6.68   8.62   1.069   1.333   1.740   2.010   2.552   2.878   3.952     1.27   2.257   3.91   5.33   6.68   8.61   1.066   1.232   1.72   2.085   2.252   2.845   3.850     1.27   2.257   3.91   5.33   6.687   8.60   1.064   1.325   1.725     1.27   2.257   3.91   5.32   6.66   8.99   1.063   1.323   1.721   2.074   2.593   2.819   3.792     21   1.27   2.256   3.90   5.32   6.66   8.89   1.061   1.321   1.717   2.064   2.492   2.797   3.745     22   1.27   2.256   3.90   5.31   6.684   8.55   1.058   1.316   1.708     23   1.27   2.256   3.90   5.31   6.684   8.55   1.058   1.316   1.708     24   1.27   2.256   3.90   5.31   6.684   8.55   1.058   1.316   1.708     25   1.27   2.256   3.90   5.31   6.684   8.55   1.058   1.316   1.708     26   1.27   2.256   3.90   5.31   6.684   8.55   1.058   1.315   1.706   2.052   2.473   2.771   3.670     27   1.27   2.256   3.99   5.30   6.683   8.54   1.055   1.313   1.701   2.045   2.467   2.753   3.674     28   1.27   2.256   3.99   5.30   6.683   8.54   1.055   1.311   1.699   2.042   2.475   2.750   3.664     28   1.27   2.256   3.99   5.30   6.633   8.54   1.055   1.310   1.697   2.042   2.475   2.750   3.664     28   1.27   2.256   3.99   5.30   6.633   8.54   1.055   1.310   1.697   2.042   2.475   2.750   3.664     28   1.27   2.256   3.99   5.30   6.633   8.54   1.055   1.310   1.697   2.042   2.475   2.750   3.664     38   1.28		120	-239	.394	.338								2.947	4.073
1.0							866	1.074	1.341	1.753	2.151	2		
16         .128         .256         .392         .515         .690         .865         1.071         1.33         1.740         2.110         2.567         2.898         3.922           18         .127         .257         .392         .534         .688         .862         1.061         1.333         1.740         2.003         2.539         2.861         3.883           1.27         .257         .391         .533         .687         .860         1.064         1.128         1.729         2.093         2.539         2.861         3.881           21         .127         .257         .391         .532         .686         .891         1.063         1.323         1.721         2.074         .508         2.819         3.792           21         .127         .256         .390         .532         .686         .891         1.063         1.331         1.712         2.008         2.518         2.831         3.819           22         .127         .256         .390         .531         .685         .851         1.018         1.171         2.060         2.518         2.831         3.792           24         .127         .256         .390		-128	.238	.393	.536	.691					2 120	2.583		
17   172   173   174   175	16	120	260	207	625	690	.865	1.071	1.337	1.740		2.567		
18         .127         .257         .392         .534         .688         .862         1.067         .1.30         .1.29         .203         .2.59         2.861         3.881           10         .127         .257         .391         .533         .688         .861         1.064         1.123         1.725         .208         .2.518         2.831         3.819           21         .127         .257         .391         .532         .686         .859         1.061         .1323         1.721         .200         2.518         2.831         3.819           22         .127         .256         .390         .532         .686         .859         1.061         .1323         1.721         .200         2.518         2.831         3.819           23         .127         .256         .390         .531         .685         .851         .1061         .1321         .171         .200         .2.518         2.831         .3.819           24         .127         .256         .390         .531         .684         .855         .1.351         .1.106         .2.062         .2.492         .2.779         .3.707           25         .127         .256												2,552		
127   257   391   533   688   861   1.066   1.323   1.725   2.086   2.538   2.843   3.859     127   257   391   533   687   860   1.064   1.323   1.725   2.080   2.518   2.831   3.819     127   257   391   532   686   859   1.063   1.323   1.721   2.080   2.518   2.831   3.792     128   127   256   390   532   686   885   1.061   1.321   1.717   2.089   2.590   2.819   3.792     128   127   256   390   531   685   887   1.099   1.318   1.711   2.084   2.492   2.797   3.745     127   256   390   531   684   8.55   1.058   1.316   1.703   2.080   2.485   2.787   3.725     127   256   390   531   684   8.55   1.058   1.315   1.706   2.052   2.479   2.719   3.707     127   256   389   530   683   855   1.058   1.311   1.701   2.054   2.467   2.763   3.659     127   256   389   530   683   855   1.058   1.311   1.701   2.045   2.467   2.763   3.659     127   256   389   530   683   854   1.035   1.311   1.699   2.045   2.457   2.750   3.646     127   256   389   530   683   854   1.035   1.311   1.699   2.045   2.457   2.750   3.646     127   256   389   350   683   854   1.035   1.311   1.699   2.045   2.457   2.750   3.646     127   256   389   350   683   854   1.035   1.311   1.699   2.045   2.457   2.750   3.646     128   255   388   3.529   681   851   1.050   1.303   1.684   2.021   2.432   2.704   3.551     128   126   225   388   3.529   681   851   1.050   1.303   1.684   2.020   2.393   2.660   3.460     128   224   336   526   677   845   1.041   1.289   1.655   1.960   2.332   2.576   3.271     129   126   224   336   526   677   845   1.041   1.289   1.655   1.665   1.960   2.326   2.576   3.271												2.539	2.861	
20												2.528	2.845	3.850
1.   1.   1.   1.   1.   1.   1.   1.							860	1.064	1.325	1.725	2.000			
21         1.27         257         391         532         686         859         1.063         1.221         1.717         2.074         2.598         2.891         37.767           22         1.27         2.56         .390         .532         .685         .858         1.061         1.312         1.714         2.096         2.590         2.897         3.767           24         1.27         .256         .390         .531         .684         855         1.081         1.318         1.711         2.064         2.492         2.797         3.725           25         1.27         .256         .390         .531         .684         8.56         1.038         1.316         1.708         2.065         2.492         2.779         3.707           26         1.27         .256         .390         .531         .684         8.56         1.038         1.315         1.706         2.052         2.473         2.771         3.670           27         1.27         .256         .390         .531         .684         .855         1.051         1.313         1.70         2.042         2.472         2.773         3.707           28         1.27	20	-14/	.257	.391	.555	.007					2.080	2.518		
22 1.17 256 390 532 .686 .838 1.061 1.319 1.714 2.069 2.590 2.807 3.745 24 1.27 2.56 390 .531 .685 .857 1.059 1.318 1.711 2.065 2.492 2.793 3.745 24 1.27 2.56 390 .531 .685 .856 1.058 1.316 1.708 2.060 2.485 2.787 3.725 25 1.27 2.56 390 .531 .684 .856 1.058 1.316 1.708 2.060 2.485 2.787 3.725 27 1.27 2.56 3.39 .531 .684 .856 1.058 1.316 1.701 2.052 2.473 2.771 3.690 2.816 2.712 2.56 3.39 .530 .683 .855 1.056 1.311 1.090 2.045 2.462 2.756 3.659 2.127 2.56 3.39 .530 .683 .854 1.035 1.311 1.699 2.045 2.462 2.756 3.659 2.127 2.56 3.39 .530 .683 .854 1.035 1.310 1.699 2.045 2.462 2.750 3.646 2.127 2.56 3.39 .530 .683 .854 1.035 1.310 1.699 2.045 2.473 2.771 3.660 2.127 2.56 3.39 .530 .683 .854 1.035 1.310 1.699 2.045 2.457 2.750 3.646 2.126 2.255 3.38 .529 .681 .851 1.050 1.303 1.684 2.000 2.390 2.650 3.666 0.126 2.254 3.36 7.527 6.79 .845 1.046 1.296 1.671 8.200 2.338 2.617 3.373 1.201 1.26 2.24 3.36 5.26 6.77 842 1.001 1.289 1.655 1.960 2.332 2.576 3.291	21	127	267	201	622	686	.859	1.063	1.323	1.721	2.074			
23   127   226   390   532   685   838   1,060   1,378   1,711   2,064   2,492   2,797   3,725   24   127   226   390   531   684   856   1,038   1,316   1,708   2,069   2,485   2,787   3,725   25   127   226   390   531   684   8,56   1,038   1,316   1,708   2,065   2,479   2,779   3,707   26   127   226   339   531   684   8,56   1,038   1,315   1,706   2,056   2,479   2,779   3,707   27   127   226   339   531   684   8,55   1,037   1,311   1,701   2,045   2,467   2,773   3,674   28   127   226   339   530   683   8,51   1,051   1,311   1,699   2,045   2,457   2,753   3,674   29   127   226   339   530   683   8,54   1,055   1,310   1,697   2,045   2,457   2,753   3,674   29   127   226   339   530   683   8,54   1,055   1,310   1,697   20   127   226   339   530   683   8,54   1,055   1,310   1,697   20   126   225   3,38   5,29   681   8,104   1,296   1,671   2,000   2,358   2,617   3,373   20   126   225   338   5,20   677   845   1,041   1,289   1,658   1,960   2,358   2,576   3,279   210   126   225   3,36   5,26   677   845   1,041   1,289   1,658   1,960   2,352   2,576   3,279   220   126   225   336   5,26   677   845   1,041   1,289   1,658   1,960   2,352   2,576   3,279											2.069			
24 1.27 2.56 3.90 5.51 6.84 8.57 1.039 1.316 1.708 2.060 2.485 2.787 3.727 2.56 3.90 5.51 6.84 8.55 1.038 1.316 1.708 2.052 2.479 2.771 3.690 2.71 1.27 2.56 3.89 5.51 6.84 8.55 1.058 1.315 1.706 2.052 2.473 2.771 3.690 2.81 2.77 2.56 3.89 5.50 6.83 8.55 1.066 1.315 1.701 2.045 2.462 2.756 3.659 2.127 2.56 3.89 5.50 6.83 8.55 1.066 1.316 1.099 2.045 2.462 2.756 3.659 2.127 2.56 3.89 5.50 6.83 8.54 1.055 1.310 1.699 2.045 2.462 2.756 3.659 2.127 2.56 3.89 5.50 6.83 8.54 1.055 1.310 1.699 2.045 2.462 2.756 3.659 2.127 2.56 3.89 5.50 6.83 8.54 1.055 1.310 1.699 2.045 2.462 2.750 3.646 2.126 2.255 3.88 5.29 6.81 8.51 1.050 1.303 1.684 2.000 2.390 2.660 3.460 2.126 2.254 3.87 5.27 6.79 8.48 1.046 1.296 1.671 8.000 2.339 2.650 3.450 2.126 2.254 3.85 5.26 6.77 8.42 1.005 1.282 1.645 1.960 2.332 2.576 3.291												2.492		
25 1.17 2.56 3.90 5.51 6.84 8.56 1.038 1.310 1.705 2.056 2.479 2.779 3.707 2.66 1.072 2.66 3.90 5.31 6.84 8.56 1.058 1.315 1.706 2.032 2.473 2.771 3.690 2.77 1.27 2.56 3.89 5.31 6.84 8.55 1.057 1.314 1.703 2.045 2.467 2.763 3.674 2.87 1.27 2.56 3.89 5.30 6.83 8.55 1.056 1.313 1.701 2.045 2.467 2.763 3.674 2.97 1.27 2.56 3.89 5.30 6.83 8.54 1.035 1.311 1.699 2.045 2.452 2.756 3.659 3.610 1.27 2.55 3.89 5.30 6.83 8.54 1.035 1.310 1.697 2.042 2.475 2.750 3.664 6.0 1.26 2.55 3.88 5.29 6.81 8.51 1.050 1.301 1.684 2.000 2.390 2.660 3.450 6.0 1.26 2.54 3.87 5.27 6.79 8.84 1.045 1.26 1.57 8.84 2.000 2.338 2.617 3.373 1.20 1.26 2.54 3.85 5.26 6.77 8.84 1.041 1.289 1.653 1.960 2.335 2.617 3.373 1.20 1.26 2.54 3.85 5.26 6.77 8.84 1.041 1.289 1.653 1.960 2.335 2.576 3.291												2.485	2.787	3.725
26 . 127 . 256 . 390 . 531 . 684 . 856 . 1.058 . 1.315 1.705 . 2.056 . 2.479 . 2.717 . 3.670 . 271 . 127 . 256 . 389 . 531 . 684 . 855 1.057 . 1.314 1.703 . 2.042 . 2.473 . 2.713 . 3.670 . 28 . 127 . 256 . 389 . 530 . 683 . 854 . 1.055 . 1.311 1.699 . 2.045 . 2.462 . 2.756 . 3.659 . 30 . 1.27 . 256 . 389 . 530 . 683 . 854 . 1.055 . 1.311 1.699 . 2.045 . 2.462 . 2.756 . 3.659 . 30 . 1.27 . 256 . 389 . 530 . 683 . 854 . 1.055 . 1.311 1.699 . 2.045 . 2.462 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.659 . 3.646 . 2.646 . 2.756 . 3.646 . 2.646 . 2.756 . 3.646 . 2.646 . 2.756 . 2.756 . 3.646 . 2.756 . 2.							,856	1.058	1,316	1,700				2.707
27 1.27 2.25 3.39 5.31 6.84 8.55 1.057 1.31 1.701 2.048 2.467 2.705 3.659 2.127 2.25 3.39 5.30 6.63 8.54 1.055 1.311 1.699 2.042 2.457 2.756 3.659 3.0 1.27 2.56 3.89 5.30 6.63 8.54 1.055 1.310 1.697 2.042 2.457 2.750 3.646 4.05 1.25 2.54 3.87 5.27 6.88 5.29 6.81 8.51 1.050 1.303 1.684 2.000 2.390 2.660 3.460 6.126 2.54 3.87 5.27 6.79 8.48 1.046 1.296 1.671 8.20 2.338 2.617 3.373 1.20 1.26 2.254 3.36 5.26 6.77 8.42 1.005 1.282 1.645 1.960 2.326 2.576 3.291		1.12/	.230	.390	.551	.00.				1 706	2.056			
28 1.27 2.56 3.89 5.51 6.83 8.55 1.056 1.31 1.699 2.045 2.452 2.750 3.645 2.501 1.27 2.56 3.89 5.30 6.83 8.54 1.035 1.311 1.699 2.042 2.457 2.750 3.646 2.501 1.27 2.55 3.88 5.29 6.83 8.54 1.035 1.310 1.697 2.042 2.457 2.750 3.646 2.501 2.50	26	127	250	200	531	684	,856	1.058	1.313	1 703	2.052			
29 1.27 .256 .389 .530 .683 .854 1.035 1.310 1.697 2.042 2.457 2.750 3.551 40 1.26 .255 .388 .529 .681 .851 1.050 1.303 1.684 2.000 2.390 2.660 3.450 60 1.26 .254 .387 .527 .679 .848 1.046 1.296 1.671 120 1.26 .254 .387 .527 .679 .845 1.041 1.289 1.653 .890 2.338 2.617 3.373 120 1.26 .254 .386 .526 .677 .842 1.036 1.282 1.645 1.960 2.326 2.576 3.291							.855	1.057	1.314	1 701	2.048	2.467		
10 127 236 389 530 683 854 1.035 1.010 1.020 2.021 2.423 2.704 3.551 1.020 1.020 2.020 2.020 2.620 3.620 1.020 2.020 2.020 2.620 3.620 2.020 2.020 2.020 2.620 3.620 2.0							.855	1.056	1.313	1.699	2.045	2.462		
10 127 236 389 530 683 854 1.035 1.010 1.020 2.021 2.423 2.704 3.551 1.020 1.020 2.020 2.020 2.620 3.620 1.020 2.020 2.020 2.620 3.620 2.020 2.020 2.020 2.620 3.620 2.0											2.042	2.457	2.750	3.040
40 126 255 388 529 .681 81 1.050 1.303 1.684 2.001 2.425 2.506 3.460 60 1.266 2.544 .387 .527 .679 8.45 1.041 1.289 1.653 1.800 2.328 2.617 3.373 120 1.26 2.54 3.86 5.526 .677 8.42 1.036 1.282 1.645 1.960 2.326 2.576 3.291							.854	1.055	1.310				2 704	3 551
126 125 254 387 527 679 848 1.046 1.289 1.658 1.980 2.358 2.617 3.373 120 1.26 2.54 .386 .526 677 842 1.051 1.282 1.645 1.960 2.326 2.376 3.291		.12/	.230	.307	,550			1.050	1.303	1.684		2,423		
120 .126 .254 .386 .526 .677 .845 1.041 1.220 1.645 1.960 2.326 2.376 3.251	40	.126	255	.388	.529									
126 .234 .386 .326 .326 .326 .326 .326	60	.126	.254	.387	.527								2 576	
∞ .126 .253 .385 .524 .674 .842 1.050	120	.126	.254	.386	.526	.677	.845	1.036	1,282	1.645	1.960	2.320	2.3.0	
	α	.126	.253	.385	.524	.674	.842	1,000				Bielen	ical Av	ricultural

Table D is adapted from Table III of Fisher & Yates: Statistical Tables for Biological, Agricultural and Madical Research, published by Oliver & Boyd Ltd., Edinburgh, and by permission of the authors and publishers.

unu publishers.

The lower percentiles are related to the upper percentiles which are tabulated above by the equation

The lower percentiles are related to the upper percentiles which are tabulated above by the equation 15df equals the negative of the 90th 15df equals the 15df

DISTRIBUTIONS	
CHI-SQUARE	
PO STIMO	

6.60	÷	6 635 10.827	-	-		_	_	_	18.475 24.322	_	_	-	_	24.725 31.264		_	_	_	÷	_	32.000 39.252	_	_	_	_		_	_	_	_	2000	_	_	700-6	_	_		_	
F	2	_		-	_	13,388 15.	_		16.622 18	_	_	-	_	22.618 24	_	_	_	_	-	_	29.633 32	_	_	_	_		_	_	_		40.270	_	_	42.856 45	_			_	н.
1	95	123	90	2815	0.488	070		12.592	14.067	15.507	16919	8 307		19 675	2010	200	705.77	23.685	24.996		26.296	_	_	_	_	_	17.5	25.01	276.55	32.175	36 415	37.652	_	38.885	_	_	_	_	
{	8	1	7,700	100	1770	2000		10.645	12017	53 565	14.50	1 0 0 0 1			200	18.04	19.812	21.064	22,307		23.542	25.55	24 080		3	28.412		29.615	30.813	32.007	33.196	34.382	_	35.563	36.741	37.916	10 087	20.00	10.23
	08	1	1.642	3.219	4.02	5,989	1.489	0 666	0.00	2000	200	777	3.44		50.4	15.812	16.985	18.151	10.1		70.466	200	20.07	27.70	23.90	25.038		26.173	27,301	28.429	29.553	30.675		31.795	12 012	34 027	36		30.430
	12		074	2,408	3,665	4.878	\$ 004		7.231	2,393	9.55	10.636	11.781		12 899	14 021	15.119	16 722	222	17.322	817 01	9.4	19.5	20.600	21.689	22.775		23.858	24,939	26 018	27.096	28.172		29.246	10	101		32.40	33,330
Percentile	9		.455	386	2,366	3,357	4,351		5,348	6.346	7,344	8.343	9,342		10.341	11.340	12 340			14 339		15,338	16 338	17.338	18,338	19.337		20.337	21,337	22 337	21.117	2117		25 116	25. 27.	200	27.730	28.330	29.336
Perc	١	3	2	713	424	2,195	3,000		3.828	4 671	5.527	6.393	7.267		8,148	0 034	9000	200	0.52	11.721		12.624	13.531	14.440	15.352	16.266		17.182	101		270	20.067	,,,,,,,	1 702		27.73	7	24.577	25.508
PERCENTILE POINTS OF CHI-SQUARE DISTRIBUTION		2	150			2	2,743		3.070	3 822	¥ 69 ¥	\$ 380	671.9		680 9	2003		3.034	9.467	10 307		11.152	12.002	12.857	11.716	14.578		16.445	71.7			100	20.740	0.00	20.00	20.703	21.388	22.475	23.364
N CHI-SO		=		.0138	7.	286		1.01	, ,00		200	2	7,000	1.00	6 470	200	6.304	7 042	7,790	8.547		9.312	0.085	10.865	17.641	2 443					4.040	200	10.473		767.71	18.14	18.939	19.768	20.599
POINTS C		s	-	.00393	10	.352		-	***	2000	2.10	7	3.355	2 340		4.575	\$ 226	5 892	6.57	7.261		1 963	6773	100						2.730	200	13.848	191	-	5.379	6.151	16.928	7.708	8.493
CENTILE				9000	20.0	185	.429	.752		*	300	2 032	2.532	3 059		3.609	4,178	4.765	891.	2002	_	_	_	_	-	2000	-	-	-	-	_	11.992	_	-!	=	=	14.847	-	908.9
		-		1	1000	1	762.	.554	_	.872	1.239	1.646	2 088	2.558	_	3 053	3.571	4 107		000	2.64.5	-	700	000	0	7.633	9,400	-	_	-	_	10.856	_	_	_	_	13.565	_	14041
2 10 4 2		1	ē		٠,	•	•			•	-	90	•	2		=	2	::	2:	•	2		9	-	8	<u>-</u>	R	-	=	=	-	7,	==	-	50	22	28	12	. 5

 $z=\sqrt{2z^4}-\sqrt{2df-1}$  is very nearly normally distributed with mean 0 and standard deviation 1. Ltd., Edinburgh, and by permission of the authors and provide .

If \( \gamma^2 \) is a chi-square variable with \( d \gamma \) greater than 30, then

523

ก็ก็ก็ก็ก็ก็ 20502

55555

54444

1955

94423

25445

27777

33377

55555 ដ្ឋាន្ធន

54535

33533 55555

**₹**5555

24244

\*\*\*\*\* \*\*\*\*\* 55555 \*\*\*\*\*\*

2442 24222 44442 24443

22,22,2

22222

22222 22652 44444 ก็ก็ก็ก็ก็

e

2

2

0

9

7

2 •

~

•

Cambridge he publisher, 3 Slometrika Trustees ž rermission of t ă Harrie), ò ٠ Pug Pearson ví ü 1962), eds. Ę 2 Spatfeliaba 5 Biomerifia Tables Reprinted fress.

322

	8	1018 39.50 13.90	2 2446	£ 22272	\$ 5555 5 555 5 2 13	188888	53458	
	921	39.49	2882	2.2.2.4 2.2.4.4 2.6.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.	2.55 2.246 2.22 2.23 2.23 2.23	2 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	855558	143
	8	39.48	1256.12	3.20 3.20 2.83 2.72	2 22 22 2 22 22 2	2 22222 2 22222 2 22222	2022	159
	2	19.47		3.56 3.06 2.79 2.79			25555	201 1.74 1.61 1.61
	30	1901 39.46 14.08	1252	1282			2213	1692
	7	997.2 39.46 14.12	25.24.2 21.14.2	E E E E E	7 7277			2014 1.38 1.76 1.76
	20	993.1 39.45 14.17	37.49	4262	2222			122202
	n	984.9 39.43 14.25		25.50 25.50 20.50				126621
	n	976.7 39.41 14.34 8.75		25555				12222
centiles	2	39.40 39.40 8.84						1227
97.5th percentiles	۰	25.55 25.55 25.55 25.55						12222
•		956.7 39.37 14.54 8.98	5.60 4.43 6.43					22422
	-	948.2 39.36 14.62 9.07			3,229	P.H.H.H.		2222
	•	937.1 39.33 14.73 9.20	86.28 8.11.4 8.11.8 8 8.11.8 8 8.11.8 8 8 8		3228			42224
	s	921 8 39.30 14.88 9.36	7.15 5.29 4.82 4.82	4.24 4.04 7.7.6 3.66	8.55 8.55 8.55 8.55 8.55 8.55			22222
	+	892.6 39.25 15.10 9.60	52225 82225 82225 8	74.44 1.00 1.89 1.89	33.23			22.22.2
	•	864.2 39.17 15.44 9.98	5.60 5.89 5.89 5.08	24444 24444 24444	24.4.0 21.95 31.90 31.90 31.90			1.50 3.23
	"	39 00 16.04 10.65	2.43 2.43 2.43 2.43 2.43	22 24 23 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	04482 74444 76632	44444		2525 2525
		647.8 38.51 17.44 12.22	10.01 12.7 12.7 12.7	25349 25348	2286620			
	<u>,                                     </u>	-4n-4	W&L@0	22222	22722	នដដងដ ដ	2222	888

8	199.5 199.5 41.83 19.32	12.14 8.88 7.08 5.93 5.93	22822	21.25 21.25	2-224		, ,,
170	25.5 25.5 25.5 25.5 25.5 25.5 25.5 25.5	72.71 9.00 6.00 5.00 8.00 8.00	25.4 10.4 10.4 10.4 10.4 10.4 10.4 10.4 10	HH 1	2000 2000 2000 2000		
\$	25.55 19.55 19.55 19.55	677.25	444E			•	
\$	848 848 11.00 87.00	22462		84505			
30	15 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	25.9 25.9 25.9 25.9 25.9 25.9 25.9 25.9		25.00			
7.	24940 199.5 42.62 20.03	12.78 7.65 6.50 5.73		3.54			
92	88.45 4.65 4.15	9.59 9.59 8.61 8.61 8.81		3.50			
52	8825 483	981 981 681 603	25.444 25.444				2.78
2	48.45 4.65	818 200 100 100 100 100 100 100 100 100 100	25.24.4 25.24.4 26.44.4	3.97 3.97 3.86			234 3 18 250 2 95 271 2 24 2 52 2 36
01	1224 199.4 13.69	2016 2018 1176 1176	25.24.4 25.00.44		3,50		22222
6	24091 199.4 21.14	13.77 10.39 13.8 13.8 6.54	5.55 4.94 4.94 5.74		3.96		SEC.
	23925 19924 44 13 21,35	13 96 10.57 7.50 6 69	5.12 5.68 5.35 8.88 8.88 8.88	25.5.4 25.5.4 25.5.4 25.5.4 25.6.4 25		94 3.78 89 3.73 85 369 77 365	¥2988
1	55 4 4 4 6 4 4	4101 0.79 0.79 0.89 0.89 0.89 0.89		28 4 4 4 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5		115 110 110 110 110 110 110 110 110 110	21.446 21.446
9	1993 1993 11.97	14.51 11.07 7.95 7.95	6.54 6.10 5.76 5.48 5.48	5 07 4 4.91 4 566	44444 2233	4444	25 5 25 E
s	25 53 3 2 45 33	26.51 8.52 7.47	66.42 5.79 5.79 5.50	5.37 5.07 4 4 8 8 8 8 8	4444 6864 864 864	44444	45.50
4	250 1992 23.15	15.56 12.03 10.05 7.96	25.58 25.59 4.88 5.55 5.55 6.69	5.64 5.54 5.37 5.37	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	23.05 23.05 23.05 23.05 23.05 23.05 23.05 23.05 23.05 23.05 24.05 25.05	444mg
r.	615 1992 24.44 24.44	16.53 12.92 10.88 9.60 8.72	7 508 7 7 7 50 6.93 6.83	6.48 6.30 6.03 5.92	5.32 5.53 5.58 5.58 5.58	0 5 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	25.00 25.00
7	9900 1990 1990 1990	12.40 12.40 11.04	8.51 8.51 7.92	07.7 2.7.7 12.7 12.7 10.0	533 6 899 533 6 811 53 6 6 81	29 248 20 241 20	9.18 8.49 7.88 1.88
-	16211 198 5 55 55 31.33	22.78 18 63 16.24 14 69	12.83 11.75 11.37	0 20 20 0 20 20 0 20 20 0 20 20 0 20 20 0 20 20 0 20 20 0 20 20 0 0 0	99999 488698 488688		
£ /		N 0 1- 10 0	2222	25 57 88 81 81 88 81	នដដដង	22 22 22	88888

## PERCENTILE POINTS OF STUDENTIZED RANGE, q, DISTRIBUTIONS FOR / AND v DEGREES OF FREEDOM\*

TABLE E

90th Percentiles

· ·	2	3	4	5	6	7	8	9	10
1	8.929	13.44	16.36	18.49	20.15	21.51	22.6	23,62	24.48
2	4.130	5.733	6,773	7.538					
3	3,328	4.467	5.199	5,738					
4	3.015	3.976	4.586	5.035	5.38				
5	2.850	3.717	4.264	4 664					
6	2.748	3.559	4.065	4.435					
7	2.680	3.451	3.931	4.280		4.780			
8	2.630	3,374	3.834	4 169					
ġ	2.592	3.316	3,761	4,084					
10	2.563	3.270	3.704	4.018				1 4 87.	3 5 007
		1	1	1		1	4.63	6 4.783	3 4.913
11	2.540	3.234	3.658	3.965					
12	2.521	3.204	3 621	3.922	4.156			1 4 652	
13	2.505	3.179	3.589	3.885	4,116			4 4,602	2 4.724
14	2.491	3.158	3.563	3.854	4.081	4.267		4 4.560	4.680
15	2.479	3.140	3.540	3.828	4.052	4.235			
16	2.469	3.124	3.520	3.804	4 026	4.207	4.360	4,492	4.608
17	2.460	3.110	3.503	3.784	4.004	4.183	4.334	4.464	4.579
18	2.452	3.098	3.488	3.767	3.984	4.161	4.311		4.554
19	2.445	3.087	3.474	3.751	3 966	4.142	4.290	4.418	4.531
20	2.439	3.078	3.462	3,736	3.950	4.124	4.271	4 398	4.510
24	2.420	3.047	3.423	3.692	3.900	4.070	4,213	4.336	4.445
30	2.400	3.017	3.386	3.648	3.851	4.016	4.155	4.275	4.381
40	2.381	2.988	3.349	3.605	3.803	3.963	4.099	4.215	4.317
60	2.363	2.959	3.312	3.562	3.755	3.911	4.042	4.155	4.254
120	2.344	2.930	3.276	3.520	3.707	3 859	3.987	4.096	4.191
œ	2.326	2.902	3.240	3 478	3 661	3.808	3.931	4.037	4.129
·	11	12	13	14	15	16	17	18	19
1	25.24	25.92	26.54	27,10	27.62	28.10	28.54	28.96	29.35
2 3 4 5 6	10.01	10.26	10.49	10 70	10.89	11.07	11.24	11.39	11.54
3	7.487	7.667	7.832	7.982	8 120	8.249	8.368	8.479	8.584
4	6.495	6.645	6.783	6.909	7.025	7.133	7.233	7.327	7.414
5	5.966	6.101	6.223	6.336	6.440	6.536	6.626	6.710	6.789
6	5.637	5.762	5.875	5.979	6.075	6 164	6.247	6 325	6.398
7 8	5.413	5.530	5.637	5.735	5.826	5.910	5.988	6.061	6 130
8	5,250	5.362	5.464	5 558	5 644	5.724	5 799	5.869	5.935
9	5,127	5.234	5.333	5.423	5.506	5.583	5.655	5.723	5.786
10	5,029	5,134	5.229	5.317	5.397	5 472	5.542	5.607	5.668
11	4.951	5.053	5.146	5.231	5.309	5.382	5.450	5.514	5.573
12	4,886	4.986	5.077	5.160	5.236	5 308	5.374	5.436	5.495
13	4.832	4.930	5.019	5.100	5.176	5.245	5.311	5.372	5 429
14	4,786	4 882	4 970	5 050	5.124	5.192	5.256	5 316	5.373
15	4.746	4.841	4.927	5.006	5.079	5.147	5.209	5.269	5.324
16	4.712	4.805	4 890	4.968	5.040	5.107	5.169	5.227	5.282
17	4.682	4.774	4.858	4.935	5 005	5.071	5.133	5.190	5.244
18	4.655	4.746	4.829	4 905	4.975 4.948	5 040 5.012	5.101 5.073	5.158 5.129	5.211 5.182
		4.721	4.803	4.879 4.855	4.948	4.987	5.073	5.103	5.182
19	4.631								
20	4,609	4.699	4.780			1		1	
20 24	4,609 4,541	4.699 4.628	4,708	4,780	4.847	4.909	4 966	5.021	5.071
20 24 30	4,609 4,541 4 474	4.699 4.628 4.559	4.708 4.635	4.780 4.706	4.770	4.830	4.886	4.939	4 988
20 24 30 40	4.609 4.541 4.474 4.408	4.699 4.628 4.559 4.490	4.708 4.635 4.564	4.780 4.706 4.632	4.770 4.695	4.830 4.752	4.886	4.939 4.857	4 988 4,905
20 24 30 40 60	4,609 4,541 4 474 4 408 4,342	4.699 4.628 4.559 4.490 4.421	4.708 4.635 4.564 4.493	4.780 4.706 4.632 4.558	4.770 4.695 4.619	4.830 4.752 4 675	4.886 4.807 4.727	4.939 4.857 4.775	4 988 4.905 4.821
20 24 30 40	4.609 4.541 4.474 4.408	4.699 4.628 4.559 4.490	4.708 4.635 4.564	4.780 4.706 4.632	4.770 4.695	4.830 4.752	4.886	4.939 4.857	4 988 4,905

Reproduced from H. Leon Harter, "Tables of range and studentized range," Annals of Mathematical Statistics, 31 (1960), 1122–47, by permission of the author and editor,

"In the one-factor ANOVA with n observations in each of 2 groups, v = l(n-1). In general, v is the number of degrees of freedom for the mean-square "within" in an analysis of variance.

	J	
	8	20114 111121 20144 444000 200000 11014000 11014000 1101400 1101400 1101400 1101400 1101400 1101400 1101400 110
	#	South 42775 Goods 44440 Edges Budge 44467
	\$	2014 Vedes 12124 44444 64444 412846 201384
	\$	2010 - 1000 - 10
		STATE STATE
	7	Month Assista Peens was at the Control of the Contr
	2	**************************************
	=	And do-or stood winds attach applied being being applied by the being applied by the being
	=	400 82445 40000 80000 80000 82484 54444 84464 84
09.518 percentites		SULA MERCIA GUINA GARANTI BERNIN STAND STAND STAND STAND STAND STAND STAND STAND STANDS STAND
PR.SUR P.		SULA HEARING MELAND ANNING MINERS LANGE BYIGHO AND ANNING MINERS CANNOT WILL AND ANNING WILL ANNING ANNING MINERS ANNING MINERS CANNOT WILL ANNING ANNING MINERS CANNOT MI
		SOLA MISTRO OSCINO GONDO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SECOS SECONO SE
		The stands of th
		SEC SULLE CORPLE PRESS CASUL SELECT MOUNTS
	1	THE COURT PROPERTY OF THE PROP
ŀ		DE CHIEFE TOOOR PETET COOR SALES ESTAD
-	7   200	THE CULTUTE CASES OFFICE RECENT ASSESSMENT A
cont.)	2 0000	
FABLE E (cont.)	7	TO ARRIED STREET
Z /	-	- werer seres seres seres seres sesse (
52	8	

97.5th Percentiles

				97.5th	Percentiles					
* \	2	3	4	5	6	7	8		,	10
1	35.99	54.00	65.69	74.2	80.87	86.2	90.	06 04	77	
2	8.776	11.94	14.01	15.54	16.75					98.20
ã	5,907	7.661	8,808							19.95
4	4.943	6.244								12.14
- 7	4.474	5.558	7.088 7.716						279	9.548
5	4.199		6.257						068	8.291
7		5.158	5.772						359	7.554
8	4.018	4.897	5.455						895	7.072
	3.892	4.714	5.233				69 6.3	82 6.:	568	6.732
. 9	3.797	4.578	5.069				50 6.1	51 6.3	325	6.479
10	3.725	4.474	4.943	5.28	7 5.55	5.7	82 5.9	72 6.1	138	6.285
11	3.667	4.391	4.843	5.17						6.130
12	3.620	4.325	4.762	5.08						6 004
13	3.582	4.269	4.694	5.00						5.900
14	3,550	4.222	4.638	4.94		3   53	74 5.5	40 5.6	84	5 811
15	3.522	4.182	4.589	4.88				71 5.6	12	5 737
16	3.498	4.148	4.548	4.838	8 <b>5.</b> 06€	5.23	53 5.4	12 5 5	50	5.672
17	3.477	4.118	4.512	4.797	7 5.020	5.20	)4 \ 5.3	61   54	96	5 615
18	3.458	4.092	4.480	4.761	4.981					5 565
19	3.442	4.068	4.451	4.728	4.945	5.12		75 5.4	05	5.521
20	3.427	4.047	4.426	4.700						5.481
24	3.381	3.983	4.347	4.610	4.816					5.358
30	3.337	3.919	4.271	4.523		4.88	1 5.01	7 5.1	34	5.238
40	3.294	3.858	4.197	4.439	4.627	4.78	0   491	0 50	22	5.120
60	3.251	3.798	4.124	4 356	4,536	4.68	2 480	6 4.9	12	5.006
120	3.210	3.739	4.053	4.276		4.58	7 4.70		D5 i	4.894
œ	3.170	3.682	3 984	4 197	4.361	4.49		5 4.70	90   0	4.784
				·						
\ J		l		l '		1	1		ı	
,	11	12	13	14	15	16	17	18	ĺ	19
1	101.3	104.0	106 5	108.8	110.8	112.7	114.5	116.2	117	
2	20 52	21.03	21.49	21.91	22,30	22.67	23.01	23,32		.62
2 3 4 5 6 7 8	12 46	12.75	13.01	13.26	13.48	13.69	13 88	14.06		.23
4	9.788	10.01	10 20	10.39	10.55	10.71	10.85	10 99		.11
5	8,490	8.670	8.834	8.984	9,124	9 253	9.374	9.480		.593
6	7,729	7.887	8.031	8.163	8.286	8.399				698
7	7,230	7.373	7.504	7 624	7.735	7.839	7.935			.111
8	6 879	7.011	7,132	7.244	7.347	7.443			5   7.	.695
9	6.617	6.742	6.856	6.961	7.058	7.148			[ 7.	.385
10	6.416	6.534	6.643	6.742	6.834	6.920	7.000	7.075	7.	.146
11	6 256	6.369	6.473	6.568	6.657	6.739	6.815	6.887	6.	955
12	6.125	6,235	6.335	6 427	6.512	6.591	6 665	6.734	6.	799
13	6.017	6.123	6.220	6.309	6.392	6 458	6.539	6.607	6.	670
14	5.926	6.029	6 123	6.210	6.290	6 364	6 434	6.499		560
15	5,848	5 949	6.041	6.125	6.203	6.276	6.344	6.407		467
16	5.781	5.879	5.969	6 052	6 128	6.199	6 265	6 328	6.	386
17	5.722	5 818	5 907	5.987	6 062	6.132	6.197	6.258		315
18	5.670	5.765	5 852	5.931	6.004	6.073	6.137	6.197		253
19	5.624	5.718	5.803	5.881	5.954	6 020	6.083	6.142		98
20	5.583	5.675	5.759	5.836	5.907	5.974	6 036	6.093	6.1	48
24	5.455	5 543	5.623	5.697	5.764	5.827	5.886	5.941 5.792	5 9	
30	5.330	5 414	5,490	5.560	5.624	5.684	5.740	5.792 5.646	5.6	
40	5 208	5.288	5.360	5.426	5 487	5.544	5.597 5.456	5.503	5.5	
60	5.089	5.164	5.232	5.295	5.352	5.406 5.271	5.436	5.362	5.4	03
120	4 972	5 043	5.107	5.166	5.221	5.139	5.183	5.224	5.7	62
× (	4.858	4 925	4.985	5.041	3.092	3.139	2.103			

				your Fert	centiles				
· · ·	2	3	4	5	6	.7	8	9	10
1	17.97	26 98	32 82	37.08	40 41	43.12	45.40		
2 3 4	6 085	8.331	9.798	10.88	11,74	12,44	45.40	47.36	49.07
3	4.501	5.910	6.825	7,502	8.037	8.478	13.03	13.54	13.99
4	3 927	5 040	5.757	6.287	6.707		8.853	9.177	9.462
5	3 635	4 602	5.218	5 673	6.033	7.053 6.330	7.347	7.602	7.826
6	3.461	4.339	4 896	5.305	5.628	5.895	6.582	6.802	6.995
7	3.344	4.165	4 681	5.060	5.359		6.122	6.319	6 493
8	3.261	4.041	4.529	4.886	5.167	5.606	5 815	5.998	6.158
9	3.199	3,949	4.415	4.756	5.024	5.399	5 597	5.767	5.918
10	3.151	3.877	4.327	4 654	4.912	5.244	5.432	5.595	5.739
				. 034	4.512	5.124	5.305	5.461	5.599
11	3.113	3.820	4.256	4 574	4 823	5.028	l		1
12	3.082	3.773	4.199	4.508	4.751		5.202	5,353	5 487
13	3.055	3.735	4.151	4.453	4.690	4 950	5.119	5.265	5.395
14	3.033	3.702	4.111	4,407	4.639	4.885	5.049	5.192	5.318
15 16	3 014	3.674	4 076	4.367	4 595	4.829	4.990	5.131	5.254
17	2.998	3.649	4.046	4.333	4.557	4.782 4.741	4.940	5.077	5.198
18	2.984	3 628	4.020	4.303	4.524	4.741	4.897	5.031	5.150
19	2 971 2,960	3 609	3.997	4.277	4.495	4.705 4.673	4.858	4.991	5.108
20		3.593	3.977	4.253	4 469		4.824	4.956	5 071
20	2 950	3.578	3 958	4 232	4.445	4.645 4.620	4.794	4.924	5 038
24	2.919		i		1 7.773	4.620	4.768	4.896	5.008
30	2.888	3.532	3.901	4.166	4,373	4.541			
40	2 858	3.486	3.845	4.102	4.302	4.341	4.684	4.807	4.915
60	2.829	3.442	3.791	4.039	4.232	4.389	4.602	4.720	4 824
120	2.800	3 399 3 356	3.737	3.977	4 163	4.389	4.521	4.635	4.735
00	2,772	3.314	3 685	3.917	4 096	4.241	4.441	4.550	4.646
	1	3.314	3 633	3.858	4.030	4.170	4.363	4.468	4.560
						4.170	4.285	4.387	4.474
\ 1	111								
"	1 "	12	13	14	15			T	
	<b>\</b>	<del> </del>		1	1.3	16	17	18	19
1	50.59	51.96	53.20	54.33				,	, ,,
2 3 4	14.39	14.75	15 08	15.38	55.36	56.32	57.22	58,04	
3	9 717	9 946	10.15	10 35	15 65	15.91	16.14	16.37	58.83
- 3	8.027	8.208	8.373	8.525	10.53	10.69	10 84	10.37	16.57
5 6 7	7.168 6 649	7.324	7.466	7.596	8 664	8,794	8.914	9.028	11.11
ž	6.302	6.789	6 917	7.034	7.717	7.828	7,932	8.030	9.134 8.122
8	6 054	6 431	6.550	6 658	6.759	7.244	7.338	7.426	7.508
9	5 867	6.175 5 983	6.287	6.389	6 483	6.852	6.939	7.020	7.097
10	5.722	5.833	6 089	6.186	6.276	6 571	6.653	6.729	6.802
	1	3.033	5.935	6 028	6.114	6 359	6.437	6.510	6.579
!!	5 605	5,713	5 811		1	6.194	6.269	6.339	6.405
12	5.511	5.615	5.710	5 901	5 984				0.403
13 14	5.431	5.533	5 625	5.798	5.878	6 062 5.953	6.134	6.202	6.265
15	5 364	5 463	5 554	5.711	5.789	5.862	6.023	6.039	6.151
16	5.306	5 404	5 493	5 637	5.714	5.786	5.931	5.995	6.055
iř	5.256 5 212	5.352	5.439	5.574 5.520	5 649	5.720	5 852	5.915	5,974
18	5 174	5.307	5 392	5.471	5.593	5 662	5.785	5.846	5.904
19	5.140	5 267	5.352	5 429	5.544	5 662 5 612	5.727 5 675	5.786	5.843
20	5 108	5 267 5.231 5.199	5.315	5.391	5.501	5,568	5.630	5.734	5.790
	3103	3.199	5 282	5.357	5 462	5 528	5.589	5.688	5,743
24	5 012	5 099	Ι.	1	5.427	5 493	5 553	5.647 5.610	5.701
30 40	4 917	5 001	5.179	5.251	5.319		- 333	3.610	5.663
	4 824		5 077	5.147	5.211	5.381	5,439	5.494	
120	4 732	4 904	4 977	5 044	5.106	5 271	5.327		5.545
120	4 732	4.714	4 878	4 942	5.106	5 163	5 216	5.379	5.429
	4 732	4 805	4.781	4 942 4 842	5.106 5.001 4 898	5 163 5 056	5 216 5.107	5.379 5.266	5.429 5.313
120	4 732	4.714	4 878	4 942	5.106	5 163	5 216	5.379	5.429

		_		99.31	a Percent	ues					
· · ·	2	3	4	5		6	7	8	$\top$	9	10
1	180.1	270.1	328.5	371.2	404	_				_	
2	19.93	26.97	31.60			. <del>4</del> .73	431.6	454			491.1
3	10.55	13.50	15.45				39.95			.46	44 89
4	7,916	9.814	11.06	11.9		.06	19.01			.53	21.15
- 3	6.751					.74	13.35			.33	14.74
5 6 7	6.105						10.88			.63	11.93
7	5.699					.135	9.52			.14	10 40
8	5.420				5 8.	339	8 67			211	9.433
9	5.218					797	8.09		54   8.	578	8.777
10				7 7.07	4 7.	405	7.68	0 7.9		120	8.303
	5.065	1	6 41:	6.80	0 7.	109	7.36	5 7.5	84 7.	775	7.944
22	4.945	5.727	6.22	2 6.58	8 6.	878 l	7.119	7.3	25 7.	505	7.664
12	4.849	5.597	6.068	6.41	6 6.	693	6,922			288	7.439
13	4.770	5.490	5.943	6.27	7 6	541	6.760			iii l	7.255
14	4,704	5.401	5.838			114	6.626			62	7.101
15	4.647	5.325	5.750			808	6 511			37	6.971
16	4,599	5.261	5 674	5.97			6.413				6.859
17	4.557	5.205	5.608				6.329	6.49		36	6.763
18	4.521	5.156	5.550				6.255	6.41			6,678
19	4,488	5.113	5,500	5.783		05 /	6.189	6.34	6 4.4		6.603
20	4.460	5.074	5.455	5.732			6 131	6.28			6.537
24	4,371	4.955	5.315	5.577	5,71	/	5.952	6 096	6.2	. 1	6.332
30	4.285	4 841	5.181	5.428	5.6		5.780	5 914			6.135
40	4.202	4.731	5.053	5.284	5.40		5.614	5.739			5,944
60	4.122	4.625	4.928	5.146	5.31		5 454	5.571	5.67		5.762
120	4.045	4.523	4.809	5.013	5.17		5.301	5.410		3	5.586
20	3,970	4.424	4.694	4,886	5,03		5.154	5.255	5.34	71	5.418
	3.570	4.424	4.074	4.000	3,03		2.134	3.233	3.54	ــــــــــــــــــــــــــــــــــــــ	3.410
\ ,	1			í	1	ſ	1		1	- (	19
<u>"</u>	11	12	13	14	15		16	17	18	_ _	19
1	506.3	520.0	532.4	543.6	554.0	1 5	63.6	572.5	580.9	1 5	88.7
,	46.16	47.31	48.35	49.30	50.17	1 -	50.99	51.74	52 45		53.12
3 4	21.70	22,20	22.66	23.08	23.46		23 82	24.15	24,46	1	24.76
4	15,10	15.42	15.72	15.99	16.24		16 48	16.70	16,90		7.09
Ś	12.21	12.46	12.69	12.90	13.09		13.27	13,44	13.60		13.75
5 6 7 8	10.63	10.83	11.02	11.20	11.36	- { i	11.51	11.65	11.78	1.	1.90
7	9,632	9.812	9,977	10 13	10 27		10.40	10 52	10.64		0 75
8	8.955	9.117	9.265	9.401	9,527		9.644	9.754	9.857		9.953
9	8,466	8.614	8.749	8.874	8,990	1	9.097	9.198	9.292		9.381
10	8.096	8.234	8.360	8.476	8.583		8.683	8.777	8.865		8 947
11	7,807	7.937	8.055	8.164	8.265	1	8.359	8.447	8,530	1	8 608
12	7.575	7.697	7.810	7.914	8.009		8.099	8.183	8.261	i :	8.335
13	7,384	7.502	7,609	7,708	7.800		7.886	7.965	8,040	J i	8.111
14	7.225	7.338	7,442	7.537	7.625	1 :	7.707	7.784	7.856	1	7.924
15	7.091	7,200	7.300	7.392	7,477	1 :	7.556	7.630	7.699	1 1	7.765
16	6.976	7.081	7,178	7.267	7 349	1 1	7.426	7.498	7.566		.629
17	6.876	6.979	7.072	7.159	7.239	1 7	7.314	7.384	7,449	1 7	.511
18	6.788	6.888	6.980	7.064	7.142	f 7	7.215	7.283	7.347		.407
19	6.711	6.809	6.898	6.981	7.057	1 2	7.128	7.195	7,257		.316
20	6 642	6.738	6.826	6.907	6.981	7	7.051	7.116	7.177		.235
24	6.431	6.520	6.602	6.677	6.747	6	.812	6.872	6 930		983 741
30	6.227	6.310	6.387	6.456	6.521	6	.360	6.638	6.691		507
40	6.030	6.108	6.179	6.039	6.304		.146	6.194	6.239	6	281
60	5.841	5.913	5.979 5.786	5.842	5 893		940	5.984	6.025	6	064
120	5 660	5.726 5.546	5.602	5,652	5,699		742	5.783	5.820	5	856
- 00	5.485	3.340	3.002	2,032	2.077				20020		

INDEE . (				95	th Perce	ntiles					
51	2	3	T	4	5	6	7		8	9	10
			-1		35.6	202.2	215.8		27.2	237.0	245.6
1 1	90.03	135.0	16		24.72	26.63	28.2		29.53	30.68	31.69
2 3 4 5 6 7 8	14.04	19,02			13.33	14.24	150		15.64	16.20	16.69
1	8.261	10 62				10 58	11.1	~	11.55	11.93	12.27
4	6.512	8.13		9,173	9.958	10 36	111.	. 1	9.669	9,972	10.24
i i	5.702	69		7.804	8 421	8.913	9.3		8.613	8.869	9.097
,	5.243	6.3		7.033	7.556	7.973	8.3		0.013		8.368
*	4,949	5.9	9	6.543	7.005	7.373		79	7.939	8.166	
:	4,746	56	15 ]	6,204 (	6.625	6.960	7.2	237	7.474	7.681	7.863
ŝ	4 596	5.4	28	5.957	6.348	6 658		15	7.134	7.325	7.495
10	4.482	5.2	70	5.769	6.136	6.428	6.6	569	6.875	7.055	7.213
11	4,392	5.1	46	5.621	5 970	6 247		476	6 672	6.842	6.992
12	4.320	5.0	46	5.502	5.836	6.101		321	6.507	6 670	6.814
13	4 26	4.	64	5.404	5.727	5.981	6.	192	6.372	6.528	6.667
14	4.21	1 4	95	5.322	5.634	5.881	6.	085	6.258	6.409	6.543
15	4.16	8 4.	336	5,252	5.556	5.796	) 5.	994	6.162	6.309	6.439
16	4.13	Ll 4.	186	5.192	5 489	5.722	1 5.	915	6.079	6.222	6.349
17	4 09	9   4.	742	5.140	5.430	5.659	5.	847	6.007	6.147	6.270
18	4.07	1 4	703	5 094	5.379	5 603	5.	788	5.944	6 031	6.201
19	4.04	6 4.	670	5 054	5.334	5.554	5.	735	5.889	6.022	6.141
20	4 02	4 4	639	5 018	5.294	5.510		688	5.839	5.970	6.087
24	3 9	6 4	546	4.907	5.168			.542	5.685	5.809	5.919
30	3.8	9 4	455	4.799	5 048		: ) 5	.401	5.536	5.653	5.756
40	3.8	25   4	367	4.696	4.931	5.114	1 ( 5	.265	5.392	5.502	5.599
60	3.7	62   4	282	4.595	4.818	4.99		.133	5.253	5.356	5.447
120	3.7	02 4	.200	4,497	4.709			.005	5.118	5.214	5.299
80	3 6	43   4	.120	4.403	4 60:	4.75	7 ( 4	882	4.987	5.078	5.157
,	1	11 12		13	14	15	-	16	17	18	19
1	253.	2 26	00	266,2	271.8	277.0		1.8	286.3	290.4	294.3
2	32.	59   3	1.40	34.13	34.81			5.00	36 53	37.03	37.50
3	17.		7.53	17,89	18.22	18.52	1 1	8.81	19.07	19.32	19.55
4	52.	57   1	2.84	13 09	13.37		1 1	3.73	13.91	14.08	14.24
	10	48   1	0.70	10 89	11.08	11.2		1.40	11.55	11.68	11.81
3		301	9 485	9 653	9.80	9.9		0.05	10 21	10.32	10.43
		548 027	8.711	8 860	8.99	7 9.1	24	9.242	9.353		9.554
	.   .	027	8.176	8,312	8.4	36 8.5 25 8.1	52	8 659	8.760		8,943
	9 1	356	7.485	7.910	80	25   8.1	22 )	8.232	8.32	8.412	8.495
	1	1		7,603	7.7	1	12	7.906	7.99	8 076	8.153
	!! ! !	.128	7.250	7,362	7.4	65 7.5	60 l	7.649	7.73	7.809	7,883
	2	5 943 5,791	7.060	7.167	7.2	65 7.3	56	7,441	7.52	7,594	7,665
	2 1	5.791	6 903	7.006	7.1	01 7.1	83	7.269	7.34	5 7.417	7,485
	14 15 16 17	6 664	6.772	7.167 7.006 6 871 6.737	6.9	62 7.0	47	7,126	7.19	7,268	7,333
	13	6.555	6 660	6.737	6 8	45 1 6.9	27 }	7.003	7.19 7.07	7,142	7,204
	!2	6 462	6.564	1 0008		44 68	23	6.898	6.96 6.87 6.79 6.71	7,032	
	16	6 381	6 480	6.572		56 6.1	134	6 806	6.87	6,937	6 997
	: 1	6.310	6 407	6 491	6:	579 61	555	6.725	6.79	2 6.854	6.912
	19	6 247	6 342	6 430		10 6.	555 585	6 654	6.71	9 6.780	6.912 6.837
	1		6 285	1	6.	630	523	6.591	6 65	4 6.714	6.771
	24	6017	6 106		5 6.	261 6	330	6.394	6.45	3 6,510	6,563
	10	5 847	5 9 3 2	600	6 6	078 6	143	6.203	6.2		
	42 )	5 686	5 764		5 5.		961	6 017	60		6,165
	120	5 528 (	2 601	5 66	7 5.		785	5 837	5.8		5,103
	ac	5 375	5 44		5 5.	562 5	614	5 662	5.7	18 5.75	5.974 3 5.790
		* +11	5 29	) 534	<b>5</b>	400 5	448	5.493	5.5	5.57	5.611

TABLE H\* DETERMINATION OF r<sub>in</sub> FOR VARIOUS VALUES OF beford OR adjbe FROM A FOUR-FOLD CONTINGENCY TABLET

r	$\frac{bc}{ad}$ or $\frac{ad}{bc}$	Fiet	$\frac{bc}{ad}$ or $\frac{ad}{bc}$	ree	$\frac{bc}{ad}$ or $\frac{ad}{bc}$	reet	bc ad	or ad
0	1.000	.26	1.941-1.993	.51	4.068-4.205	.76	11.513	-12.177
.010	1.013-1.039	.27	1.994-2.048	.52	4.206-4.351	.77	12.178	-12.905
.02	1,040-1.066	.28	2.049~2.105	.53	4.352-4.503	.78	12.906	-13.707
.03	1.067-1.093	.29	2.106-2.164	.54	4.504-4.662	.79	13.708	-14.592
.04	1.094-1.122	.30	2,165-2,225	.55	4.663-4.830	.80	14.593-	-15.574
.05	1.123-1.151	.31	2.226-2.288	.56	4.831-5.007	.81	15.575-	-16.670
.06	1.152-1.180	.32	2.289-2.353	.57	5.008-5.192	.82	16.671-	-17.899
.07	1.181-1.211	.33	2.354-2.421	.58	5.193-5.388	.83	17.900	19.287
.08	1.212-1.242	.34	2,422-2,491	.59	5.389-5.595	.84	19.288-	20.865
.09	1.243-1.275	.35	2.492-2.563	.60	5.596-5.813	.85	20.866-	22.674
.10	1.276-1.308	.36	2.564-2.638	.61	5.814-6.043	.86	22.675~	24.766
.11	1.309-1.342	.37	2.639-2.716	.62	6.044-6 288	.87	24.767-	
.12	1.343-1.377	.38	2.717-2.797	.63	6.289-6.547	.88	27.213-	30.105
.13	1.378-1.413	.39	2,798-2.881	.64	6.548-6 822	,89	30.106-3	
.14	1.414-1.450	.40	2.882-2.968	.65	6.823-7.115	.90	33.578-3	
.15	1.451-1.488	.41	2.969-3.059	.66	7.116-7.428	.91	37.816-4	
.16	1.489-1.528	.42	3.060-3.153	.67	7.429-7.761		43.097-4	
.17	1.529-1.568	.43	3.154-3.251	.68	7.762-8.117		49.847-5	
.18	1.569-1.610	.44	3.252-3.353	.69	8.118-8.499		58.759-7	
.19	1.611-1.653	.45	3.354-3.460	.70	8.500-8.910		71.036-8	
.20	1.654-1.697	.46	3.461-3.571	.71	8.911-9.351		88.965-1	
.21	1.698-1.743	.47	3.572-3.687	.72	9.352-9 828		17.480-1	
.22	1.744-1.790	.48	3.688-3.808	.73	9.829-10.344		59,504-29	
.23	1.791-1.838	.49	3.809-3.935	.74	10.345-10.903		2.865-92	
.24	1.839-1.888	.50	3.936-4.067	.75	10.904-11.512	1.00 92	23.688-	00
.25	1.889-1.940		- 1		ì			


Values in this table were calculated by Thomas O. Maguire.
 † If bc/ad is greater than 1, the value of r<sub>ui</sub> is read directly from this table. If ad/be is greater than 1, the table is entered with ad/be and the value of r<sub>ui</sub> is negative.

TABLE G. FISHER'S Z-TRANSFORMATION OF ray

11311									
,	Z,	,	2,	<i>r</i>	z,	,	2,	r	z, ———
.000	,000	.200	.203	.400	.424	.600	.693	.800	1.099
.003	.005	.205	.208	.405	.430	.605	.701	.805	1.113
.010	.010	,210	.213	.410	.436	.610	.709	.810	1.127
.015	.015	.215	.218	.415	.442	.615	.717	.815	1.142
,020	.020	.220	.224	.420	.448	.620	.725	.820	1.157
.025	.025	225	.229	.425	.454	,625	.733	.825	1,172
.030	.030	.230	.234	.430	.460	.630	.741	.830	1.188
.035	.035	.235	.239	.435	.466	.635	.750	.835	1,204
.040	.040	,240	.245	.440	.472	.640	.758	.840	1.221
.045	.045	.245	.250	.445	.478	.645	.767	.845	1.238
.050	.050	.250	.255	.450	.485	.650	.775	.850	1.256
.055	.055	.255	.261	.455	.491	.655	.784	.855	1.274
.060	.060	.260	.266	.460	.497	.660	.793	.860	1.293
.065	,065	.265	.271	.465	.504	.665	.802	.865	1,313
.070	.070	.270	.277	.470	.510	.670	.811	.870	1.333
.075	.075	.275	.282	475	,517	.675	.820	.875	1,354
.080	.080	.280	.288	.480	,523	.680	.829	.880	1.376
.085	.085	.285	.293	.485	.530	.685	.838	.885	1.398
.090	.090	.290	.299	.490	.536	.690	.848	.890	1.422
.095	.095	.295	.304	.495	.543	.695	.858	.895	1.447
.100		.300	.310	.500	.549	.700	.867	.900	1,477
.105		.305	.315	.505	.556	.705	.877	.905	1.499
.110		.310	.321	.510	.563	.710	.887	.910	1.52
.115		.315	.326	.515	.570	.715	.897	.915	1.55
.120	.121	.320	.332	.520	.576	.720	.908	.920	1.58
.12		.325	.337	.525	.583	.725	.918	.925	1.62
.13	.131	.330	.343	.530	.590	.730	.929	.930	1.65
.13		.335	.348	.535	.597	.735	.940	.935	1.69
.14		.340			.604	.740	.950	,940	1.73
		.345	.360	.545	.611	.745	.962	.945	1.78
-15		.350			.618	.750	.973	.950	1.83
.15					.626	.755	.984	.955	1.88
.16					.633		.996	.960	
.10					.640	.765	1,008	.965	
.1	70 .172	.37	.388	.570	.648		1.020	.970	
1.	75 .177 80 .183					.775	1.033	.975	2.18
	85 .18					.780	1.045	.980	
	90 .19							.985	
	95 .19							.990	
_			5 .41	8 .595	.68	.795	1.085	.995	2.99

<sup>6</sup> Values reported in this table were calculated by Thomas O. Maguire and are reproduced with his kind permission.

TABLE J CONFIDENCE INTERVALS AROUND r ON p FOR n = 3,4,...,400
(FIND UPPER-LIMIT VALUE ABOVE THE PRINCIPAL DIAGONAL,
AND LOWER-LIMIT VALUE BELOW IT)



THE CORRELATION COEFFICIENTS CRIT TABLE I

df = n - 2	$\alpha = .10$	.05	.02	.01
1	,988	.997	,9995	.9999
2	,900	.950	.980	.990
3	,805	.878	,934	.959
4	,729	.811	.882	.917
5	,669	.754	,833	.874
6	,622	.707	,789	.834
7	,582	.666	.750	.798
8	,549	.632	.716	.765
ŷ	,521	.602	.685	.735
10	.497	.576	.658	.708
11	,476	.553	.634	.684
12	,458	.532	.612	.661
13	,441	.514	,592	.641
14	,426	.497	.574	.623
15	412	.482	,558	.606
16	,400	.468	.542	.590
17	,389	.456	.528	.575
18	.378	.444	.516	.561
19	.369	.433	.503	.549
20	,360	.423	.492	.537
21	,352	.413	.482	.526
22	.344	.404	.472	.515
23	.337	.396	.462	.505
24	.330	.388	.453	.496
25	.323	.381	.445	.487
26	.317	.374	.437	.479
27	.311	.367	.430	.471
28	.306	.361	.423	.463
29	.301	.355	.416	.456
30	.296	.349	.409	.449
35	.275	.325	.381	.418
40	.257	.304	.358	,393
45	.243	-288	.338	.372
50	.231	.273	.322	.354
60	.211	-250	.295	.32
70	.195	.232	.274	.30
80	.183	.217	-256	.28
90 100	,173 ,164	.205 .195	.242	.26

Table I is reprinted from Table V.A. of Fisher & Yates, Stantical Methods for Research Workers, published by Oliver and Boyd Ltd., Edinburgh, and by permission of the author and publishers. If the absolute raile of an r from a sample of size n exceeds the tabled value of a reason of the first form of the research of the standard of the following the sample of size not accept the tabled value of a significance; the alternative properties of the sample role of 39 with n = 20 leads to rejection of the physicists p = 0 in the 51 level of significance;


TABLE K ABSOLUTE VALUES OF THE CRITICAL VALUES OF SPEARMAN'S RANK CORRELATION COEFFICIENT, I., FOR TESTING THE NULL HYPOTHESIS OF NO CORRELATION WITH A TWO-TAILED TEST.

n	α = .10	$\alpha = .0$	$\alpha = 0$	$\alpha = .01$
5	0.900	T -	_	
6	0.829	0.886	0.943	
7	0.714	0.786	0.893	! —
8	0.643	0.738	0,833	0.881
9	0.600	0.683	0,783	0.833
10	0.564	0.648	0.745	0.818
11	0.523	0.623	0.736	0.794
12	0.497	0.591	0.703	0 780
13	0.475	0 566	0.673	0.745
14	0.457	0.545	0.646	0.716
15	0.441	0.525	0.623	0.689
16	0.425	0.507	0,601	0.666
17	0.412	0.490	0.582	0.645
18	0.399	0.476	0.564	0.625
19	0.388	0.462	0.549	0.608
20	0.377	0.450	0.534	0.591
21	0.368	0,438	0.521	0.576
22	0.359	0.428	0.508	0.562
23	0.351	0.418	0.496	0.549
24	0.343	0.409	0 485	0.537
25	0.336	0.400	0.475	0.526
26	0.329	0.392	0.465	0.515
27	0.323	0.385	0.456	0.505
28	0.317	0,377	0.448	0.496
29	0.311	0.370	0.440	0.487
30	0.305	0.364	0.432	0.478

Adapted from E. G. Olds, "Distributions of sums of squares of rank differences for small numbers of individuals," Annals of Mathematical Statistics, 9 (1938), 133-48, and "The 5% signifi-cance levels for sums of squares of rank differences and a correction." Annals of Mathematical Statistics, 20 (1949), 117-18, by permission of The Institute of Mathematical Statistics.

\* The tabled values are absolute values of the critical values for two-tailed tests. For example, the critical values of  $r_n$  for n = 10

and  $\alpha = .10$  are +0.564 and -0.564.



Reprinted from Table 15, Biometrika Tables for Statisticians (2nd ed., 1962), eds. E. S. Pearson and H. O. Hartley, by permission of the Biometrika Trustees and the publisher University Press.

Į×,	10.2956 10.3441 10.3923 10.4403	10.4881 10.5357 10.6301	10.7703 10.7703 10.8167	10.9087 10.9545 11.0000 11.0454 11.0905	11.1355	11.4018 11.485 11.485	11.5758 11.6190 11.6619 11.7047 11.7898 11.8322
ž	11236 11449 11664	12321 12321 12769	13225 13456 13689 13924	14641 14884 15129	15376 15625 15876 16129	16641 16900 17424 17689	17956 18225 18496 18769 19044 19321 19600
ν	901 108	9 3355	113 113 113 113 113	2 222	75. 25. 25. 25. 28.	137 137 137 137 137 137 137	134 135 137 138 140
[X	8.4261 8.4853 8.5440 8.6023	8.7178 8.7750 8.8318 8.8882	8.9443 9.0000 9.0554 9.1104 9.1652	9.2195 9.3274 9.3808	9,5394 9,5394 9,5917 9,6437	9.6954 9.7468 9.8489 9.8995	10.0499 10.0499 10.0895 10.1489 10.1980
\$	5041 5184 5329 5476	5776 5929 6084 6241	6561 6724 6889 7056	7225 7396 7569 7744	8281 8464 8649	925 925 9409 9604	10000 10201 10404 10609 10816 11025
≷	52525	5 2 2 8 2 5	88321	88888	8 2823	8 9 8 8 8	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
√N N	6,0000 6,0828 6,1644 6,2450 6,3246	6.4031 6.4807 6.5574 6.6332	6.7823 6.9282 7.0000	7.1414 7.2111 7.2801 7.3485	7.4162 7.4833 7.5498 7.6158	7.7460 7.8102 7.8740 7.9373 8 0000	8.0623 8.1240 8.1854 8.2462 8.3066
*	1296 1369 1444 1521 1600	1681 1764 1849 1936	2209 2304 2401 2500	2501 2704 2809 2916	3136 33249 3364	3721 3721 3969 4096	4356 4489 4624 4761 4900
N	36 33 39 40	<del>2</del> 4444	4 <b>4</b> 8468	25225	3,5888	3 26838	98898
IX	1.000 1.4142 1.7321 2.0000 2.2361	2,4495 2,6458 2,8284 3,0000 3,1623	3.3166 3.4641 3.6056 3.7417 3.8730	4.0000 4.1231 4.2426 4.3589 4.4721	4.5826 4.6904 4.7958 4.8990	5.0990 5.1962 5.2915 5.3852 5.4772	5.5678 5.6569 5.7446 5.8310 5.9161
£	14695	36 49 100 100	225 255 255 255 255	289 324 361 400	441 484 529 576 625	727 729 784 841 900	961 1024 1089 1156 1225
≈	WW4W	97.86Q	122212	202222	ដដដដដ	328278	EEEEE

TABLE L PROBABILITIES ASSOCIATED WITH VALUES AS LARGE AS OBSERVED VALUES OF 5 IN KENDALL'S 7 WHEN THE NULL HYPOTHESIS OF NO CORRELATION IS TRUE

_		1'0	lues of N		s		Values of N				
S	4	5	8	9	"	6	7	10			
0	.625	_592	.548	.540	1	,500	.500	.500			
2	.375	403	.452	.460	3	,360	.386	.431			
4	.167	.242	360	.381	3	.235	.281	.364			
6	.042	.117	.274	306	7	.136	.191	.300			
8		.042	.199	.238	9	.068	.119	.242			
10		.0033	.138	.179	11	.028	.068	.190			
12	1	ì	.039	.130	13	.0083	.035	.146			
14		1	.054	.090	15	.0014	.015	.103			
16	1	Į.	.031	.060	17	i	.0054	.078			
18	1	ì	-016	.038	19		.0014	.054			
20	1	)	.0071	.022	21		.00020	.036			
22	1	1	.0028	.012	23	}		.023			
24	1	ì	.00037	.0063	25	]	)	.014			
26	1		.00019	.0029	27		1	.0083			
28	1	1	.000025	.0012	29	Ī	1	.0046			
30	1		1	.00043	31	l	1	.0023			
32	ì	ł	1	.00012	33	l	į	.0011			
34		1	1	.000025	35	ĺ	I	.00047			
36	1	1	}	.0000028	37	1	1	.00018			
	1		ı	1	39	1	.000058				
	1	1	ì	i	41	1	i	.000015			
	1	1	1	1	43	1	1	.0000002			
_	3	1	1	1	45	ì	1	,0000000			

Adapted from M. G. Kendall, Rank Correlation Methods, 3rd ed. (London: Charles Griffin & Company Limited, 1962), Appendix Table 1, by permission of the publisher.

	1																						
\ \ V	19.6469	19.7231	19.7737	19.7990	19.8494	19.8997	19.9249	20,0000	03.00	20.0499	20.0749	20.1246	20.1494	20.1742	20.2237	20.2485	20.2731	20 2978	20.3470	20.3715	20.3961	20.450	20 4695 20.4939
ž.	148996	151321	152881	154449	156025	156816	158404	160000	160801	161604	162409	164025	164836	165649	167281	168100	168921	170569	171396	1/2225	173056	174724	176400
×	386	389	391	252	395	396	398	88	401	405	50	402	406	664	409	2	145	413	444	?	416	8 5	420
_	)				-		-	_					_										=
N/	18.7350 18.7617 18.7883	18.8149	18.8680	18.9209	18.9737	19 0000	19.0526	19.1050	19 1311	19.1572	19.2094	45.C7.CT	19.2614	19.3132	19 3391		19.3907	19.4422	19.4936	10 5101	19.5448	19.5704	19.6214
*	123201 123904 124609	125316	126736	128164	129600	130321	132496	133225	133956	135424	136161		138384	139129	140625	141376	142129	142884	144400	145161	145924	147456	148225
×	351 352 353	355	356	358	360	361	363	365	366	368	369		371	373	375	376	377	378	380	381	387	384	£
√N N	17.7764 17.8045 17.8326	17.8885	17.9165	17.9722	18.0278	18.0555	18.1108	18.1639	18.1934	18.2483	18.2757	10 2303	18.3576	18.3848	18 4391	18.4662	18.4932	18.5472	18.5742	18.6011	18 6279	18.6815	Sec.
۳.	99856 100489 101124	102400	103041	104976	103625	106276	108241	006807	109561	110889	112225	112806	113569	114244	115600	116281	116964	118336	119025	119716	121104	121801	
N	316 317 318	320	321 322	324	į į	327	329	95	332	333	332	336	337	339	340	341	342	7	3	346	78	320	
		=			_	_																	
ν,	16.7631 16.7929 16.8226	16.8819	16.9115	17.0000	17.0587	17.0880	17.1464	17 2047	17.2337	17.2627	17.3205	17,3494	17,3781	17.4356	17.4642	17.4929	17.5499	17.5784		17.6635	17.6918	17.7482	
N3	78961 79524 80089 80656	81228	82369	83521 84100	84681	85264	86436	87616	88209	89401	00006	10906	91204	92416	5000	93634	94864	96100	16230	97344	97969	99225	
>	281 283 283 284	285	286 287	388 380 380	291	292	295	296	297	532	300	25	303	Š	300	302	308	310	311	315	35	315	

(N)	15.6844	15.7480	15.8114	15.8430	15.9060	15.957	16 0000	16 0312	16.0935	16.1245	16,1555	16.2173	16.2481	10:4100	16 3095	16.3707	16.4012	10.431	16,4621	16.5227	16.5529	16,5831	16.6132	16,6733	16,7033	16.7332
ž	60516	2000	625001 62500	63001	63504	64516 65025	65536	66049	67081	67600	68121	69169	96969	702.55	70756	71824	72361	00677	73441	74529	75076	15625	76176	77784	77841	78400
×	246	248	256	251	252	255	,	257	23.53	260	197	262	50	265	566	268 268 268	569	2/0	17	72.5	274	275	276	77.	273	280
500	t					-	=	_			==		=	_	_	_	-	=	_	_	_	_				=
Ş	9969	2095	14 6287	700 *1	14,7309	14.7986	14 0254	14.8661	14.9332	15.0000	150333	15,0665	15,1327	15.1658	15.1987	15.2315	15.2971	15.3297	15,3623	15,3948	15 4596	15,4919	15.5242	15.5563	15.6205	15.6525
ž		11611	45796	57.04	47089	47961	48400	48841	49729	20625	\$1076	\$1529	252	52900	53361	53824	54756	\$5225	53696	56169	5212	57600	58081	58564	20536	60025
>	: \	122	82	â	326	25	220	ដូន	គ្រ	12	354	222	35	ន្ត	231	333	35	235	236	237	35	ia a	241	55	32	£
Į	2	13,2665	13,3417	13.4164	13 4536	13.5277	13.6015	13.6382	13.7113	13.7477		13.8203	13.8924	13.9642	14 0000	14.0357	14.0712	14.1421	14,1774	14.2127	14.2478	143178	14.3527	14.3875	7,722	14.4914
		30976	1987	32400	32768	33489	34225	34596	3534	35721		36481	37249	38025	311.31	38809	39204	10000	10707	40804	4 100	42025	ASASA	42849	43264	120
	×.	178	8,0	8	181	183	183	186	183	681	2	162	8	19.5	301	197	861	8	303	502	203	ğ	ý	35	500	512
		<u> </u>									_	-	_	=	~	=	-	-	-	-	-	-		===		_
(cout )	X.Y	11 8743	26.7	11.00	12.0830	17.1	17.74	12,3882	12,3288	1097	7.44.7	12,4900	12.5698	12 6095	,	12,7279	12.7671	12.8062	11 00 11	12.9228	12.9615	13000		13.1149	13.1529	13.1909
	×	19861	7 ST OF	20736	21316	1190	51 51 51 51 51 51 51 51 51 51 51 51 51 5	10311	10.0	3116	1907	24336	1965	£15.		12.5	26569	36896		27889	28274	28561	200	1850	29929	30276
TABLE M	×	. 3	23	<b>Ξ</b> Ξ	<u> </u>	72	25	3 5	52	34	22	951	25	65	3	<u> </u>	29	<u> </u>	3	25	89	<u>2</u> 5	2 :	ΞĒ	12	55
			543	t																						

(2	N V	25.8763	25.8457	25.8650		25.9037	25.9422	25.9615	43.9808	26.0000	26.0192	26 0576	26.0768	0200.30	26.1151	26.1343	26.1534		26.1916	26 2298	26.2488	26.2679	26,2869	26.3059	26.3249	26.3629		26.3818 26.4008	26.4197	26.4575
7/4	771888	444889	446224	448900	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	451584	452929	454276		456976	458329	461041	462400	463761	465124	466489	469225	10000	471969	473344	474721	000	477481	478864	481636	483025			487204 488601	
>	999	299	888	670	S	672	6	675	. ;	979	628	679	080	189	85	888	685	289	687	889	88		169	7 E	694	695	969	697	866	3
1×	25.1197	25.1396	25.139	25.1992	25.2190	25.2389	25.2587	25.2982	35 2100	25.3377	25.3574	25.3772	600	25.4165	25 4558	25.4755	25.4951	25.5147	25.5343	25.533	25.5930	35,613,5	25.6320	25.6515	25.6710	5069.67	25.7099	25.7488	25.7682	
*	398161	400680	401956	403225	404496	405769	408321	409600	410881	412164	413449	416025		417316	419904	421201	477200	423801	425104	427716	429025	430336	431649	432964	435600	00000	436921	439569	440896	
×	159	229	634	635	989	637	636	640	2	642	£3	3	;	£ 2	648	649	20	651	709	654	655	929	657	658	99	;	26 26 26 26 26 26 26 26 26 26 26 26 26	663	665 665	
Ň	24.4131	24.4540	24.4745	V4X4.42	24.5153	24.5561	24.5764	7962.77	24.6171	24.6374	24.6779	24 6982	24 7184	24.7386	24.7588	24.7992	2010	24,8395	24.8596	24.8797	-1.0770	24.9199	24 9600	24 9800	25.0000	25,000	25.0400	25.0599	25.0998	
<b>₹</b> /	355216	357604	358801	361701	362464	363609	364816	C7Dnoc	367236	369664	370881	3/2100	373321	374544	376996	378225	379456	380689	381924	384400		386884	388129	389376	390625	391876	393129	395641	396900	
*	596 597	288	£ 6	5	38	93	\$ 5	3	95	88	\$	010	119	613	614	615	919	219	819	620	5	729	623	624	670	979	627	628	630	
V.N	23 6854 23.7065	23.7276	23.7697	23.7908	23.8118	23.8328	23.8747	23 006 5	23 9165	23.9374	23.9583		24 0000	24.0416	24 0624	7580.67	24.1039	24.1247	24.1661	24.1868	24,2074	24.2281	24.2487	24.2899		24.3105	24.3516	24.3721		
N <sub>3</sub>	314721	318096	319225	320356	321489	322624	324900	326041	327184	328329	330625		33176	334084	335241	00+000	337561	339880	341056	342225	343396	344569	346921	348100	14030	350464	351649	354025		
>	262 262 262 263	5 % 5	565	999	267	28	270	571	215	573	575	31.3	572	578	28	2	8	283	284	285	286	282	289	290	401	262	203			
																												54	5	

	TABLE	Σ	(tout)	=	3	ķ	2	ž	Í	2	N	ξ.
	×	ž,	>	۶	ė			100176	22.1585	525	276676	22.9347
	;		20 00	739	207936	21.3542	103	19027	22.1811	527	277729	22.9565
	7	78084	20.5426	457	61880	11,300	161	243049	22.2036	225	279841	23,0000
54	12	789.30	20 5670	200	189012	21,4243	107	244036	70111	530	280900	23 0217
4	7	27,776	20 20 20	3	211600	21,4476	465	242022	200		170104	23 0414
	÷	CTOWN	****			0027	496	246016	22.2711	231	281901	190
		A75.181	20 6398	97	212321	2007	207	247009	22.2935	7	10000	33 0868
	1	23.5	20 6640	45	13	17.	498	248004	22.3159	25	20100	21.1084
		183184	20 6882	2:	9000	1107	664	240001	22.3383	200	386728	23,1301
	2	184071	20.71	12	216225	21.5639	200	250000	22,3607	3		
	9	184900	10.73	=				30000	0191 14	536	287296	23.1517
	:	176401	\$072.04	994	217156	21.5870	25	1000	22.4054	537	288369	23,1733
	Ţ:	0,000	28.70	467	218089	21.6102	700	2000	22 4277	238	289444	23.1948
		87.589	20 8087	468	219024	21.6333	25	254016	22 4499	233	29052	23.2
	35	88356	20 8327	9	219961	2000	Ş	255025	22.4722	240	291600	23.23 (7
	2	89228	20 8267	470	22000	27.07.12					107000	1030 00
	,			:		3,000	206	256036	22,4944	Ī.	292681	73.73
	436	960061	20 8806	7	77	7367	203	257049	22.5167	242	1075	77.70
	437	19006	20.0013	777	107771	7386	808	258064	22,5389	543	204849	2000
Ī	438	101844	20.9284	2,5	35355	2177.15	\$00	259081	22.5610	7	295936	22.52.5
•	2	222	20.025	722	225625	21.7945	510	260100	22.5832	950	570767	20.07
•	9	193661	*0.2.0*				;		****	277	108116	23.3666
	;	194491	2,000	476	226576	21.8174	S	261121	77.0023	25	200200	23.3880
•	:	24.50	11 0238	411	227529	21.8403	25	707	170.07	878	300304	23,4094
	:5	06230	21.0476	478	228484	21.8632	25	20102	717966	249	301401	23.4307
7	3	97136	21.0713	479	229441	21.880	218	265225	22.6936	820	302500	23.4521
4	45	198025	21.0950	490	20402	71:300	:					,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
		,,,,,,,		187	131116	21 9317	316	266256	22.7156	331	30360	77.7
٩.	_	00000	200	785	232324	21.9545	517	267289	22,7376	225	100	22.4
7.		2000	9	483	233289	21.9773	218	268324	12.7596	25	202202	5255
7 *		103101	1 KGK	484	234256	22.0000	219	269361	22.7816	100	20000	22 4884
7	25	005	21 2132	485	235225	22.0227	520	270400	22.8035	? =	30905	
,							5	1777.120	130000	356	300136	23.5797
7	12	203401	21.2368	486	236196	22.0434	125	272484	22 8473	557	310249	23.6008
₹.	22	25	21.2603	48/	23/165	22.000	223	273529	22.8692	558	311364	23.6220
٠.	2	50.50	11.020	CSP	23012	27 1133	224	274576	22,8910	559	312481	23.6432
7 4	,,,	207025	21.3307	6	240100	22.1359	525	275625	22.9129	260	313600	23.6643
1	:		=									

18	31.0000 31.0161 31.0322 31.0483	31.0644 31.0805 31.0966 31.127 31.1288	31.1448 31.1609 31.1769	31.2410 31.2570 31.2570	31,32890 31,3289 31,3528	31.3847 31.4006 31.4166 31.4325	31,4484 31,4643 31,4802 31,4960 31,5119	31.5278 31.5436 31.5595 31.5753 31.5911 31.6070	31.6228
•	923521 925444 927369 929296	933156 935089 937024 938961	940900 942841 944784 946729	952576 954529 956484	962361 964324 964324 966289	970225 972196 974169 976144	978121 980100 982081 984064 986049	990025 992016 994009 996004	1000000
×	963 963 963 963 963	98828	272 272 272 273	576 578 878 878	983 58 883 186 883 r>86 86 86 86 86 86 86 86 86 86 86 8	986 987 9887	86 86 86 86 86 86 86 86 86 86 86 86 86 86 86 86 8	999 997 998 998	1000
1×	30.3480 30.3645 30.3809 30.3974	30,4467 30,4467 30,4631 30,4531	30.5123 30.5287 30.5450 30.5614	30,5778 30,5941 30,6268 30,6268	30.6594 30.6757 30.6920 30.7083	30.7409 30.7571 30.7734 30.7896	30.8221 30.8221 30.8545 30.8707 30.8869	30,9031 30,9192 30,9354 30,9516 30,9577	20,7037
*	848241 850084 851929 853776 85574	859329 861184 863041	866761 868624 872336	874225 876096 877969 879844 881721	883600 885481 887364 889136	894916 896809 898704 900601	902500 904401 906304 908209 910116	912025 913936 915849 917764 919681 921600	
×	922 923 924 924	926 927 929 930	931 933 934 934	938 938 938 938	2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	948 948 948 948	953 953 953 953 953	958 958 959 960	
	r								
X	29.6816 29.6985 29.7153 29.7321 29.7489	29.7658 29.7825 29.7993 29.8161 29.8329	29.8496 29.8664 29.8831 29.8998	29.9333 29.9500 29.9666 29.9866	30 0167 30 0333 30 0500 30.0666	30.0998 30.1164 30.1330	30.1828 30.1993 30.2159 30.2324 30.2490	30.2655 30.2985 30.2985 30.3150 30.3150	
*	776161 777924 779689 781456 783225	784996 786769 788544 790321 792100	793881 795664 797449 799236 801025	802816 804609 806404 808201	813604 813604 815409 817216 819025	820836 822649 824464 826281	829921 831744 833569 835396 837225	839056 840889 842724 844561 846400	
×	881 882 883 884 885	888 888 888 890 890	891 892 893 894 895	898 898 898 898	98888 8643 8448	28824 10884 10884	911 913 914 915	916 917 918 920	
-									
$\sqrt{N}$	29.0000 29.0172 29.0345 29.0517 29.0689	29 0861 29.1033 29.1204 29.1376 29.1348	29.1719 29.1890 29.2062 29.2233 29.2233	29.2575 29.2746 29.2916 29.3087 29.3258	29.3428 29.3598 29.3769 29.3939 29.4109	29.4279 29.4449 29.4618 29.4788 29.4988	29.5127 29.5296 29.5466 29.5635 29.5804	29.5973 29.6142 29.6311 29.6479 29.6648	
N.	707281 708964 710649 712336 714025	715716 717409 719104 722500	724201 725904 727609 729316 731025	732736 73449 736164 737881 739600	741321 743044 744769 746496 748225	749956 751689 753424 755161 756900	758641 760384 762129 763876 765625	767376 769129 770884 772641 774400	
>	841 842 844 845	846 847 849 849	851 852 853 854 855	856 858 858 859 860	863 863 864 865	866 867 870 870	871 872 873 874 875	878 878 878 880	
								547	

-	TABLE M (C	(cont.)		ž	(×,	×	ž	N/N	>		:
×	Ş.	X, 1	٤	ς		F	177702	27,7669	908	649636	28,3901
ē	101161	26 4764	736	271696	27.133	12	\$95984	27,7849	200	652864	28 4233
į,	108.64	26 495	282	7977	27.1662	25	20,02	27,8209	803	654481	28.44.05
ē	919564	26 5330	23	546121	17,1810	ži.	600625	27,8388	⊋ ==	930	
ê	\$101A	26.5518	<b>}</b>			ř	451503	27.8568	811	657721	28.4784
2	Attact	16.5707	7	\$4908	27.2213	17.0	603729	27.8747	200	090099	28.5132
į,	578657	26 5895	:53	33030	17.2580	778	605284	27.8927	2	662596	28,5307
o.	50136	26 6083	£	553536	27.2764	779	606841	27,9106	813	664223	28.5482
ē	90.00	26 6458	7	\$55025	22.2947	08/	00000		,	770377	18 6657
?	5		,		01117	781	196609	27.9464	22	662330	28.5832
ž	10.5	36 6646	272	\$58000	27,3313	787	611524	25025	00	669124	28.6007
2	0,000	101.45	748	\$59504	27,3496	783	61,5039	36,000	618	670764	28.6182
:	9611.00	26.7.08	£.	26100	77.3679	785	616225	28 0179	820	672400	28 6520
2	211235	26.7395	2	20700					ŝ	673031	28.6531
	,		5	(0079)	27.4044	286	617796	28.035/	35	474K84	28.6705
2	200	20.735	.02	\$65504	27.4226	787	616369	28 0355	15	677329	28,6880
÷	14034	76 7935	753	\$67009	27,4408	788	7600	1000 00	824	678976	28,7054
25	96915	26 8142	757	568516	27.4591	702	001525	28,1069	825	680625	28.7228
2	518400	26.8328	35	3/00/2	211411	2			1		000
		* 1 20 22	756	471436	77.4955	162	625681	28,1247	820	02770	20.7402
7	5 984	26 8314	25	671010	27.51.16	792	627264	28.1425	827	693929	20.00
1	2	2000	24	272564	27.5318	793	628849	28,1603	200	150000	28.702
32	57176	26.9072	259	\$76081	27.5500	794	630436	28.1/80	830	688900	28.8097
Į.	525625	26,9258	92	\$17600	27.3681	S	034043		=		
		-	5	10101	CY35 LX	706	613616	28.2135	831	690561	28.827
91	527076	26.74	5	250033	17 6043	797	635200	28,2312	22	69224	7778,61
2	2.60	10.00	12	691.88	17.6225	798	636804	28.2489	23	66,260	10.00
37	21.513	22 0000	125	583696	27,6405	799	638401	28,2666	634	2000	2000
12	312900	27,0185	765	585225	27.6586	88	640000	28.2843	6	07170	0.010
		Q_50 55	386	495785	17 6767	108	641601	28,3019	836	968869	28,9137
5;	111014	22.00	191	588289	27.6948	802	643204	28.3196	837	700509	
:=	\$17289	27.0740	268	\$59824	27.7128	803	644809	28.3373	200	10000	
3	538756	27,0924	692	391361	27.7308	200	44	25,204	200	105600	
ĭ			-								





TABLE N (cont.)



3




TABLE O 2-SCORE EQUIVALENTS OF SELECTED PERCENTILES IN THE UNIT NORMAL DISTRIBUTION\*

Percentile	z	Percentile	z į	Percentile	z
0.01	-3,719	28	-0.583	74	0 643
0.05	-3.291	30	-0.524	75	0.675
0.1	-3.090	32	-0.468	76	0.706
0.5	-2.576	34	0,412	78	0.772
1	-2.326	36	-0.358	80	0.842
2	-2.054	38	-0.305	82	0.915
3	1.881	40	-0.253	84	0.994
4	~1.751	42	-0.202	86	1.080
5	-1.645	44	-0.151	88	1.175
6	-1,555	46	0.100	90	1,282
7	-1,476	48	0 050	91	1,341
8	-1,405	50	0,000	92	1,405
9	1,341	52	0.050	93	1.476
10	-1,282	54	0.100	94	1.555
12	-1,175	56	0.151	95	1.645
14	-1,080	58	0.202	96	1.751
16	-0.994	60	0.253	97	1.881
18	-0.915	62	0.305	98	2.054
20	-0.842	64	0.358	99	2.326
22	-0.772	66	0.412	99,5	2.576
24	-0.706	68	0.468	99,9	3.090
25	-0.675	70	0.524	99.95	3.29[
26	-0 643	72	0,583	99.99	3.719

<sup>\*</sup> Values in this table were found by interpolating in Table B.

# TABLE P CONFIDENCE INTERVALS AROUND A SAMPLE PROPORTION, p, ON A POPULATION PROPORTION, P

From E. S. Pearson and C. J. Clopper, "The use of confidence intervals or fiducial limits illustrated in the case of the binomial," *Biometrika*, 26 (1934), 404. Reproduced by permission of the *Biometrika* Trustees.





TABLE P (cont.)



# APPENDIX B

# ANSWERS TO PROBLEMS AND EXERCISES

# CHAPTER 2

```
2. a. nominal b. ordinal 3. a. 6 yr 4.5 mos. -6 yr 5.5 mos. b. 2 ib 12.75 oz-2 ib 13.25 oz c. $342.50-$343.50  
4. a. X_{12} b. X_{11} c. X_{22} 5. a. 19 b. 10 c. 11 d. 13  
6. a. 60 b. 30.5 c. 5 d. 15² = 225  
7. a. 26 b. 15 c. 7 d. 9 + 4 + 1 + 25 = 39  
8. a. X_1^2 + X_2^2 + X_3^2 + X_4^2 b. X_4 + X_5 + X_6 c. (X_1 + X_2 + X_3)^2 9. a. 3\frac{5}{c^2}X_1 b. \left(\frac{15}{c^2}X_1^2\right)^2 c. \frac{5}{c^4}(X_1 + T) d. \frac{5}{c^2}X_1(X_1 + 1) = \frac{5}{c^2}(X_1^2 + X_2)
```

I. a. nominal b. ordinal c. ratio d. nominal

#### CHAPTER 3

2. 
$$P_{50} \approx 108.41$$

# CHAPTER 4

- t. Mean = 2.56; median = 2.8; mode = 3.05.
- 2. Mean = 3.06; median = 3.3.
  - 3. Mean = 14.34; median = 14.45.
- 4. Mean = 3(14.34) = 43.02; median = 43.35.
- 5. Mean = 8.86; median = 9.65.
- Mean = 13.3; combined median can't be determined from a knowledge of individual medians and n's only.

# 7. At point D.

CHAPTER 5

# Inclusive range = 21; variance, s<sup>2</sup> = 32.34; standard deviation, s = 5.68; mean deviation = 4.22.

2. 
$$Q = \frac{126.17 - 96.17}{2} = 15$$
.

3. 
$$s_x^2 = 0.039$$
.  $s_x = 0.198$ .

- 4. a. positively skewed b. positively skewed
- c. positively skewed d. negatively skewed
- 5, "greater than"; variance of combined groups is 66.9.
- 6. b. 0.73 c. 2.60 d. ~2.60
- 7. a. Student B. b. Test 1:  $z_A = -.64$ ,  $z_B = +.41$ . Test 2:  $z_A = 2.49$ ,  $z_B = -.07$ .
  - c. Student A.

8. 
$$\sum_{i=1}^{n} z_{i}^{2} = \sum_{i=1}^{n} \frac{(X_{i} - \bar{X})^{2}}{x_{i}^{2}} = \frac{1}{x_{i}^{2}} \sum_{i=1}^{n} (X_{i} - \bar{X})^{2} = \frac{1}{x^{2}} (n-1)s_{x}^{2} = n-1.$$

# CHAPTER 6

5. Variance of Y for X = 1.50 is also 0.50.

#### CHAPTER 7

1. If  $z_x = z_y$ , then  $r_{xy} = (\sum z_x z_y)/(n-1) = (\sum z_x^2)/(n-1)$ . In Prob. 8 in Chapter 5, it was shown that  $\sum z_x^2 = n - 1$ . Hence

$$r_{xy} = \frac{\sum z_x^2}{n-1} = \frac{n-1}{n-1} = 1$$
, when  $z_x = z_y$ .

- 2. Brown and Smith will obtain the same value for the correlation coefficient, since height in inches is 12 times height in feet. A linear transformation of X and/or Y will not change the value of the correlation coefficient.
- 3. a. positive b. negative c. positive d. positive e. negative
- 4. Since  $r_{xy}$  can't exceed +1,  $r_{xy} = (s_{xy})/(s_x s_y) < 1$ . We know that  $s_x = 5$  and  $s_y = 4$ ; hence  $(s_{xy})/(5 \cdot 4) < 1$ , which implies that  $s_{xy} < 20$ .
- 5. X is more closely linearly related to Z than to Y.
- 6. The researcher is inferring a causal relationship solely from correlational evidence. He has no justification for doing so. It may well be the case—and probably is so—that teachers' salaries and the "drop-out rate" are both a function of the social and economic status of the community and that increasing teachers' salaries in a given school would not bring about a decrease in the "drop-out rate."
- 7. a. r = -.04.
  - c. X and Y are curvilinearly related. There is almost no linear relationship between them as indicated by the Pearson product-moment correlation coefficient of -04.
- 8. a.  $r_{xy} = .92$ . b.  $r_{xy} = .33$ .

## CHAPTER 8

1. c. Y = 2 + X



- 2. As the line moves from X = 2 to X = 3, it drops one unit, from Y = 2 to Y = 1. Hence the slope, b<sub>1</sub>, of the line is -1. If the line falls one unit on the Y axis for each unit it moves to the right on the X axis, then the line must rise one unit for each unit it moves to the left on the X axis. Thus, as X moves from 2 to 0, the line rises 2 units on the Y axis from 2 to 4. Hence, b<sub>0</sub> = 4.
- 3. a. 2.93 b. 1.80
- b. 1.80 c. 2.06
- d. 3.24

4.  $b_0 = 46.76$ ,  $b_1 = .194$ . Thus  $\hat{Y} = .194X + 46.76$ .  $\hat{Y}$  for an X of 36 is 53.16.

5. 
$$s_{\bullet} = 7.16$$
.  
7.  $Y_1 = 99.37$ ,  $Y_{\bullet} = 93.63$ .

- 8. No.
- 9. Smith's regression line will have the equation  $\hat{Y} = 5X + 250$ .
- 10. Transforming X into cX + d divides  $b_1$  by c, i.e., for the transformed X's, the slope of the regression line is  $b_1/c$ . Since  $b_0 = Y - b_1X$ , the Y intercept for the transformed data, denoted by ba, is

$$b_0^* = \overline{Y} - \frac{b_1}{c}(c\overline{X} + d) = \overline{Y} - b_1\overline{X} - b_1\frac{d}{c} = b_0 - b_1\frac{d}{c}$$

#### CHAPTER 9

- a. point biserial b. Spearman's rho c. rank biserial
- d. Pearson's r e. Phi coefficient f. rank biserial
- 2. a.  $\phi = 0.08$ . b.  $\phi = 0.35$ .
- 3.  $r_{pb} = 0.25$ .
- 4.  $r_{bis} = 0.47$ .
- 5.  $r_{ki} = -0.75$
- 6.  $\tau = 0.60$ .
- 7.  $r_{ut} = 0.50$ .
- 8. a.  $r_s = 0.87$ . b.  $\tau = 0.76$ .
- 9.  $r_{rb} = 0.36$ .
- 10. a. Partial r equals 0.32. b.  $R_{xy,\epsilon} \approx 0.38$ .
- 11.  $b_0 = 0$ ,  $b_1 = .32$ ,  $b_2 = .12$ .  $R_{y_{1,2}} = 0.41$ .
- 12. No.

# CHAPTER ID

- 1. a. P(A) = 13/52 = 1/4. b. P(A) = 1/52. c.  $P(A \cap B) = 0/52 = 0$ . d.  $P(A \cup B) = P(A) + P(B) - P(A \cap B) = (13/52) + (4/52) - (1/52) =$
- A: a child has perceptual problems.
  - B: a child has emotional problems.
- $A \cap B$ : a child has both perceptual and emotional problems.
- $P(B) = .06, P(A \cap B) = .015.$  $P(A \cup B) = .03 + .06 \sim .015 = .075.$
- b. 210
- d. If n = 19, then (n + 1)! = 20! = 20(19!).
- 4. No.

6. 
$$\binom{13}{5} = 1287$$
.

7. 
$$1 + 4 + 6 + 4 + 1 = 16$$
:  $2^4 = 16$ .

8. 
$$\frac{1}{2}\binom{10}{5} = \frac{1}{2}(252) = 126.$$

9. 
$$P(X=0) = {10 \choose 0} (1/2)^0 (1/2)^{10} = (1/2)^{10}$$

$$P(X=1) = {10 \choose 1} (1/2)^{1} (1/2)^{2} = 10(1/2)^{10}.$$

$$P(X = 2) = {10 \choose 2} (1/2)^2 (1/2)^8 = 45(1/2)^{10}.$$

$$P(X = 3) = {10 \choose 3} (1/2)^3 (1/2)^7 = 120(1/2)^{10}.$$

$$P(X = 0 \text{ or } 1 \text{ or } 2 \text{ or } 3) = (1 + 10 + 45 + 120)(1/2)^{10} = 176(1/2)^{10}.$$

$$P(X = 4 \text{ or more}) = 1 - 176(1/2)^{10} = 1 - 176/1024 = 1 - .17 = .83.$$

12. 
$$(1/2)^5 = 1/32$$
.

13. a. .79 b. .006 c. 
$$(.6915)^3 = .331$$
.

14. 
$$E(X) = 2\frac{1}{5}$$
.

15. a. 
$$E(X) = 2$$
 b.  $\sigma_x^2 = 1$ 

# CHAPTER II

2. 
$$_{.95}F_{8.\infty} = 1.94$$
;  $\frac{.95\chi_8^2}{9} = \frac{15.507}{9} = 1.94$ .

4. 
$$_{.05}F_{4.8} \approx 1/_{.95}F_{8.4} = 1/6.04 = 0.166.$$

5. Positively skewed.

- 6. Median is below n.
- 7.  $_{85}\chi_{20}^2 = 31.41$ . Setting (X 20)/6.32 equal to 1.64, the 95th percentile of the unit normal distribution, gives a value of X equal to 30.36.

8. 
$$\sigma_t^2 = E(t_n^2) = E(F_{1,n}) = n/(n-2)$$
.

#### CHAPTER 12

- 2. a. same b. different c. same d. same e. different
- The probability that s<sup>2</sup> exceeds σ<sup>2</sup> is less than .50.

4. a. 
$$\frac{\sigma_{I}^{*}}{25.0} \frac{\sigma_{I}}{5.00}$$

- 5. A sample of size 40 would have to be taken.
- 6. a. .68 b. .90 c. .99 d. \_50 e. .997
- 7. The 95% confidence interval is 105.97 to 107.53.

9. The statistic in b has the smaller variance error; however, as an estimator of the population mean, it is biased.

# CHAPTER 13

- 1. a. The null hypothesis  $H_0$  is a hypothesis about a parameter or parameters of a distribution that may or may not be "nullified" (rejected) on the basis of evidence provided by a sample from the distribution.
  - b. The alternative hypothesis  $H_1$  is a hypothesis about a parameter of a distribution that specifies values of the parameter other than that specified in  $H_{\Phi}$ . If  $H_0$  is rejected  $H_1$  is accepted, and vice versa.
  - A type I error is the rejection of H<sub>a</sub> when it is true.
  - A type II error is the acceptance of H<sub>0</sub> when it is false.
  - e. The level of significance, a, is the probability of committing a type I error.
  - f. The power of a test,  $1 \beta$ , is the probability of rejecting  $H_0$  when it is false. g. The critical region is all of those values of a sample statistic for which the investigator will reject  $H_{\bullet}$  if his sample yields one such value.

- 2. In none of the cases can it be concluded with certainty that  $H_0$ :  $\rho = 0$  is false.
- 3. a. type I error b. no error c. no error d. type II error
- A type II error cannot be committed if H<sub>0</sub> was rejected since the definition of a type II error is that it is the acceptance of a false H<sub>0</sub>.
- 5. Probability of a type I error is approximately .16.
- 6. a. Critical region is split between both tails.
- b. upper tail c. lower tail
- 7. a. The probability of a type I error is the same for both researchers;  $\alpha=.05$ . b. Rowe has a larger probability of committing a type II error when  $\rho=.10$ . c. null
- 8. a. .10 b. .95 c. .95 d. above .99
- 9. Probability of a type II error when ρ = .10 is greater than .99; hence, the power of the test of H<sub>0</sub>: ρ = 0 against the alternative H<sub>2</sub>: ρ = .10 is less than .01. If there is believed to be a high probability that ρ surely deviates no more from 0 than +.10 or -.10 if at all, then the investigator is wasting his time; he is almost certain to end up accepting H<sub>0</sub>. He should either increase a, which would have the effect of moving the critical values of r closer to zero, or he should increase π.

#### CHAPTER 14

- 1.  $t = \frac{-3.54}{10.47} = -.34$ , which is not significant at the .01 level since  $.995t_{40} = 2.704$ .
- 2. t = (7.90/3.10) = 2.55. Since  $_{.975}t_{18} = 2.101$ , the value of t is significant at the 05 level.
- 3.  $Z_r \pm 2.58/\sqrt{n-3} = (-.221, .605)$ , which equals (-.217, .540) when converted back to the scale of  $r_{ev}$ .
- 4. a.  $t = r\sqrt{(n-2)/(1-r^2)} = 1.033$ . Since  $_{978}t_{73}$  is approximately equal to 2.00, the value of r of .12 is nonsignificantly different from zero at the .05 level. (Corroborate this result by use of Table I in Appendix A.)
  - b.  $z = \frac{.12 (-.01)}{\sqrt{1/72 1.161}} = 0.81$ . Since .975 z = 1.96, the difference between the two
  - V1/72 + 1/81r's is not significant at the. 05 level. (Note that the numerator of z is in terms of Z,'s and not r's. It just happens that in this instance the two Z,'s
- 5. z = (3.178/1.128) = 2.82. Since 2.82 exceeds <sub>x75</sub>z = 1.96, r<sub>xy</sub> and r<sub>xz</sub> are significantly different. The results indicate that the Miller Analogies Test (X) is more highly related to a vocabulary test (Y) than to a nonverbal reasoning test (Z).
- 6. z = -4.72, which is significant far beyond the .01 level.

and r's are identical to two decimal places.)

7.  $z = \sqrt{n} \phi = \sqrt{69}$  (.24) =  $\sqrt{3.97} = 1.99$ . Since  $_{.95}z = 1.64$  and  $_{.05}z = -1.64$ , reject the null hypothesis at the .10 level.

8. 
$$z = \frac{.28 - .27}{\sqrt{(.28)(.72)[(1/873) + (1/837)]}} = 0.47$$
, which is nonsignificant at the .01

level.

- 9. The obtained value of x² is 24.69. The 95th percentile in the x² distribution with 6 degrees of freedom is 12.59. Hence, reject the null hypothesis of "no receivition" between the two factors of classification at the .05 level.
- 10. The obtained value of x² is approximately 2.50. The 90th and 95th percentiles in the x² distribution with 1 degree of freedom are 2.71 and 3.84 respectively; hence the obtained x² is nonsignificant at both the .10 and .05 levels.
- 11. The obtained value of x² is 13.27. The 99th percentile in the x² distribution with 4 degrees of freedom is 13.28. While the obtained value of x² is strictly non-significant at the .01 level, one should have few compunctions about announcing significance at the .02 level.
  - 12. Since zero lies between the limits of the confidence interval, it is true that

$$(\bar{X}_{.1} - \bar{X}_{.2}) - {}_{.875}t_{18}^{5}\bar{\chi}_{.1} - \bar{\chi}_{.3} < 0 < (\bar{X}_{.1} - \bar{X}_{.2}) + {}_{.875}t_{18}^{5}\bar{\chi}_{.1} - \bar{\chi}_{.3}^{2}$$

By subtracting  $(X_1 - X_2)$  from all three sides of the above inequality and then dividing all sides of the inequality by  $-s_{\mathcal{I},1-\mathcal{I},1}$  (which reverses the direction of the inequality), one obtains

$$+_{.975}t_{18} > \frac{(\overline{X}_{.1} - \overline{X}_{.2})}{s_{\overline{X}_{.1} - \overline{X}_{.2}}} > -_{.975}t_{18},$$

which shows that the t-statistic for testing  $H_0$ :  $\mu_1 = \mu_2$  lies between the critical values for testing the null hypothesis at the .05 level.

#### CHAPTER IS

1. a. 
$$df_b = 1$$
,  $df_w = 6$ .  
b.  $df_b = 4$ ,  $df_w = 5$ .  
c.  $df_b = 2$ ,  $df_w = 10$ .  
d.  $df_b = 2$ ,  $df_w = 7$ .

2. a. 
$$_{.95}F_{1.10} = 10.04$$
. b.  $_{.90}F_{4.30} = 2.14$ . c.  $_{.95}F_{2.15} = 3.68$ .

- 3. a. Yes, this statement is equivalent to the statement of the alternative hypothesis,  $H_1$ .
  - b. No. c. Yes. d. No.
- 4.  $MS_B = 20.31$ ;  $MS_w = 6.61$ ; F = 3.07.
- 5.  $MS_B = 190.68$ ;  $MS_w = 221.56$ ; F = 0.86. Since  $_{.99}F_{2.21} = 5.49$ , F is not significant at the .01 level.
- 6.  $MS_B = 312.05$ ;  $MS_w = 48.20$ ; F = 6.5;  $(2.55)^2 = 6.50$ .
- 7. ANOVA table:

Since  $_{48}F_{8.44} = 2.74$ ,  $H_{6}$  is rejected at the .05 level.

8. ANOVA table:

Source of var.	đf	MS	F
Between groups Within groups	2 21	233.79 28.02	8.34

Since  $.95F_{2.21} = 3.47$ ,  $H_0$  is rejected at the .05 level,

9. ANOVA table:

Source of var.	df	MS	F
Between groups	2	9.733	4.38
Within groups	2.7	2.222	

An F-ratio of 4.38 exceeds the 95th percentile in  $F_{2,27}$ , but fails to exceed the 99th percentile in that distribution.

10. ANOVA table:

Reject  $H_0$  at the .01 level.

- 11. The value of  $\phi$  is  $\sqrt{2} = 1.414$ ;  $n_1 = J 1 = 4$ ;  $n_2 = N J = 20$ ; and  $\alpha = .05$ . From Table N in Appendix A (the power of the F-test) we see that the power,  $1 \beta$ , is approximately, 50.
- 12. Since  $s_{\Lambda}^2$  estimates  $\sigma^2/n$ ,  $ns_{\Lambda}^2$  estimates  $\sigma^2$ . Hence, the estimate of  $\sigma^2$  is 20(2.40) = 48.0.

13. 
$$E(MS_0) = \sigma^2 + \frac{n \sum_{i=1}^{J} (\mu_j - \bar{\mu}_i)^2}{J - 1} = 20 + 250 = 270.$$

## CHAPTER 16

1. a. 
$$.95q_{4.20} \approx 3.845$$
. b.  $.95q_{6.120} \approx 4.872$ . c.  $.96q_{10,10} \approx 4.913$ . d.  $.99q_{12,6} = 9.485$ . e.  $.95q_{2,60} \approx 2.829$ .

2, a. Probability equals .05; note the value of .9595,30-

b. One percent would exceed 5.048; note the value of ,9995,30

 Any pair of means must differ by at least <sub>56</sub>q<sub>e,50</sub>√MS<sub>w</sub>/n = 8.32 to be judged significantly different at the .05 level. By this criterion, the following differences between means are statistically significant:

$$X_3 - X_1 = 9.32$$
  $X_5 - X_2 = 16.94$   
 $X_4 - X_1 = 12.17$   $X_6 - X_2 = 17.45$   
 $X_5 - X_1 = 20.41$   $X_5 - X_3 = 11.09$   
 $X_6 - X_4 = 20.92$   $X_6 - X_3 = 11.60$ 

$$X_{.4} - X_{.2} = 8.70$$
  $X_{.6} - X_{.4} \approx 8.75$ 

4.  $MS_e = 5.49$ , n = 18, J = 4,  $_{36}q_{1.68} \pm 3.72$ . The value of  $_{36}q_{1.68} \sqrt{MS_e/n} = 2.05$ . If any two means differ by more than 2.05 units, then they will be judged significantly different at the .05 level by the T-method. By this criterion, the following pairs of means are significantly different:

$$\overline{X}_{1} - \overline{X}_{4} = 3.33$$
,  $\overline{X}_{2} - \overline{X}_{4} = 2.21$ ,  $\overline{X}_{3} - \overline{X}_{4} = 2.38$ .

We see that the "control group," group 4, differs from the three experimental groups that cannot be judged to differ among themselves.

5.  $MS_w = 102.05$ , n = 20, J = 3,  $_{SP}q_{3.57} \pm 4.282$ . The value of  $_{SP}q_{3.57}\sqrt{MS_w/n}$  is 9.68. Any pairs of means differing by at least 9.68 units will be judged significantly different at the .01 level. The following pairs of means are significantly different:

$$\bar{X}_1 - \bar{X}_2 = -13.70$$
  $\bar{X}_1 - \bar{X}_2 = -15.75$ .

6. a. ANOVA table:

Source of var.	df	MS	F
Between groups	3	276.731	15.94
Within groups	24	17.365	

Since the obtained F-ratio exceeds  $_{.85}F_{3.24} = 3.01$ , the null hypothesis of no differences among the four population means is rejected at the .05 level,

b. MS<sub>e</sub> = 17.37, J = 4, n = 7. <sub>M</sub>F<sub>3 21</sub> = 3.01, <sub>M</sub>g<sub>1.24</sub> = 3.901. For the two means to differ significantly by the T-method, the absolute value of the difference between them must exceed <sub>M</sub>g<sub>1.21</sub>√MS<sub>w</sub>/n = 6.14.

For two means to differ significantly by the S-method, the absolute value of the difference between them must exceed  $\sqrt{(J-1)_{Ab}F_{a,2}t_a}$ ,  $\sqrt{MS_a(2/n)} = 6.69$ . The results of the multiple comparisons appear below:

Comparison	Is difference significant by T-method?	Is difference significant by S-method?
$X_{.1} - X_{.2} = -2.57$	No	No
$X_1 - X_2 = -9.16$	Yes	Yes
$X_{.1} - X_{.4} = -13.84$	Yes	Yes
$X_{.2} - X_{.3} = -6.59$	Yes	No
$X_{.2} - X_{.4} = -11.27$	Yes	Yes
$X_{.3} - X_{.4} = -468$	No	No

c. In the contrast  $(\mu_2 + \mu_3)/2 - (\mu_1 + \mu_4)/2$ , the coefficients are

$$c_1 = -\frac{1}{2}, \quad c_2 = \frac{1}{2}, \quad c_3 = \frac{1}{2}, \quad c_4 = -\frac{1}{2}.$$

Thus the value of a; is

$$\delta_{\varphi} = \sqrt{MS_{e}(1+1+1+1)/7} = 1.575.$$

The value of é is

$$\hat{\mathbf{y}} = (21.25 + 27.84)/2 - (18.68 - 32.52)/2 = -1.05.$$

The 95% confidence interval is

$$\psi \pm \sqrt{(J-1)_{.95}F_{3.24}}\,\hat{\sigma}_{\hat{\psi}} = -1.05 \pm 4.73 = (-5.78, 3.68).$$

7. a. ANOVA table:

Source of var.	df	MS	F
Between groups	4	695.06	10.57
Within groups	95	65.75	

The obtained value of F is significant at the .05 level.

b.  $MS_w = 65.75$ , J = 5, n = 20,  $_{59}g_{5.35} \approx 3.94$ . Any difference between two means which is larger in absolute value than  $_{58}g_{5.85} \sqrt{MS_w/n} = 7.13$  is significant at the .05 level by the *T*-method. The significant differences are identified by an asterisk in the following table of differences:

Differences between means

	Acc	SY	so	AY
SY	-1,25			
SO	-8.40*	-7.15*		
AY	8.00*	9.25*	16.40*	
AO	1.70	2.95	10.10*	-6.30

\* Difference is significant at the .05 level.

c. The 95% confidence interval on  $\mu_1 - (\mu_2 + ... + \mu_5)/4$  is (-6.35, 6.37).

## CHAPTER 17

#### 1. ANOVA table:

Source of var.	df	SS	MS	F
Factor A	4	64.26	16.07	1.70
Factor B	5	46.85	9.37	0.99
$A \times B$	20	1164.05	58.20	6.15
Within	120	1136.53	9.47	
Total	149	2411.69		

	Critical value	Obtained value	Decision
Factor A	$_{.99}F_{4.120} = 3.48$	1.70	Do not reject H <sub>b</sub>
Factor B	$_{.99}F_{5.120} = 3.17$	0.99	Do not reject Ha
$A \times B$	$_{.99}F_{20.120} = 2.03$	6.15	Reject H <sub>0</sub>

2. a. Yes b. Yes c. No

Females studying French by the aural-oral method should score 8 points above the general mean on the language mastery test:

$$\mu_{11} = \mu + \alpha_1 + \beta_1 + \alpha \beta_{11} = \mu + 6 + 2 + 0 = \mu + 8.$$

- Case a shows a disordinal interaction. Case b shows an ordinal interaction. In case c there is no interaction since the two lines of the graph are parallel.
- When "treatments" is placed on the abscissa, the ordinal interaction graphed in case b of 5 becomes disordinal.

#### 7. ANOVA table:

Source of var.	df	MS	F
Type of question	1	18.38	0.91
Position of question	1	532.04	26.43
Interaction	1	77.04	3.83
Within cells	20	20.13	

The critical values against which each of the three F-ratios is compared are all the sense: 16F1.50 = 2.97. Hence, we see that the F-tests for "Position of Question" and "Interaction" result in rejection of the null hypothesis.

df

MS

#### 8. ANOVA table:

Š.	ntelligence		1	92.04	3.56
N	lethod of in-	truction	2	233.79	9.03
I	nteraction		2	15.29	0.59
١	Vithin cells		18	25.88	
Hypoth	esis tested	Critical t	alue	Obtained F	Decision
H <sub>4</sub> : ∑ α		.95F1.19 =	4.41	3.56	Do not reject H <sub>p</sub>
$H_0: \sum E$		.s4F2.15 =	3.55	9.03	Reject IIa
$H_{\bullet}: \sum_{i} \sum_{j} \sum_{i} f_{ij}$	$\sum \alpha \beta_{ij}^2 = 0$	.95F2,19 =	3.55	0.59	Do not reject Ho

#### 9. ANOVA table:

Source of var.	dſ	MS	F
Reading achievement	2	28,103,97	158.86*
Reading difficulty	1	730.81	4.13*
Interaction	2	48.02	0.27
Within cells	234	176.91	

<sup>\*</sup> Significant at the .05 level.

Source of var.

t----

# 10. ANOVA table:

Source of vae.	dſ	MS	F
Sex		1.020	
Reading readiness activity	•	1.028	6.31*
Interaction	2	0.033	0.45
Within cells	2	0.042	0.26
ייוניווח כפווג	24	0.163	

<sup>.</sup> Significant at the .05 level.

- a. proportional
   b. disproportional
   c. proportional
   d. disproportional
- 12. If two observations are randomly discarded from the cell at the intersection of the third row and third column, the cell frequencies would be proportional.
- 13. The layout of cell means is as follows:

	Factor B	
	9.675	10.050
Factor A	10.242	10.512
	9.011	9.883

From these data, the following sums of squares are found:

$$SS'_{A} = 0.868$$
,  $SS'_{B} = 0.384$ ,  $SS'_{AB} = 0.104$ .

From the original layout of 40 observations,  $MS_w$  is found to be equal to 0.7830.

The value of c in Eq. (17.15) is 0.1604. Hence,  $MS'_{\omega} = 0.126$ . The remainder of the analysis is reported in the ANOVA table below:

Source of var.	đf	MS'	F
Socio-economic status (A)	2	0.434	3.44
Type of school (B)	1	0.384	3.05
$A \times B$	2	0.052	0.41
Within cells	34	0.126	

The critical value of F for testing the null hypothesis for factor B at the .05 level is about  $_{85}F_{1.34} = 4.12$ . Hence, the hypothesis of no difference between types of school cannot be rejected.

For testing factor A and the interaction of factors A and B at the .05 level, a critical value of  ${}_{36}F_{2}$   ${}_{34} = 3.27$  is used. The null hypothesis for factor A can be rejected, but no evidence exists for rejecting the hypothesis of no interaction.

### CHAPTER 18

- 1. a.  $\hat{\sigma}_a^2 = .042$ .
  - b. Based upon our estimates, the variance of ratings for children being rated by the same judge is about 250 times as large as the variance of judges.
  - c. The 95% confidence interval on  $\sigma_s^2/\sigma^2$  is (-.03, .13). Since it is impossible for  $\sigma_s^2/\sigma^2$  to be negative we could set the interval from 0 to .13.
- 2. a.  $MS_A = 237.25$ ;  $MS_w \approx 43.92$ .

 $\hat{\sigma}_a^2 = (MS_A - MS_w)/12 = 16.11.$ 

 $\hat{\sigma}^2 = 43.92$ .

b. The 95% confidence interval on  $\sigma_a^2/\sigma^2$  is (.16, 2.66).

3.  $\phi_a^2 = (MS_A - MS_{AB})/(nJ)$ 

## 4. ANOVA table:

Source of var.	df	MS	F
VIQ vs. PIQ	1	10.42	0.10
Persons	29	183.84	
Interaction	29	108.52	

The critical value for the F-test is  $.95F_{1,29} = 4.18$ . Obviously the obtained Fratio is far below this critical value. Hence, the hypothesis of no difference between Verbal and Performance IQ can not be rejected.

# 5. ANOVA table:

Source of var.	đf	MS	F
Between subtests	3	23.41	3.22
Persons	11	19.07	
Interaction	33	7.26	

Set  $\alpha = .01$ . Apparent test: The critical value for the F-ratio with the "apparent test" is 22F2.33 = 4.40. Since the F-ratio of 3.22 does not exceed the critical value, the null hypothesis of no differences among population means on the four types of test is not rejected at the .01 level.

### 6. a. $20 \times 32 \times 8 = 5120$ .

b. "Raters" and "ratees" are essentially random-effects factors, because 20/10,000 and 32/20,000 are negligibly different from 0. "Traits" is a fixedeffects factor, because 8/8 = 1. This is a fully crossed design; there is no

- c. There are seven sources of variation in these data: three main effects, three two-factor interactions, and one three-factor interaction (untestable residual, here). The sources of variation are as follows: between raters, between ratees, between traits, raters × ratees, raters × traits, ratees × traits, and raters × ratees × traits
- d. Let raters be factor 1, ratees factor 2, and traits factor 3. Then

$$\begin{split} E(MS_1) &= \sigma^2 &+ 8\sigma_{12}^2 &+ 256\sigma_1^2 \\ E(MS_2) &= \sigma^2 &+ 8\sigma_{12}^2 &+ 166\sigma_2^2 \\ E(MS_2) &= \sigma^2 + \sigma_{123}^2 &+ 32\sigma_{12}^2 + 20\sigma_{22}^2 + 640\sigma_2^2 \\ E(MS_2) &= \sigma^2 + \sigma_{123}^2 &+ 32\sigma_{12}^2 + 20\sigma_{22}^2 + 640\sigma_2^2 \\ E(MS_{12}) &= \sigma^2 + \sigma_{123}^2 &+ 32\sigma_{13}^2 \\ E(MS_{22}) &= \sigma^2 + \sigma_{123}^2 &+ 20\sigma_{23}^2 \\ E(MS_{12}) &= \sigma^2 + \sigma_{123}^2 &+ 20\sigma_{23}^2 \end{split}$$

$$F_{19.589} = \frac{MS_1}{MS_{12}}. \qquad F_{31.589} = \frac{MS_2}{MS_{12}}$$

$$F_{1,j_{2}} = \frac{MS_{3}}{MS_{13} + MS_{23} - MS_{123}}$$

where  $f_2$  is approximately

$$\frac{(MS_{13} + MS_{23} - MS_{123})^2}{\frac{MS_{13}^2}{133} + \frac{MS_{23}^2}{217} + \frac{MS_{123}^2}{4123}}$$

(see Brownlee, 1965, p. 301).

$$F_{589\ 4123} \ge \frac{MS_{12}}{MS_{123}}$$
,

there being no exactly appropriate F for testing the interaction of raters with ratees.

$$\begin{split} F_{133.4123} &= \frac{MS_{13}}{MS_{123}}, \qquad F_{217.4123} &= \frac{MS_{23}}{MS_{123}}, \\ \delta_1^2 &= (MS_1 - MS_{12})/256, \\ \delta_2^2 &= (MS_2 - MS_{12})/160, \\ \delta_3^3 &= (MS_3 - MS_{13} - MS_{23} + MS_{123})/640, \\ \delta_{12}^1 &\geq (MS_{12} - MS_{123})/8, \\ \delta_{13}^2 &= (MS_{12} - MS_{123})/22, \\ \delta_{23}^2 &= (MS_{23} - MS_{123})/20, \\ \delta_{231}^2 &= (MS_{23} - MS_{123})/20, \\ \delta_{231}^2 &= (MS_{23} - MS_{123})/20, \\ \delta_{231}^2 &= (MS_{23} - MS_{123})/20, \\ \delta_{231}^2 &= (MS_{23} - MS_{123})/20, \\ \delta_{231}^2 &= (MS_{23} - MS_{123})/20, \\ \delta_{232}^2 &= (MS_{23} - MS_{23})/20, \\ \delta_{231}^2 &= (MS_{23} - MS_{23})/20, \\ \delta_{232}^2 &= (MS_{23} - MS_{23})/20, \\ \delta_{232}^2 &= (MS_{23} - MS_{23})/20, \\ \delta_{233}^2 &= (MS_{23} - MS_{23})/20, \\ \delta_{23$$

- 7. a. The two nested factors are raters (nested within political party of rater) and ratees (nested within political party of ratee). It happens here that the political parties are the same for raters and ratees. This is a doubly nested design.
  - b. Yes. Raters cannot cross political party of rater, and ratees cannot cross political party of ratee, but they can cross everything else. See Stanley (1961a) for further details.
  - c. "Political party of rater" crosses everything except "rater."
  - d. Rater and ratee are likely to be considered random-effects factors.
  - e. As you probably surmised, Democrat raters tended to rate Democrat raters much higher than they rated Republican ratees, and Republican raters tended to rate Republican ratees much higher than they rated Democrat rates. Thus raters tended to prefer the ratees who were prominent in their own party.
  - f. The interaction of party of rater with party of ratee with trait rated suggests that raters' bias in favor of their own party was not uniform across traits Raters tended to rate ratees of their own party relatively higher on some traits than on others, and similarly for ratees of the other party.



# VERIFICATION OF STANDARD SOLUTION TO LEAST-SQUARES CRITERION

See page 149 first. We begin here with

$$\hat{Y}_i = b_1 X_i + b_0 = b_1 X_i + (\bar{Y}_i - b_1 \bar{X}_i) = b_1 (X_i - \bar{X}_i) + \bar{Y}_i$$

Then consider adding the constant c to  $b_1$  and the constant d to  $b_0$ , where c and d may be any real numbers, negative, zero, or positive:

$$\hat{Y}'_{t} = (b_{1} + c)X_{t} + [(\bar{Y}_{t} - b_{1}\bar{X}_{t}) + d] = [b_{1}(X_{t} - \bar{X}_{t}) + \bar{Y}_{t}] + (cX_{t} + d).$$

We need to show that  $\sum_{i=1}^{n} (Y_i - \hat{Y}_i)^2 \le \sum_{i=1}^{n} (Y_i - \hat{Y}_i)^2$ . By substituting for  $\hat{Y}_i$  and  $\hat{Y}'_i$  their values shown above, we secure the following inequality to be proved:

$$\sum_{i=1}^{n} [Y_i - b_i(X_i - \bar{X}) - \bar{Y}]^2 \le \sum_{i=1}^{n} \{ [Y_i - b_i(X_i - \bar{X}) - \bar{Y}] - (cX_i + d) \}^2.$$

After squaring and summing, we have

$$\begin{split} &\sum_{i=1}^{n} \{Y_i - b_i(X_i - \bar{X}) - \bar{Y}_i\}^{\epsilon} \leq \sum_{i=1}^{n} \{Y_i - b_i(X_i - \bar{X}) - \bar{Y}_i\}^{\epsilon} \\ &+ \sum_{i=1}^{n} \{cX_i + d\}^{\epsilon} - 2 \sum_{i=1}^{n} \{(Y_i - \bar{Y}_i) - b_i(X_i - \bar{X}_i)\}(cX_i + d) \cdot \\ \end{split}$$

571

Subtract  $\sum_{i=1}^{n} [Y_i - b_i(X_i - \overline{X}) - \overline{Y}]^2$  from both sides and obtain

$$0 \leq \sum_{i=1}^{n} (cX_{i} + d)^{2} - 2\sum_{i=1}^{n} [(Y_{i} - \overline{Y}) - b_{i}(X_{i} - \overline{X})](cX_{i} + d).$$

 $\sum_{i=1}^{n} (cX_i + d)^2$  cannot be negative. To complete the proof we shall show in a straightforward manner that, when  $b_1 = s_-/s_-^2$ .

$$\sum_{i=1}^{n} [(Y_{i} - \bar{Y}_{i}) - b_{i}(X_{i} - \bar{X}_{i})](cX_{i} + d) = 0.$$

First, express  $cX_i + d$  as  $(cX_i + d - c\overline{X}) + c\overline{X}$ , which is equivalent to  $c(X_i - \overline{X}) + (c\overline{X} + d)$ . Then

$$\begin{split} &\sum_{i=1}^{n} \left[ (Y_{t} - \bar{Y}_{i}) - b_{1}(X_{i} - \bar{X}_{i}) \right] \left[ c(X_{t} - \bar{X}_{i}) + (c\bar{X}_{i} + d) \right] \\ &= c \sum_{i=1}^{n} (X_{i} - \bar{X}_{i})(Y_{t} - \bar{Y}_{i}) + (c\bar{X}_{i} + d) \sum_{i=1}^{n} (Y_{t} - \bar{Y}_{i}) - b_{1}c \sum_{i=1}^{n} (X_{t} - \bar{X}_{i})^{2} \\ &- b_{1}(c\bar{X}_{i} + d) \sum_{i=1}^{n} (X_{i} - \bar{X}_{i}) \\ &= c(n - 1)s_{xy} + (c\bar{X}_{i} + d)0 - \frac{s_{xy}}{s_{x}^{2}}c(n - 1)s_{x}^{2} - b_{1}(c\bar{X}_{i} + d)0 \\ & \stackrel{>}{=} c(n - 1)s_{xy} + 0 - c(n - 1)s_{xy} - 0 = 0. \end{split}$$

Note that this is equivalent to saying, as in Sec. 9.4, that the covariance of the  $X_i$ 's—or of the linear transformation of the  $X_i$ 's, which is  $cX_i + d$  here—with the discrepancies between the actual  $Y_i$ 's and the  $Y_i$ 's predicted from the  $X_i$ 's via  $b_1$  and  $b_0$  is always exactly zero. Thus the correlation between initial status,  $X_i$  or  $cX_i + d$ , and the  $Y_i - \hat{Y}_i$  is zero.

Therefore,

$$0\leq \sum_{i=1}^n(cX_i+d)^2,$$

which is a correct statement. This completes the proof.

Thus, if one were to use as his regression-slope coefficient  $b_1 + c$  rather than  $b_1 = s_{xy}/s_x^2$ , and as his intercept  $b_0 + d$  rather than  $b_0 = \overline{Y} - b_1\overline{X}$ , he would increase the sum of the squared errors of estimate by the nonnegative amount  $\sum_{i=1}^{n} (cX_i + d)^2$ .

This proves the contention, but it does not show how  $b_1$  and  $b_0$  were derived (via the differential calculus) in the first place. That derivation appears in most textbooks of basic mathematical statistics.

# BIBLIOGRAPHY

- Acton, F. S. Analysis of Straight-line Data. New York: John Wiley, 1959.
- Adkins, Dorothy. Statistics. Columbus, Ohio: Charles E. Merrill, 1964.
- Arkin, H. and R. R. Colton. Graphs: How to Make and Use Them. New York:
- Asher, John W. and Marian N. Schusler. "Students' grades and access to cars." Journal of Educational Research, 60 (1967), 435-37.
- Baker, Frank B. "An investigation of the sampling distributions of item discrimination indices." Psychometrika, 30 (1965), 165-78.
- "Experimental design considerations associated with large-scale research projects. Improving Experimental Design and Statistical Analysis, ed. Julian C. Stanley. Chicago: Rand McNally, 1967.
- Bartlett, M. S. "The effect of non-normality on the t distribution." Proceedings of the Cambridge Philosophical Society, 31 (1935), 223-31.
- Society of London, Series A, 160 (1937), 268-82.
- Bennett, Carl A. and Norman Franklin. Statistical Analysis in Chemistry and the Chemical Industry. New York: John Wiley, 1954.
- Bereiter, Carl E. "Some persisting dilemmas in the measurement of change." Problems in Measuring Change, ed. Chester W. Harris. Madison, Wisconsin: University of Wisconsin Press, 1963.

- Binder, Arnold. "The choice of an error term in analysis of variance design." Psychometrika, 20 (1955), 29-50.
- ——. "Considerations of the place of assumptions in correlational analysis." American Psychologist, 14 (1959), 504-10.
- "Further considerations on testing the null hypothesis and the strategy and tactics of investigating theoretical models." Psychological Review, 70 (1963), 107-15.
- Blalock, Hubert M., Jr. Social Statistics. New York: McGraw-Hill, 1960.
- Causal Inferences in Nonexperimental Research. Chapel Hill: University of North Carolina Press, 1964.
- Bock, R. Darrell, "Multivariate analysis of variance of repeated measurements." Problems in Measuring Change, ed. Chester W. Harris. Madison, Wisconsin: University of Wisconsin Press, 1963.
- "Contributions of multivariate experimental designs to educational research." Handbook of Multivariate Experimental Psychology, ed. Raymond B. Cattell. Chicago: Rand McNally, 1967.
- and Ernest A. Haggard. "The use of multivariate analysis of variance in behavioral research." Handbook of Measurement and Assessment in Behavioral Sciences, ed. Dean K. Whitla. Reading, Massachusetts: Addison-Wesley, 1968.
- Bolch, B. W. "More on unbiased estimation of the standard deviation." American Statistician, 20 (June, 1968), 27.
- Boneau, C. A. "The effects of violations of assumptions underlying the t-test," Psychological Bulletin, 57 (1960), 49-64.
- Boring, Edwin G. "The logic of the normal law of error in mental measurement." American Journal of Psychology, 31 (1920), 1-33.
- Bouvier, E. A., Norman C. Perry, William B. Michael, and A. F. Hertzka. "A study of the error in the cosine-pi approximation to the tetrachoric coefficient of correlation." Educational and Psychological Measurement, 14 (1954), 690-99.
- Box, George E. P. "Non-normality and tests on variances." Biometrika, 40 (1953), 318-35.
- "Some theorems on quadratic forms applied in the study of analysis of variance problems. I, Effect of inequality of variance in the one-way classification." Annals of Mathematical Statistics, 25 (1934a), 290-302.
- "Some theorems on quadratics forms applied in the study of analysis of variance problems. II, Effects of inequality of variance and of correlation between errors in the two-way classification." Annals of Mathematical Statistics, 25 (1954b), 484–98.
- Box, G. E. P. and Anderson, S. L. "Permutation theory in the derivation of robust criteria and the study of departures from assumption." *Journal of the Royal Statistical Society*, Series B, 17 (1955), 1-26.
- Bracht, Glenn H. and Gene V. Glass. "The external validity of experiments." American Educational Research Journal, 5 (1968), 437-74.
- Braithwaite, Richard B. Scientific Explanation. Cambridge, England: Cambridge University Press, 1953.

- Bravais, A. "Analyse mathématique sur les probabilités des erreurs de situation d'un point." Mémoires Présentés par Divers Savants à L'Académie des Sciences de L'Institut de France, 9 (1846), 255-332.
- Brown, George I. "The relationship between barometric pressure and relative humidity and classroom behavior." Journal of Educational Research, 57 (1964),
- Brownlee, K. A. Statistical Theory and Methodology, 2nd Ed. New York; John Wiley, 1965.
- Burke, Cletus J. "Letter to the editor on Peters' reply to Lewis and Burke." Psychological Bulletin, 48 (1951), 81-82.
- -. "A brief note on one-tailed tests." Psychological Bulletin, 50 (1953), 384-87.
- "Further remarks on one-tailed tests." Psychological Bulletin, 51 (1954).
- Campbell, Donald T. "Factors relevant to the validity of experiments in social settings." Psychological Bulletin, 54 (1957), 297-312.
- ------ and Julian C. Stanley. "Experimental and quasi-experimental designs for research on teaching." Handbook of Research on Teaching, ed. N. L. Gage. Chicago: Rand McNally, 1963, pp. 171-246. Also appears as Experimental
- and Quasi-experimental Designs for Research. Chicago: Rand McNally, 1966. Carroll, John B. "The nature of the data, or how to choose a correlation coefficient." Psychometrika, 26 (1961), 347-72.
- Clark, Cherry A. "Hypothesis testing in relation to statistical methodology."
- Review of Educational Research, 33 (1963), 455-73. Cochran, William G. "Some consequences when the assumptions for the analysis
- of variance are not satisfied." Biometrics, 3 (1947), 22-38.
- "The comparison of percentages in matched samples." Biometrika, 37
- ———. "Testing a linear relation among variances." Biometrics, 7 (1951), 17–32. — Sampling Techniques. New York: John Wiley, 1963.
- ----- and Gertrude M. Cox. Experimental Designs, 2nd ed. New York: John
  - Coleman, James S. et al. "Equality of Educational Opportunity." Washington, D.C.: U.S. Government Printing Office, 1966.
- Comfield, Jerome and John W. Tukey. "Average values of mean squares in
- factorials," Annals of Mathematical Statistics, 27 (1956), 907-49. Cox, David R. "The use of a concomitant variable in selecting an experimental
- design." Biometrika, 44 (1957), 150-58.
- —. Planning of Experiments. New York: Wiley, 1958.
- "A remark on multiple comparison methods." Technometrics, 7 (1965),
- Cronbach, Lee J. "The two disciplines of scientific psychology." American
- Cureton, Edward E. "Rank-biserial correlation." Psychometrika, 21 (1956),

- Cureton, Edward E. "Note on  $\phi/\phi$  max." Psychometrika, 24 (1959), 89-91.
- "Rank-biserial correlation when ties are present." Educational and Psychological Measurement, 28 (1968a), 77-79.
- 22 (February, 1968b), 22.
- ... "Priority correction to 'Unbiased estimation of the standard deviation." American Statistician, 22 (June, 1968c), 27.
- David, Florence N. and N. L. Johnson. "The effect of non-normality on the power
- function of the F-test in the analysis of variance." Biometrika, 38 (1951). 43-57. Davidoff, M. D. and Howard W. Goheen. "A table for the rapid determination of the tetrachoric correlation coefficient." Psychometrika, 18 (1953), 115-21.
- Delacato, C. H. Neurological Organization and Reading. Springfield, Illinois: Thomas, 1966.
- Diamond, Solomon. Information and Error. New York; Basic Books, 1960.
- ---. The World of Probability. New York: Basic Books, 1965.
- Dispensa, J. "Relationship of the thyroid with intelligence and personality." Journal of Psychology, 6 (1938), 181-86.
- Dixon, Wilfred J. and Frank J. Massey. Introduction to Statistical Analysis, 2nd ed, New York; McGraw-Hill, 1957. (3rd edition, 1969)
- Dizney, Henry F, and Lauren Gromen. "Predictive validity and differential achievement on three MLA Comparative Foreign Language Tests." Educational and Psychological Measurement, 27 (1967), 1127-30.
- Draper, Norman R. and H. Smith. Applied Regression Analysis. New York: John Wiley, 1966.
- DuBois, Philip H. Multivariate Correlational Analysis. New York: Harper, 1957.
- Dunçan, David B. "Multiple range and multiple F tests." Biometrics, 11 (1955),
- ..... "A Bayesian approach to multiple comparisons." Technometrics, 7 (1965), 171-222.
- Dunn, Olive J. "Multiple comparisons among means." Journal of the American Statistical Association, 56 (1961), 52-64.
- Dunnett, C. W. "A multiple comparison procedure for comparing several treatments with a control." Journal of the American Statistical Association, 50 (1955), 1096-1121.
- Durbin, J. and Alan Stuart. "Inversions and rank correlation coefficients." Journal of the Royal Statistical Society, Series B, 13 (1951), 303-9.
- Dyson, Ernest. "A study of ability grouping and the self-concept." Journal of Educational Research, 60 (1967), 403-5.
- Edwards, Allen L. "On 'the use and misuse of the chi-square test'-the case of the 2 × 2 contingency table." Psychological Bulletin, 47 (1950), 341-46.
- New York: John Wiley, 1964.

- Edwards, Allen L. Experimental Design in Psychological Research, 3rd ed. New York: Holt, Rinehart and Winston, 1968. Edwards, Ward, Harold Lindman, and Leonard J. Savage. "Bayesian statistical
- inference for psychological research." Psychological Review, 70 (1963), 193-242.
- Erlenmeyer-Kimling, L. and L. F. Jarvik. "Genetics and intelligence: a review." Science, 142 (1963), 1477-79.
- Ezekiel, M. and Karl A. Fox. Methods of Correlation and Regression Analysis, 3rd ed. New York: John Wiley, 1959. (4th edition, 1963)
- Feller, William. An Introduction to Probability Theory and Its Application, Volume I, 2nd ed. New York: John Wiley, 1957.
- Ferguson, George A. Statistical Analysis in Psychology and Education, 2nd ed. New York: McGraw-Hill, 1966. (First edition, 1959)
- Fisher, Ronald A. "On the 'probable error' of a coefficient of correlation deduced from a small sample." Metron, 1, Part 4 (1921), 3-32.
- —. Statistical methods for research workers, eds. 1-13. Edinburgh: Oliver and Boyd, 1925-58.
- The Design of Experiments, eds. 1-8. Edinburgh: Oliver and Boyd, 1935-66.
- Statistical Methods and Scientific Inference, 2nd ed. New York: Hafner, 1959.
- ——and Frank Yates. Statistical Tables for Biological, Agricultural, and Medical Research, 2nd ed. Edinburgh: Oliver and Boyd, 1943.
- Forchand, G. A. and W. L. Libby, Jr. "Effects of educational programs and perceived organizational climate upon changes in innovative administrative behavior." In Innovative Behavior. Chicago: University of Chicago Center for Programs in Government Administration, 1962.
  - Fryer, H. C. Concepts and Methods of Experimental Statistics. Boston: Allyn and Bacon, 1966.
- Furfey, Paul H. "Comment on 'the needless assumption of normality in Pearson's r." American Psychologist, 13 (1958), 545-46.
- Gayen, A. K. "The distribution of 'Student's' t in random samples of any size drawn from non-normal universes." Biometrika, 36 (1949), 353-69.
- ——. "The distribution of the variance ratio in random samples of any size drawn from non-normal universes." Biometrika, 37 (1950a), 236-55.
  - "Significance of difference between the means of two non-normal samples."

    Biometrika, 37 (1950b), 399-403.
- Geary, R. C. "The distribution of 'student's' ratio for non-normal samples." Journal of the Royal Statistical Society, Series B, Supplement (1936), 178-84.
  Glass, Gene V. "Testing benefit of the Property of the P
- Glass, Gene V. "Testing homogeneity of variances." American Educational Research Journal, 3 (1966a), 187–90.
- "Note on rank-biserial correlation." Educational and Psychological Measurement, 26 (1966b), 623-31.
- and James R. Collins. "Geometric proof of the restriction on the

- possible values of  $r_{xy}$  when  $r_{xz}$  and  $r_{yz}$  are fixed." Educational and Psychological Measurement, 29 (1969).
- Glass, Gene V and A. Ralph Hakstian. "Measures of association in comparative experiments: their development and interpretation." American Educational Research Journal, 6 (1969), 403-14.
- Godard, R. H. and E. F. Lindquist. "An empirical study of the effect of heterogeneous within-groups variance upon certain F-tests of significance in analysis of variance." Psychometrika, 5 (1940), 263-74.
- Goldfried, M. A. "One-tailed tests and 'unexpected' results." Psychological Review, 66 (1959), 79-80.
- Goodman, Leo A. "Simultaneous confidence intervals for contrasts among multinomial populations." Annals of Mathematical Statistics, 35 (1964) 716– 25.
- Goolsby, T. M. "Comparability and validity of three forms of SCAT." Educational and Psychological Measurement, 27 (1967), 1041-45.
- Grant, David A. "Testing the null hypothesis and the strategy and factics of investigating theoretical models." Psychological Review, 69 (1962), 54-61.
- Graybill, Franklin A. An Introduction to Linear Statistical Models, Volume 1. New York: McGraw-Hill, 1961.
- Green, Bert F. and John W. Tukey. "Complex analysis of variance: general problems." Psychometrika, 25 (1960), 127-52.
- Greenhouse, S. W. and S. Geisser. "On methods in the analysis of profile data." Psychometrika, 24 (1959), 95-112.
- Gronow, D. G. C. "Test for the significance of difference between means in two normal populations having unequal variances." Biometrika, 38 (1951), 252-56.
- Guenther, W. C. Analysis of Variance. Englewood Cliffs, N.J.: Prentice-Hall, 1964.
- ----. Concepts of Statistical Inference. New York: McGraw-Hill, 1965.
- Guthrie, John T. "Expository instruction versus a discovery method." Journal of Educational Psychology, 58 (1967), 45-49.
- Haggard, Ernest A. Intra-class Correlation and the Analysis of Variance. New York: Dryden Press, 1958.
- Harrington, Scott A. "Sequencing organizers in meaningful verbal learning." Research Paper No. 10. Boulder: University of Colorado, Laboratory of Educational Research, 1968.
- Harris, Chester W. Problems in Measuring Change. Madison, Wisconsin: University of Wisconsin Press, 1963.
- Harter, H. Leon. "Tables of range and studentized range." Annals of Mathematical Statistics, 31 (1960), 1122-47.
- Hays, William L. Statistics for Psychologists. New York: Holt, Rinehart and Winston, 1963.
- Henderson, C. R. "Design and analysis of animal husbandry experiments," In Techniques and Procedures of Animal Production Research. American Society of Animal Production, Belisville, Maryland, 1959.

- Herman, William L. "Teaching attitude as related to academic grades and athletic ability of prospective physical education teachers." Journal of Educational Research, 61 (1967), 40-42,
- Hick, W. E. "A note on one-tailed and two-tailed tests." Psychological Review, 59 (1952), 316-18.
- Hinton, R. T., Jr. "A further study on the role of the basal metabolic rate in the intelligence of children." Journal of Educational Psychology, 30 (1939), 309-14. Hopkins, Kenneth D. "An empirical analysis of the efficacy of the WISC in the
- diagnosis of organicity in children of normal intelligence." Journal of Genetic Psychology, 105 (1964), 163-72.
  - and Russel A. Chadbourn. "A scheme for proper utilization of multiple comparisons in research and a case study." American Educational Research Journal, 4 (1967), 407-12,
  - Horsnell, G. "The effect of unequal group variances on the F-test for the homogeneity of group means," Biometrika, 40 (1953), 128-36.
  - Horst, Paul. "A proof that the point from which the sum of the absolute deviations is a minimum is the median." Journal of Educational Psychology, 22 (1931), 463-64.
  - ... Matrix Algebra for Social Scientists. New York: Holt, Rinehart and Winston, 1963,
  - Huff, Darrell. How to Lie With Statistics. New York: Norton, 1954.
  - Jarrett, Rheem F. "A minor exercise in history." American Statistician, 22 (June, 1968), No. 3, 25-26.
  - Jaspen, Nathan. "Serial correlation." Psychometrika, 11 (1946), 23–30.
  - Jeffrey, W. E. and S. Jay Samuels. "Effect of method of reading training on initial learning and transfer." Journal of Verbal Learning and Verbal Behavior, 6 (1967), 354-58.
  - Jenkins, William L. "An improved method for tetrachoric r." Psychometrika,
  - Jensen, Arthur R. "Social class, race, and genetics: implications for education." American Educational Research Journal, 5 (1968), 1-42.
  - Johnson, Palmer O. Statistical Methods in Research. Englewood Cliffs, N.J.:
  - Jones, Lyle V. "Tests of hypotheses: one-sided vs. two-sided alternatives." Psychological Bulletin, 49 (1952), 43-46.
    - --. "A rejoinder on one-tailed tests." Psychological Bulletin, 51 (1954),
  - Kaiser, Henry F. "Directional statistical decisions." Psychological Review, 67
    - Kaplan, Abraham. The Conduct of Inquiry. San Francisco: Chandler, 1964.
    - Keeping, E. S. Introduction to Statistical Inference. Princeton, N.J.: Van
  - Kelley, Truman L. "A new measure of dispersion." Publications of the American Statistical Association, 17 (1921), 743-49.

- Kelley, Truman L. "The principles and techniques of mental measurement." American Journal of Psychology, 34 (1923), 408-32.
- ----- Fundamentals of Statistics. Cambridge, Massachusetts; Harvard University Press, 1947.
- Kendall, Maurice G. Rank Correlation Methods, 3rd ed. London: Griffin, 1962.
- and William Buckland. A Dictionary of Statistical Terms. Edinburgh; Oliver and Boyd, 1957,
- and Alan Stuart. The Advanced Theory of Statistics, Volumes 1-3. London; Griffin, 1961-66.
- Kennedy, W. A., V. Van de Riet, and J. White. "A normative sample of intelligence and achievement of Negro elementary school children in the Southeastern United States." Monographs of the Society for Research in Child Development, 28 (1963), No. 6.
- Kenyon, Gerald S. "Multiple comparisons and the analysis of variance: an empirical illustration." Research Quarterly of the American Association for Health, Physical Education, and Recreation, 36 (1965), 413-19.
- Kerlinger, Fred N. Foundations of Behavioral Research, New York; Holt, Rinehart and Winston, 1964.
- Keuls, M. "The use of the 'studentized range' in connection with an analysis of variance." Euphytica, 1 (1952), 112-22.
- Kimmel, Herbert D. "Three criteria for the use of one-tailed tests." Psychological Bulletin, 54 (1957), 531-33.
- Klausmeier, Herbert J. "Effects of accelerating bright older elementary pupils." Journal of Educational Psychology, 54 (1963), 165-71.
- Kruskal, William H. "Ordinal measures of association." Journal of the American Statistical Association, 53 (1958), 814-61.
- ... "Tests of significance." International Encyclopedia of the Social Sciences, Vol. 14. ed. David L. Sills. New York: Macmillan and Free Press, 1968.
- LaForge, R. "Comment on 'the needless assumption of normality in Pearson's r." American Psychologist, 13 (1958), 546.
- Lana, Robert D. and A. Lubin. "The effect of correlation on the repeated measures design." Educational and Psychological Measurement, 23 (1963), 729-39.
- Lancaster, H. O. and M. A. Hamden. "Estimate of the correlation coefficient in contingency tables with possibly nonmetrical characters." Psychometrika, 29 (1964), 383-91.
- Lerner, Daniel (ed.) Cause and Effect. New York: Free Press, 1965.
  - Levene, H. "Robust tests for equality of variances." Contributions to Probability and Statistics, ed. I. Olkin. Stanford, California: Stanford University Press. 1960, pp. 278-92.
  - Lewis, Donald and Cletus J. Burke. "The use and misuse of the chi-square test." Psychological Bulletin, 46 (1949), 433-89.
    - "Further discussion on the use and misuse of the chi-square test." Psychological Bulletin, 47 (1950), 347-55.
  - Leyman, Laretha. "Prediction of freshman and sophomore grade-point averages

k

of women physical education major students." Educational and Psychological Measurement, 27 (1967), 1139-41.

- .-- -- ---

- Lindquist, E. F. Statistical Analysis in Educational Research. New York: Houghton Mifflin, 1940.
- Design and Analysis of Experiments in Psychology and Education. Boston:
   Houghton Mifflin, 1953.
- Lord, Frederic M. "Biserial estimates of correlation." Psychometrika, 28 (1963a), 81-85.
  - "Elementary models for measuring change." Problems in Measuring Change, ed. C. W. Harris. Madison, Wisconsin: University of Wisconsin Press. 1954). 21-38.
    - and Melvin R. Novick. Statistical Theories of Mental Test Scores. Reading, Massachusetts: Addison-Wesley, 1968.
  - Lubin, Ardie. "The interpretation of significant interaction." Educational and Psychological Measurement, 21 (1962a), 807-17.
  - Marascuilo, Leonard A. "Large sample multiple comparisons." Psychological Bulletin, 65 (1966), 280-90.
  - Marks, Melvin R. "Two kinds of experiment distinguished in terms of statistical operations." Psychological Review, 58 (1951), 179-84.

  - McHugh, Richard B. and D. S. Ellis. "The 'post-mortem' testing of experimental comparisons." Psychological Bulletin, 52 (1955), 425-28.
  - McLean, Leslie D. "Phantom classrooms." School Review, 74 (1966), 139-49.
  - "Some important principles for the use of incomplete designs in behavioral research." Improving Experimental Design and Statistical Analysis, ed. Julian C. Stanley. Chicago: Rand McNally, 1967.
  - McNamara, Walter J. and J. W. Dunlap. "A graphical method for computing the standard error of biserial r," Journal of Experimental Education, 2 (1934), 274-77.
  - McNemar, Quinn. "Note on the sampling error of the difference between correlated proportions or percentages." Psychometelka, 12 (1947), 153-57.
  - ----. Psychological Bulletin, 37 (1940), 747.
  - Psychological Statistics, 3rd ed. New York: John Wiley, 1962.
  - Michael, William B., Henry F. Kaiser, and Cherry Ann Clark. "Research tools: statistical methods." Review of Educational Research, 27 (1957), 498-527.
  - Milholland, John E. "Comment on 'the needless assumption of normality in Pearson's r." American Psychologist, 13 (1958), 544-45.
    - Miller, Rupert G. Simultaneous Statistical Inference. New York: McGraw-Hill, 1966.
    - Millman, Jason and Gene V. Glass. "Rules of thumb for writing the ANOVA table." Journal of Educational Measurement, 4 (1967), 41-51.
    - Mises, Richard von. Probability, Statistics, and Truth, 2nd ed. New York: Macmillan, 1957.

- Mood, Alexander and Franklin A. Graybill. Introduction to the Theory of Statistics, 2nd ed. New York: McGraw-Hill, 1963.
- Moses, Lincoln E. "Statistical theory and research design." Annual Review of Psychology, 7 (1956), 233-58.
- Nefzger, M. D. and James Drasgow. "The needless assumption of normality in Pearson's r." American Psychologist, 12 (1957), 623-25.
- Newman, D. "The distribution of range in samples from a normal population, expressed in terms of an independent estimate of standard deviation." Biometrika, 31 (1939), 20-30.
- Norris, Raymond C. and Howard F. Hjelm. "Non-normality and product moment correlation." Journal of Experimental Education, 29 (1961), 261-70.
- North, George E. and O. Lee Buchanan. "Teacher views of poverty area children." Journal of Educational Research, 61 (1967), 53-55.
- Nunnally, J. C. "The place of statistics in psychology." Educational and Psychological Measurement, 20 (1960), 641-50.
- Ohnmacht, Fred W. "Achievement, anxiety, and creative thinking." American Educational Research Journal, 3 (1966), 131-38.
  - Olds, E. G. "Distributions of sums of squares of rank differences for small numbers of individuals." Annals of Mathematical Statistics, 9 (1938), 133-48.
- "The 5% significance levels for sums of squares of rank differences and a correction." Annals of Mathematical Statistics, 20 (1949), 117-18.
- Olkin, Ingram. "Correlations revisited." Improving Experimental Design and Statistical Analysis, ed. Julian C. Stanley. Chicago: Rand McNally, 1967.
- and John W. Pratt. "Unbiased estimation of certain correlation coefficients." Annals of Mathematical Statistics, 29 (1958), 201-11.
- ----- and M. Siotani. "Asymptotic distribution functions of a correlation matrix." Stanford, California: Stanford University Laboratory for Quantitative Research in Education, Report No. 6, 1964.
- Pastore, Nicholas. "Some comments on the use and misuse of the chi-square test." Psychological Bulletin, 47 (1950), 338-40.
- Pearson, Egon S. "The analysis of variance in cases of non-normal variation." Biometrika, 23 (1931), 114-33.
- and C. J. Clopper. "The use of confidence intervals or fiducial limits illustrated in the case of the binomial." Biometrika, 26 (1934), 404-13.
- and H. O. Hartley. Biometrika Tables for Statisticians, 3rd ed. Cambridge, England: Cambridge University Press, 1966.
- "Charts of the power function for analysis of variance tests, derived from the non-central F-distribution." Biometrika, 38 (1951), 112-30.
- Pearson, Karl. "On further methods of determining correlation." Drapers' Company Memoirs, Biometric Series, IV (1907).
- Peckham, Perc D., Gene V Glass, and Kenneth D. Hopkins. "The experimental unit in statistical analysis: comparative experiments with intact groups." Journal of Special Education, 3 (1969).
- Peizer, David B. "A note on directional inference." Psychological Bulletin, 68 (1967), 448.

- Peters, C. C. "The misuse of chi-square—a reply to Lewis and Burke." Psychological Bulletin, 47 (1950), 331-37.
- Pitman, E. J. G. "Significance tests which may be applied to samples from any populations. III: The analysis of variance test." *Biometrika*, 29 (1937), 322-35.
- Platt, John R. "Strong inference." Science, 146 (1964), 347-53.
- Pugh, R. C. "The partitioning of criterion-score variance accounted for in multiple correlation." American Educational Research Journal, 5 (1968), 639-46.
- RAND Corporation. A Million Random Digits with 100,000 Normal Deviates. New York: Free Press, 1955.
- Rozeboom, W. W. "The fallacy of the null-hypothesis significance test." Psychological Bulletin, 57 (1960), 416-28.
- Foundations of the Theory of Prediction. Homewood, Illinois; Dorsey Press, 1966.
- Rugg, H. O. Statistical Methods Applied to Education. Boston: Houghton Mifflin, 1917.
- Ryan, Thomas A. "Multiple comparisons in psychological research." Psychological Bulletin, 56 (1959), 26–47.
  - "The experiment as the unit for computing rates of error." Psychological Bulletin, 59 (1962), 301-5.
- Samuels, S. Jay. "Attentional process in reading: the effect of pictures on the acquisition of reading responses." Journal of Educational Psychology, 58 (1967), 337-42.
- Sawrey, W. L. "A distinction between exact and approximate nonparametric methods." Psychometrika, 23 (1958), 171-77.
- Scannell, D. P. and J. C. Marshall. "The effect of selected composition errors on grades assigned to essay examinations." American Educational Research Journal, 3 (1966), 125–30.
- Scates, Douglas E. "Reporting, summarizing, and implementing educational research." Review of Educational Research, 12 (1942), 558-74.
- Scheffe, Henry. The Analysis of Variance. New York: John Wiley, 1959.
- Schultz, E. F., Jr. "Rules of thumb for determining expectations of mean squares in analysis of variance." Biometrics, 11 (1955), 123-35.
- Sears, Pauline S. "Levels of aspiration in academically successful and unsuccessful children." Journal of Abnormal and Social Psychology, 35 (1940), 498-536.
- Senders, Virginia L. Measurement and Statistics. London: Oxford University Press, 1958.
- Sigel, Sidney. Nanparametric Statistics for the Behavioral Sciences. New York: McGraw-Hill, 1956.

- Snedecor, George W. and William G. Cochran. Statistical Methods, 6th ed. Ames, Iowa: Iowa State University Press, 1967.
- Soper, H. E. "On the probable error of the bi-serial expression for the correlation coefficient," Biometrika, 10 (1914), 384-90.
- Sparks, Jack N. "Expository notes on the problem of making multiple comparisons on a completely randomized design." Journal of Experimental Education, 31 (1963), 343-49.
- Spear, Mary E. Charting Statistics. New York: McGraw-Hill, 1952.
- Srivastava, A. B. L. "Effect of non-normality on the power of the analysis of variance test." Biometrika, 46 (1959), 114-22.
- Staff, Harvard University Computational Laboratory. Harvard Tables of the Cumulative Binomial Distribution. Cambridge, Massachusetts: Harvard University Press, 1956.
- Stanley, Julian C. "Additional 'post-mortem' tests of experimental comparisons." Psychological Bulletin, 54 (1957a), 128-30.
  - "Index of means vs. mean of indices." American Journal of Psychology, 70 (1957b), 467-68.
- "Analysis of a double nested design." Educational and Psychological Measurement, 21 (1961a), 831-37. Errata, 22 (1962), ii.
- "Analysis of unreplicated three-way classifications, with applications to rater bias and trait independence." Psychometrika, 26 (1961b), 205-19.
- Prentice-Hall, 1964.
- "Quasi-experimentation." School Review, 73 (1965), 197-205.
- -. "A common class of pseudo-experiments." American Educational Research Journal, 3 (1966a), 79-87.
- research thirty years later." American Educational Research Journal, 3 (1966b). 223-29.
- "On improving certain aspects of educational experimentation." Improving Experimental Design and Statistical Analysis, ed. Julian C. Stanley. Chicago: Rand McNally, 1967a.
  - "Elementary experimental design-an expository treatment." Psychology in the Schools, 4 (1967b), 195-203.
- "Problems in equating groups in mental retardation research." Journal of Special Education, 1 (1967c), 241-56.
- "Linear hypotheses. II. Analysis of variance." International Encyclopedia of the Social Sciences, Vol. 9, ed. David L. Sills. New York: Macmillan and Free Press, 1968a, pp. 324-36.
- "An important similarity between biserial r and the Brogden-Cureton-Glass biserial r for ranks." Educational and Psychological Measurement, 28 (1968b), 249-53.
- "Plotting anova interactions for ease of visual interpretation." Educational and Psychological Measurement, 29 (1969a), 793-97.

- Peters, C. C. "The misuse of chi-square—a reply to Lewis and Burke." Psychological Bulletin, 47 (1950), 331-37.
- Pitman, E. J. G. "Significance tests which may be applied to samples from any populations. III: The analysis of variance test." Biometrika, 29 (1937), 322-35.
- Platt, John R. "Strong inference." Science, 146 (1964), 347-53.
- Pugh, R. C. "The partitioning of criterion-score variance accounted for in multiple correlation." American Educational Research Journal, 5 (1968), 639-46.
- RAND Corporation. A Million Random Digits with 100,000 Normal Deviates. New York: Free Press, 1955.
- Rozeboom, W. W. "The fallacy of the null-hypothesis significance test." Psychological Bulletin, 57 (1960), 416-28.
- ----- Foundations of the Theory of Prediction. Homewood, Illinois: Dorsey Press, 1966.
- Rugg, H. O. Statistical Methods Applied to Education. Boston: Houghton Mifflin, 171.
  - Ryan, Thomas A. "Multiple comparisons in psychological research." Psychological Bulletin, 56 (1959), 26-47.
  - "The experiment as the unit for computing rates of error." Psychological Bulletin, 59 (1962), 301-5.
- Samuels, S. Jay. "Attentional process in reading: the effect of pictures on the acquisition of reading responses." Journal of Educational Psychology, 58 (1967), 337-42.
- Sawrey, W. L. "A distinction between exact and approximate nonparametric methods." Psychometrika, 23 (1958), 171-77.
- Scandura, Joseph M. and Jay N. Wells. "Advance organizers in learning abstract mathematics." American Educational Research Journal, 4 (1967), No. 3.
- Scannell, D. P. and J. C. Marshall. "The effect of selected composition errors on grades assigned to essay examinations." American Educational Research Journal, 3 (1966), 125-30.
- Scates, Douglas E. "Reporting, summarizing, and implementing educational research." Review of Educational Research, 12 (1942), 558-74.
- Scheffe, Henry. The Analysis of Variance. New York: John Wiley, 1959.
- Schultz, E. F., Jr, "Rules of thumb for determining expectations of mean squares in analysis of variance." Biometrics, 11 (1953), 123-35.
  - Scars, Pauline S. "Levels of aspiration in academically successful and unsuccessful children." Journal of Abnormal and Social Psychology, 35 (1940), 498-536.
  - Senders, Virginia L. Measurement and Statistics. London: Oxford University Press, 1958.
  - Sigel, Sidney. Nonparametric Statistics for the Behavioral Sciences. New York: McGraw-Hill, 1956.

- Snedecor, George W. and William G. Cochran. Statistical Methods, 6th ed. Ames, Iowa: Iowa State University Press, 1967.
- Soper, H. E. "On the probable error of the bi-serial expression for the correlation coefficient." Biometrika, 10 (1914), 384-90.
- Sparks, Jack N. "Expository notes on the problem of making multiple comparisons on a completely randomized design." Journal of Experimental Education, 31 (1963), 343-49.
- Spear, Mary E. Charting Statistics. New York: McGraw-Hill, 1952.
- Srivastava, A. B. L. "Effect of non-normality on the power of the analysis of variance test." Biometrika, 46 (1959), 114-22.
- Staff, Harvard University Computational Laboratory. Harvard Tables of the Cumulative Binomial Distribution. Cambridge, Massachusetts: Harvard University Press, 1956.
- Stanley, Julian C. "Additional 'post-mortem' tests of experimental comparisons." Psychological Bulletin, 54 (1957a), 128-30.
  - "Index of means vs. mean of indices." American Journal of Psychology, 70 (1957b), 467-68.
- "Analysis of a double nested design." Educational and Psychological Measurement, 21 (1961a), 831-37. Errata, 22 (1962), ii.
- —. "Analysis of unreplicated three-way classifications, with applications to rater bias and trait independence." Psychometrika, 26 (1961b), 205–19.
- Measurement in Today's Schools, 4th ed. Englewood Cliffs, N.J.: Prentice-Hall, 1964.
- ——. "Quasi-experimentation." School Review, 73 (1965), 197-205.
- "A common class of pseudo-experiments." American Educational Research Journal, 3 (1966a), 79-87.
- research thirty years later." American Educational Research Journal, 3 (1966b). 223-29.
- "On improving certain aspects of educational experimentation." Improving Experimental Design and Statistical Analysis, ed. Julian C. Stanley. Chicago: Rand McNally, 1967a.
  - "Elementary experimental design-an expository treatment." Psychology in the Schools, 4 (1967b), 195-203.
  - "Problems in equating groups in mental retardation research." Journal of Special Education, 1 (1967c), 241-56.
- "Linear hypotheses. II. Analysis of variance." International Encyclopedia of the Social Sciences, Vol. 9, ed. David L. Sills. New York: Macmillan and Free Press, 1968a, pp. 324-36.
- "An important similarity between biserial r and the Brogden-Cureton-Glass biserial r for ranks." Educational and Psychological Measurement, 28 (1968b), 249-53.
- "Plotting anova interactions for ease of visual interpretation." Educational and Psychological Measurement, 29 (1969a), 793-97.

- Stanley, Julian C. and Marilyn D. Wang. "Restrictions on the possible values of r<sub>10</sub> given r<sub>13</sub> and r<sub>22</sub>." Educational and Psychological Measurement, 29 (1969b), 579-81.
- Steel, R. G. and J. H. Torrie. Principles and Procedures of Statistics. New York: McGraw-Hill, 1960.
- Stennett, R. G. "Absence from school: norms by sex and grade." Journal of Educational Research, 60 (1967), 351-54.
- Stevens, S. S. (ed). Handbook of Experimental Psychology. New York: John Wiley, 1951.
- "Student." "The probable error of a mean." Biometrika, 6 (1908), 1-25.
- Tables of the cumulative binomial probabilities. Ordnance Corps Pamphlet ORDP 20-1. Washington, D.C.: U.S. Government Printing Office, 1953.
- Tate, Merle W. and R. C. Clelland. Nonparametric and Shortcut Statistics in the Social, Biological, and Medical Sciences. Danville, Illinois: Interstate Printers, 1957
- Terman, Lewis M. and Maud A. Merrill. Stanford-Binet Intelligence Scale:
  Manual for the Third Recision, Form L-M. Boston: Houghton Mifflin, 1960.
- Thalberg, Stanton P. "Reading rate and immediate versus delayed retention." Journal of Educational Psychology, 58 (1967), 373–78.
- Townsend, John C. Introduction to Experimental Method. New York: McGraw-Hill, 1953.
- Travers, Robert M. W., R. K. Van Wagenen, D. H. Haygood, and Mary McCormick. "Learning as a consequence of the learner's task involvement under different conditions of feedback." Journal of Educational Psychology, 55 (1964) 167-73.
- Tucker, L. R., Fred Damarin, and Samuel Messick. "A base-free measure of change." Psychometrika, 31 (1966), 457-73.
- Varberg, D. E. "The development of modern statistics." The Mathematics Teacher, 56 (1963), 252-57, 344-48.
- Walker, Helen M. Studies in the History of Statistical Method. Baltimore: Williams and Wilkins, 1929.
- "Degrees of freedom." Journal of Educational Psychology, 31 (1940),
- Mathematics Essential for Elementary Statistics, revised ed. New York:
  Holt, Ringhart and Wington, 1981
- —— and Walter N. Durost. Statistical Tables: Their Structure and Use. New York: Teachers College, Columbia University, 1936.
- —— and Joseph Lev. Statistical Inference. New York: Holt, Rinehart and Winston, 1953.
- Wallen, Norman E. and Mary Lou A. Campbell. "Vocabulary and non-verbal reasoning components of verbal analogies tests (Miller Analogies Test and Concept Mastery Test)." Journal of Educational Research, 61 (1967), 87-89.
- Wallis, W. Allen and Harry V. Roberts. Statistics: A New Approach. Glencoe, Illinois: Free Press, 1956.
- The Nature of Statistics. New York: Free Press, 1965.

- Webb, Eugene J., Donald T. Campbell, Richard D. Schwartz, and Lee Sechrest.

  Unobtrustice Measures: Nonreactive Research in the Social Sciences. Chicago:
  Rand McNally, 1966.

  Webster, Harold A. and Coal E. Benitse. Webster, Harold A. and Coal E. Benitse.
- Webster, Harold A. and Carl E. Bereiter. "The reliability of changes measured by mental test scores." Problems in Measuring Change, ed. Chester W. Harris. Madison, Wisconsin: University of Wisconsin Press, 1963.
- Welch, B. L. "The significance of the difference between two means when the population variances are unequal." Biometrika, 29 (1937), 350-62.
- Wert, J. E., Charles O. Neidt, and J. Stanley Ahmann. Statistical Methods in Educational and Psychological Research. New York: Appleton-Century-Crofts, 1954.
- Werts, Charles E. "The partitioning of variance in school effects studies," American Educational Research Journal, 5 (1968), 311-18.
- Whitfield, J. W. "Rank correlation between two variables, one of which is ranked, the other dichotomous." Biometrika, 34 (1947), 292-96.
- Wilks, S. S. Mathematical Statistics. New York: John Wiley, 1962.
- Williams, D. L. "Rewritten science materials and reading comprehension." Journal of Educational Research, 61 (1968), 204-6.
- Williams, E. J. "Linear hypotheses. I. Regression." International Encyclopedia of the Social Sciences, Vol. 9, ed. David L. Sills. New York: Macmillan and Free Press, 1968, pp. 310–24.
- Wilson, W. R. "A note on the inconsistency inherent in the necessity to perform multiple comparisons." *Psychological Bulletin*, 59 (1962), 296-300.
- and H. Miller. "A note on the inconclusiveness of accepting the null hypothesis." Psychological Review, 71 (1964), 238-42.
- Wine, R. Lowell. Statistics for Scientists and Engineers. Englewood Cliffs, N.J.: Prentice-Hall, 1964.
- N.J.: Frentice-Hall, 1904.
  Winer, Ben J. Statistical Principles in Experimental Design. New York: McGraw-Hill, 1962.
- Yamamoto, Kaoru. "Creativity and unpredictability in school achievement." Journal of Educational Research, 60 (1967), 321-25.
- Yates, Frank. "Contingency tables involving small numbers and the χ² test." Journal of the Royal Statistical Society, Supplement, 1 (1934), 217-35.

Drasgow, James, 283
DuBois, Philip H., 130, 187
Duncan, David B, 382-83
Dunlap, J, W, 320
Dunnett, C. W., 382
Durbin, J., 178
Durost, Walter N., 54
Dyson, Ernest, 336

Edwards, Allen L., 129, 174, 252, 333, 495, 498, 499

Ellis, D. S., 382 Entermeyer-Kiming, L., 118 Euler, Leonhard, 296

Ezekiel, M., 130, 186

Ferguson, George A., 71, 72, 332 Fermal, Pierre de, 96 Fisher, Ronald A., 250, 265, 297, 403, 491, 492, 499, 520, 521, 536

Forehand, G. A., 309 Fox, Karl A., 130, 186 Franklih, Norman, 472 Furfey, Paul H., 283 Gage, N. L., 508, 509

Galton, Francis, 113, 172
Gauss, Karl, 139, 173
Geisser, S., 370, 470
Getzel, Jacob., 273
Glass, Gene V, 152, 163, 179, 181, 190, 320, 376, 472, 509
Godard, R. H., 371
Goldfried, M. A., 289
Goodman, Lo. D., 226

Goldfried, M. A. 289 Goodman, Leo A., 326 Goolsby, T. M., 194 Gosset, W. S. (Student), 260 Grant, David A., 290 Graybil, Franklin A., 229 Greenhouse, S. W., 370, 470 Gromow, D. G. C., 297 Guenther, W. C., 333, 397, 472 Guthner, John T., 378, 398

Haggard, Ernest A., 152 Handen, M. A., 167 Handen, M. A., 167 Harter, H. Leon, 529-53 Hartley, H. O., 209, 374, 375, 522-28, 538, 548-51 Hartley, H. O., 209, 374, 375, 522-28, 538, 548-51 Hasting, H. O., 209, 374, 375, 522-28, 538, 548-51 Hasting, M. T., 170, 312, 316 Henderson, C. R., 472 Herman, William L., 312, 316 Hock, W. E., 202, 312 Honton, R. T., 172, 312 Honton, R. T., 172, 313 Honton, R. T., 172, 310 Hookins, Kenneth D., 396, 483, 484, 509

Hjelm, Howard F., 283, 310 Hopkins, Kenneth D., 396, 483, 484, 509 Horsneth, G, 371 Horst, Paul, 67 Hovland, C. I., 508 Hsu. P. Le, 371

Huff, Darrell, 52

Jackson, Philip, 273
Jarrett, Rheem, 253
Jarvik, L. F., 118
Jaspen, Nathan, 171
Jeffrey, W. E., 398
Jenkins, William L., 167
Jensen, Arthur R., 191
Johnson, Palmer O, 306
Jones, Lyle V., 289

Jones, Lyle V., 289

Kabot, Ruth R., 193

Kasser, Henry F., 12, 285

Kapfan, Abraham, 14

Kepping, E. S., 63, 333

Kelley, Truman L., 54, 77, 104

Kendall, Maurice G., 4-5, 175, 176–79, 181,

193, 316, 317, 488, 540 Kennedy, W. A., 191 Kenyon, Gerald S., 382 Kimmel, Herbert D., 289 Klausmeier, Herbert J., 399 Kruskal, Wilham H., 130, 178, 290

Kuhlmann, F., 294

LaForge, R., 283

Lana, Robert D, 370, 470

Lancaster, H. O, 167

Lerner, Daniel, 186

Lev, Joseph, 107, 169

Levene, H., 372-75

Levene, H., 372-75

Levene, H., 712-75

Levene, H., 712-75

Levene, H., 714-75

Levene, H., 714-75

Levene, H., 715-75

Levene, H., 716-75

Levene, H., 716-75

Levene, H., 716-75

Levene, H., 716-75

Levene, Laretha, 194

Lord, Frederic M., 12, 181, 307, 320, 509

Lubu, Ardie, 370, 410, 411, 470

Lumsdane, A., 508, 509

Marshall, J. C., 383
Massey, Frank J., 143
Merc, Chevalier de, 96
Merrill, Maud A., 104, 191
Messick, Samuel, 183
Misholland, John F., 283
Miller, H., 290
Miller, Rubert G., 397
Millman, J. 3800, 472
Nefger, M. D., 283

Neidt, Charles O., 171 Newman, D., 382, 388 Norris, Raymond C., 283, 310 North, George E., 335 Novick, Melvin R., 500 Nunnally, J. C., 290

Ohnmacht, Fred W., 272

	To contact man
Applied statistics, 4	Correlation:
Area probability as, 218-20	causation and, 121-23
"Argument for Divine Providence, Taken from	homogeneous, 466
the Constant Regularity Observed in	multiple, 186-91
the Constant Regularity Control of	part, 182-86
the Births of Both Sexes, An" (Arbuth-	partial, 182-86
not), 272	Constant and Scients
Arithmetic mean, 60, 65, 341	Correlation coefficient:
Ars Conjectandi (Bernaulli), 96, 207	inferential techniques, 315-21
Assumption of homoscedasticity, 358	biserial, 163-64, 318, 319-20
Asymmetrical frequency polygons, 70	Kendall's tau, 316-17
	partial, 321
Average deviation, 85	phi, 315
	point-biserial, 163-64, 318
Bar graphs, 46, 47	
Bartlett's test, 506	population product-moment, 308-10
Bernoulli trials, 207	rank-biserial, 320
Bimodal scores, 58	Spearman's rank-, 172-75, 176, 316
Binomial distribution, 207~12	tetrachoric, 166, 318-19
	interpreting, 121-27
Binomial trials, 207	multiple, 186-91
Biometrika Tables for Statisticians (eds.	
Pearson and Hartley), 372, 374, 522-28,	Pearson product-moment, 109-13, 122, 155,
538, 548-51, 553-54	158, 159, 161, 163, 166, 186
Biserial correlation coefficient, 319-20	standard error of, 249
point-, 163-64, 318	Correlation ratio, 150-51
rank-, 320	Covatiance, 112
Bivariate frequency distribution, 105	analysis of, 499-501
Bivariate normal distribution, 105-7, 140, 141,	Covarying, 490
274	Critical region, 279-83
Bivariate normal population, 264	Cumulative frequency distribution, 34-35
Blocking, 490, 491-95	Curves:
on a nominal-scale, 493	leptokurtic, 91, 98
•	mesokurtic, 91, 98
California Test of Mental Maturity, 242	normal, 98-104
Cartesian coordinate system, 133-54	family of, 100-101
Causation, correlation and, 121-23	kurtosis of, 98
Cell frequencies:	skewness of, 98
disproportional, 439-43	unit, 98, 100
proportional, 432–39	uses of, 102-4
Central Limit Theorem, 229, 244-45	percentile, 45, 47
Central tendency, measurement of, 57-74	for representing a series of distributions,
choice of, 68-71	48-50
geometric mean, 7t	platykurtic, 91, 98
harmonic mean, 71-72	
the mean, 60-67	smooth, 44-45
the mean, co-or	Curvilinearity, 124-25
mean of ratios, 72	
the median, 59-60, 65-67	Date:
the mode, 58-59, 65-67	graphing, 38-42
ratio of means, 72	layout of:
Charting Statistics (Spear), 40-42	one-factor analysis, 338-39
Children's Manifest Anxiety Scale, 273, 274, 279	two-factor analysis, 400-402
Chi-square distributions, 229-32, 236-38	a model for:
noncentral, 425-26	
Circle graphs, 47	one-factor analysis, 339-42
Coefficient:	two-factor analysis, 402-3
confidence, 260	symbolizing, 17-18, 400-402
mb, 150 to 210	tabulation of, 26-32
phi, 158-59, 315	form of the table, 31-32
Column diagrams, 43-44	frequency distribution, 27-31
Combinations, 203-6	rank order, 26-27
Confidence coefficient, 260	Deciles, 33
Confidence interval, 260-68, 333-34	
around contrasts by the Symethod 202 ne	
around contrasts by the T-method 195.98	
Contingency table, 160	for sources of variation, 475-76
independence of classifications in, 329-33	two-factor, 416-18
Continuity correction, 332	Density function, probability, 219
Continuous random variables, 217-18	Dependent samples, 297-300, 306-7, 313-14,
Contraharmonic mean, 72	326-28
mean, /2	Dependent variables, 136

	SUBJECT INDEX 59
Descriptive statistics, 2	
Design and Analysis of Experiments in Psycho	Estimation (cont.)
ogy and Education (Lindquist), 509	
Deviation:	samples, 240–42
average, 85	sampling distribution concept, 243-50
mean, 79, 85-86	Y from X, 135-40
score, 64	Estimators, 242
standard, 82-83	properties of, 250-56 Events, 197
unit normal, 102	elementary, 198
Diagrams, column, 43-44	Exclusive range, 76-77
Digit Symbol subtest, 150-52	Expectations, 220-23
Directional alternative hypotheses, 288-89	of mean squares, 479-81
Discrete random variables, 217-18	of MS, and MS, 347-50
Disordinal interaction, 410-11	Experimental design, 486-509
Distribution:	analysis of covariance, 499-501
binomial, 207-12	blocking, 490, 491-95
chi-square, 229-32, 236-38	on a nominal-scale, 493
noncentral, 425-26	comparative, 501-8
F-, 233-34, 236-38	controlled experiments, 496-98
frequency:	example of, 488-90
bivariate, 105 cumulative, 34-35	factorial, 490
grouped, 28-31, 37	natural experiments, 496-98
representing graphically, 42-50	ordered levels of factors, 495
tabulation of data, 27-31	quasi-, 501
marginal, shapes of, 124-25	random selection of factor levels, 495-96 Experiments on Mass Communication (Hov-
mean squares, 424-28	land, Lumsdaine, and Sheffield), 508
normal, 95-108, 228-29, 236-38	
bivarrate, 105-7, 140, 141, 274	Factorial design, 490
history of, 95-97	Factor-level combinations, 487
the normal curve, 98-104	Factors, 338
as a standard, 102	crossed, 473 fixed, 473-74
probability, 218-19	nested, 473
sampling, 243-50	ordered levels of, 495
in statistical inference, 228-39 Student's 1-, 234-38, 293-94, 296, 298, 303, 318	random, 473-74
Duncan's New Multiple Range Test, 382-83	random selection of levels, 495-96
Duncan's New Manapie reange sess, 502 05	F-distributions, 233-34, 236-38
Economic statistics, 4	Fisher's Z-transformation, 265-68, 308-10, 311
Educational statistics, 4	Frequency, cell:
Elementary events, 198	disproportional, 439-43
Elementary Science Study (ESS), 463, 465	proportional, 432-39 Frequency distribution:
Equiprobability, 213	bivariate, 105
Error:	cumulative, 34-35
of the correlation coefficient, 249	grouped, 28-29, 37
of estimate, 137-38, 139 standard, 141-44	constructing, 29-31
of the mean, 248	representing graphically, 42-50
sampling, 275, 490	tabulation of data, 27-31
type I, 279, 283	Frequency polygons, 44
type II, 283-88	asymmetrical, 70
variance, 254-56	F-test, 353-57, 366 power of, 357, 376-77
Essay on Criticism (Pope), 6	Function, 215-16
Estimate, 242	
error of, 137-38, 139	Geometric mean, 71
standard, 141-44	Grand mean, 342
interval, 256 least-squares, 342, 403-6	Graphs, 38-42 bar, 46, 47
point, 256	circle, 47
of terms in the model, 342-43	of a frequency distribution, 42-50
Estimation, 133-54, 240-70	misleading, 50-52
interval, 256-68	nicture, 4/
hypothesis testing and, 333-34	suggestions for constructing, 52-54
least-squares, 403-6	typewriter, 46, 47
point, 256 population, 240–42	Handbook of Research on Teaching (ed Gage),
properties of estimators, 250-56	508, 509

Harmonic mean, 71-72	Inferential techniques (cont.)
defined, 72	mean of a population, 293-95
Height, 125	population proportion, 321-24
Heterogeneous variances, 371-72	variance of a population, 301-3
Histograms, 43-44, 47, 68	Interaction:
Homogeneous correlations, 466	absence of, 408-9
Homogeneous populations, 333	disordinal, 410-11
Homogeneous variances, 295, 372-74	nature of, 406-11
testing, 374-76	ordinal, 410-11
Homoscedasticity, 141-44, 295	two types of, 410-11
assumption of, 358	Interaction term, 403
How to Lie with Statistics (Huff), 52	International Encyclopedia of the Social
Hypotheses:	Sciences, 290
alternative, 284	Interquartile range, 78
directional, 288-89	Interval, confidence, 260-68, 333-34
null, 284, 341, 502-4	around contrasts by the S-method, 393-95
expected values of mean squares and, 423	around contrasts by the T-method, 385-88
F-test of, 353-57, 366	simultaneous, 385, 387, 394
statement of, 411-14	Interval estimate, 256
in terms of population means, 345 tests of, 428-31	Interval estimation, 256-68
See also names of hypotheses	hypothesis testing and, 333-34
Hypotheses testing, 271-91	interval measurement, 10-11 156-57
critical region, 279-83	interval-scale, leveling on 491.05
directional alternatives, 288-89	IQ, 13, 16, 51, 76, 103-4, 110, 196, 260-61,
interval estimation and, 333-34	294, 312, 335, 483-84
level of significance, 279-83	
nonuncctional atternatives 200 on	Journal of Special Education, 509
	Kendall's tau, 176-79, 316-17
one-tailed, 288-89	Kolmogorov-Smirnov test, 506
power, 283-88	Kuhimaan Anderson test, 506
scientific, 272-74	Kuhimann-Anderson intelligence test, 182, 184 Kurtosis, 90-92
significance, 481-82	of the normal curve, 98
statistical, 272-79	
two-tailed, 288-89	Law of large numbers, 202
type I error, 279-83	10-451-3QUATES, 138, 149_40 470 71
type II error, 283-88	
Inchesses	Leptokurtic curves, 91, 98
Inclusive range, 76-77	Leveling, 490, 497
Independent samples, 295-97, 303-6, 311-13, 324-26	on interval-scale, 494-95
Independent variables, 124	on ratio-scale, 494_95
Inference, 212	variable, 494-95
statistical:	Level of significance, 279-83
estimation 240 70	Levene's test, 506
AVDOthesis teering 274 Av	
	lower real, 76 upper real, 76
Inferential statistics, 2-3	Linear regression
	Linear regression equation, 187 Linear relationships, 118, 124
	Linear transformation, 119-20
	Lorge-Thorndike Intelligence Test, 335
Kendall's tau, 176-79, 316-17 partial, 321	Lorge-Thorndike Picture Reasoning Test, 335
phi, 315	Test, 335
Doint-hiserial 173 co	Manifest Anxiety Scale, 123
population product-moment, 308-10 rank-biserial, 320	
rank-biserial, 320	Mathematical Association of America, 332
Spearman's rank tra	Mathematical statistics, 4
tetrachoric, 166, 318-19	
dependent samples, 297-200, 204 -	arithmetic, 60, 65, 341
dependent samples, 297-300, 306-7, 313-14,	
	of combined groups, 65-66 contraharmonic, 72
	defined, 60
tingency table, 329-33	deviation 70 es oc
independent samples, 295-97, 303-6, 311-	geometric, 71
interval estimation, 333-34	
	harmonic 71 70
	defined, 72

Mean (cont.)	Measurement (cont.)
interpretation of, 67	standard deviation, 82-83
of a population, 293-95	standard scores, 86-88
null hypothesis in terms of, 345	the variance, 79-82, 83-85
properties of, 64-65	Measurement in Today's Schools (Stanley), 70,
of ratios, 72	266
sample, 60	Measuring Intelligence (Terman and Merril), 104
standard error of, 248	Median, 59-60, 65-67
unweighted analysis, 440	calculation of, 59-60
variance error of, 248	of combined groups, 65-66
Mean square (MS), 347, 365	defined, 59
determining the expectations of, 479-81	interpretation of, 67
distribution of, 424-28	Median interval, 60
expected values of, 421-24	Mesokurtic curves, 91, 98
null hypotheses and, 423	Metropolitan Achievement Tests, 399, 453
two-factor, 418	Metropolitan Reading Test, 459-60
Measurement, 7-16	Miller Analogies Test, 335
of central tendency, 57-74	Million Random Digits with 100,000 Normal
choice of, 68-71	Deciates, A, 510-12
geometric mean, 71 harmonic mean, 71–72	Minnesota Teacher Attitude Inventory, 334,
the mean, 60-67	335
the median, 59-60, 63-67	Minnesota Tests of Creative Thinking, 335 Mode, 58-59, 65-67
the mode, 58-59, 65-67	
ratio of means, 72	of combined groups, 65-66 conventional use of, 58-59
defined, 7	interpretation of, 67
nonlinear relationships between variables,	major, 59
150-52	minor, 59
of relationships, 109-32, 155-94	Modern Language Association Foreign
computational formula for ray, 113-14	Language Proficiency Tests, 189
differences of variables, 127-29	Moments, 223-24
effect on reg of transforming scores, 119-	
20	the S-method, 388-93
illustration of the calculation of ray, 114-	compared to the T-method, 395-97
16	confidence intervals around contrasts by,
interpretation and use of r <sub>s</sub> , 175-76	193-95
interpreting correlation coefficients, 121-	the T-method, 383-84
27	compared to the S-method, 395-97
interval (ratio), 156-57	confidence intervals around contrasts by, 385-88
Kendall's tau, 176-79	Multiple correlation coefficient, 186-91
multiple correlation, 186-91	Multiple prediction, 186-91
multiple prediction, 186-91 nominal-dichotomous, 156	Multiple-prediction equation, 187
ordinal, 8-10, 156	Multiple-regression equation, 187
part correlation, 182-86	Multiplicative rule of probabilities, 202-3
partial correlation, 182-86	Multivariate prediction, 187
Pearson product-moment correlation	*
coefficient, 109-13, 122, 155, 158, 159,	National Industrial Conference Board, 39
161, 163, 166, 186	Newman-Keuls procedure, 382, 388
properties of \$\phi\$, 161-62	Neyman-Pearson hypothesis-testing theory,
range of values of res, 117-19	287
Spearman rank-correlation coemcient,	Nominal-dichotomous measurement, 156 Nominal measurement, 8
172-75, 176, 316	Noncentral chi-square distribution, 425-26
tied ranks problem, 175	Nondirectional alternative hypotheses, 288-
variance of sums, 127-29	89
scales of, 8-14 characteristics, 12	Nonlinear relationships between variables,
examples of, 12	150-52
interval, 10-11	Normal curves:
nominal, 8	distribution, 98-104
ordinal, 8-10, 156	family of, 100-101
ratio, 11-14	kurtosis of, 98
of variability, 14-16, 75-94	skewness of, 98 unit, 98, 100
kurtosis, 90–92	uses of, 102-4
mean deviation, 79, 83-86	Normal distribution, 95-108, 228-29, 236-38
the range, 76-78	bivariate, 105-7, 140, 141, 274
skewnese, 88-90	

Normal distribution (cont.)	Probability, 195-227
history of, 95-97	as an area, 218-20
the normal curve, 98-104	binomial distribution, 207-12
as a standard, 102	combinations, 203-6
Notation, sigma, 18-24	combining, 198-203
Null hypotheses, 284, 341, 502-4	first addition rule, 198-99
expected values of mean squares and,	multiplicative rule, 202-3
423	second addition rule, 200-202
F-test of, 353-57, 366	defined, 197
statement of, 411-14	density function, 219
in terms of population means, 345	distributions, 218-19
tests of, 428-31	equi-, 213
	expectations, 220-23
Ogive, 45	as a mathematical system, 196-98
One-tailed test, 288-89	moments, 223-24
One-tailed t-test, 295	permutations, 203-6
Operations, symbolizing, 17–18	randomness, 212-15
Ordinal interaction, 410-11	random sampling, 212-15
Ordinal measurement, 8-10, 156	random variable, 215-17
	types of, 217-18
Parameters, 240-42	Product-moment biserial, 163
Part correlation, 182-86	Product-moment correlation coefficient, 308-10
Partial correlation, 182-86	Psychological Bulletin, 333
Pascal's triangle, 208-9	Psychological statistics, 4
Pearson product-moment correlation coeffi-	
cient, 109-13, 122, 155, 158, 159, 161,	Quantiles, 33
163, 166, 186	Quarterly Publications of the American Statis-
Percentile curves, 45, 47	tical Association, 53
for representing a series of distributions, 48-50	Quartiles, 33
Percentile range, 77	Quasi-experimentation, 501
Percentiles, 33	DANID Comments and
defined, 34	RAND Corporation, 510
determining, 34-38	Randomized-block design, 491-92 Randomness, 212-15
Permutations, 203-6	Random sampling, 212-15, 242-43
φ, properties of, 161-62	estimation, 242-43
Phi coefficient, 158-59, 315	Random variable, 215-17
Picture graphs, 47	continuous, 217-18
Planning of Experiments (Cox), 508	discrete, 217-18
Platykuriic curves, 91, 98	types of, 217-18
Point-biserial correlation coefficient, 163-64,	Range, 76-78
318	exclusive, 76-77
Point estimate, 256	inclusive, 76-77
Point estimation, 256	interquartile, 78
Polygons:	percentile, 77
frequency, 44	semi-interquartile, 77-78
asymmetrical, 70	defined, 78
for representing a series of distributions, 48,	
Population, 212, 240-42	Rank Correlation Methods (Kendall), 540
bivariate normal, 264	Rank order, 26-27
estimation, 240-42	Ratio, correlation, 150-51
homogeneous, 333	Ratio measurement, II-I4
mean of, 293-95	Ratio of means, 72
null hypothesis in terms of, 345	Ratio-scale, leveling on, 494-95
product-moment contributor coefficient	Region of rejection, 281 Relationships:
209-10	between be and bi. 144-49
proportion, 321–24	Incar, 118, 124
variance of, 301-3	measurement of, 109-32, 155-94
Power:	computational formula for res. 113-14
of the F-test, 357, 376-77	differences of variables, 127-29
hypotheses testing, 283-88 Prediction, 133-54	effect on ray of transforming scores, 119-2
multiple, 186-91	illustration of the calculation of rep, 114-1
multivariate, 187	interpretation and use of r., 175-76
univariate, 187	interpretation of correlation coefficients
Prediction line, 137	121-27
• • • •	interval (ratio), 156-57

Detector to the contract of th	
Relationships (cont.)	Skewness, 88-90
Kendali's tau, 176-79	negative, 90
multiple correlation, 186-91	of the normal curve, 98
multiple prediction, 186-91	positive, 90
nominal-dichotomous, 156	S-method, 388-93
ordinal, 8-10, 156	compared to the T-method, 395-97
part correlation, 182-86	confidence intervals around contrasts by,
partial correlation, 182-86	393-95
Pearson product-moment correlation	
coefficient, 109-13, 122, 155, 158, 159	, Sociological statistics, 4
161, 163, 166, 186	Spearman's rank-correlation coefficient, 172-
properties of φ, 161-62	75, 176, 316
range of values of ray, 117-19	Standard deviation, 82-83
Spearman rank-correlation coefficient	, Standard error of estimate, 141-44
172-75, 176, 316	Standard error of the correlation coefficient,
tied ranks problem, 175	249
variance of sums, 127-29	Standard error of the mean, 248
nonlinear, between variables, 150-52	Standard scores, 86-88
Repeated measures design, 469	Stanford Achievement Test, 307-8
Replication of the experiment, 424	Stanford-Binet Intelligence Test, 51, 103-4, 110
Residual change score, 182-83	Statistical analysis, units of, 505-9
r, interpretation and use of, 175-76	Statistical hypotheses, 272-79
r <sub>sy</sub> :	Statistical inference:
computational formula for, 113-14	estimation, 240–70
effect of transforming scores, 119-20	interval, 256-68
illustration of the calculation of, 114-16	population, 240-42
range of values of, 117-19	properties of estimators, 250-56
	random sampling, 242-43
Sample mean, 60	samples, 240-42
Samples, 212, 240-42	sampling distribution concept, 243-50
	hypothesis testing, 271-91
dependent, 297-300, 306-7, 313-14, 326-28	
estimation, 240-42	critical region, 279-83
independent, 295-97, 303-6, 311-13, 324-26	directional alternatives, 288-89
Sample space, 196-97, 215, 216	level of significance, 279-83
Sample variance, 342	nondirectional alternatives, 288-89
Sampling:	one-tailed, 288-89
distributions, 243-50	power, 283-88
	scientific, 272-74
error, 275, 490	statistical, 272-79
random, 212-15, 242-43	
estimation, 242-43	two-tailed, 288-89
Sampling Techniques (Cochran), 241	type I error, 279-83
Scales of measurement, 8-14	type 11 error, 283-88
characteristics, 12	theoretical distributions, 228-39
examples of, 12	Statistical Methods for Research Workers
	(Fisher and Yates), 536
interval, 10-11	Statistical Methods in Research (Johnson), 306
nominal, 8	Statistical Tables for Biological, Agricultural
ordinal, 8-10, 156	and Medical Research (Fisher and
ratio, II-I4	Yates), 520, 521
Scatter diagrams, 110, 111, 116, 123-24, 125,	
138, 141, 247	Statistics, 1-6
Scholastic Aptitude Test, 188	applied, 4
School Mathematics Study Group (SMSG),	defined, 4-5
487	descriptive, 2
Science Curriculum Improvement Study	economic, 4
(SCIS), 463, 465	educational, 4
	inferential, 2-3
Scientific hypotheses, 272-74	mathematical, 4
Score class, 28	psychological, 4
Semi-interquartile range, 77–78	sociological, 4
defined, 78	300101080444
Series, ungrouped, 26	summary, 57
Sigma notation, 18-24	Statisticulation, 1
Significance:	Straight line
level of, 279-83	general equation for, 135
testing, 481-82	intercept of, 135
Simple random sampling, 212-15	slope of, 135
Simultaneous confidence intervals, 385, 387,	Service 490 492
304	on ordinal-scale, 493-94

	Variance (cont.)
Stratifying variable, 493-94 Student's 1-distribution, 234-38, 293-94, 296,	expectations of mean squares, 475
298, 303, 318	heterogeneous, 371-72
Summary statistics, 57	homogeneous, 295, 372-74
Sums of squares, 143-45, 364-65	testing, 374-76
computing, 476-79	lines of ANOVA table, 474-73
between groups, 345	mixed-effects, 463-71
with groups, 345	model for the data, 402-3
total 343-45	one-factor analysis of, 338-80
in the two-factor ANOVA, 414-16	degrees of freedom, 345-47, 365
Systematic biases, 214	distribution theory, 350-53
-,	estimates of terms in the model, 342-43
Tables, \$10-54	expectations of MS, and MS, 347-50
the form of, 31-32	failure to meet assumptions of, 368-74
Tau, 176-79, 316-17	F-test of the null hypothesis, 353-57,
1-distributions, 234-38, 293-94, 296, 298, 303,	366
318	layout of data, 338-39
"Tests of Significance" (Kruskal), 290	mean squares, 347, 365 model for the data, 339-42
Tetrachoric correlation coefficient, 166, 318-19	with n observations per cell, 357-62
T-method, 383-84	mult hypothesis in terms of population
compared to the S-method, 395-97 confidence intervals around contrasts by,	means, 345
385-88	power of the F-test, 376-77
Transformation:	sums of squares, 343-45, 364-65
Fischer's Z-, 265-68, 308-10, 311	testing homogeneity, 374-76
Imear, 119-20	with unequal n's, 362-68
Tukey method, 388, 397-98	of a population, 301-3
Two-tailed test, 288-89	properties of, 83-85
Two-tailed 4-test, 295	random-effects, 452-62
Typewriter graphs, 46, 47	sample, 342
	significance testing, 481-82
Ungrouped series, 26	of sums, 127-29
Unit normal curves, 98, 100	two-factor analysis of, 400-451
Unit normal deviation, 102	computational procedures, 418-20
Units of statistical analysis, 505-9	degrees of freedom, 416-18
Univariate prediction, 187 Unweighted means analysis, 440	distribution of mean squares, 424-28 with equal numbers of observation per
Uses of Things test, 273	cell, 431-32
Oses of strongs test, 2-0	expected values of mean squares, 421-24
Value, absolute, 85	fixed-effects with unequal n's, 432-43
Variability, measurement of, 14-16, 75-94	hypothesis tests of the null hypotheses,
kurtosis, 90-92	428-31
mean deviation, 79, 85-86	layout of data, 400-402
the range, 76-78	least-squares estimation of the model,
skewness, 88-90	403-6
standard deviation, 82-83	mean squares, 418
standard scores, 86-88	model for data, 339-42
the variance, 79-82, 83-85 Variables:	multiple comparisons in, 443-45
dependent, 136	nature of interaction, 405-11
differences of, 127-29	sources of variation, 414
exact value of, 15-16	statement of null hypotheses, 411-14
independent, 136	sums of squares in, 414-16
leveling, 494-95	symbolization of data, 400-402 writing the ANOVA table, 471-72
nonlinear relationships between, 150-52	Variance error, 254-56
random, 215-17	of the mean, 248
continuous, 217-18	Venn diagrams, 200, 201
discrete, 217-18 types of, 217-18	
reported value of, 15-16	Wechsler Adult Intelligence Scale, 150
stratifying, 493-94	
Variance, 79-82, 83-85, 452-85	Weensier Intelligence Scale for Children
analysis of, are ANOVA	
calculation of, \$1-82	
computing sums of squares 424 70	Wide-Range Vocabulary Test, 335
degrees of freedom for sources of, 475-	Z-transformation, Fisher's, 265-68, 308-10,